

# FLORIDA INFLUENZA SURVEILLANCE

Week 31: August 2nd 2009– August 8th 2009



Kateesha McConnell, MPH, Respiratory Disease Surveillance Epidemiologist  
 Kate Goodin, MPH, Surveillance Epidemiologist  
 Lea Wansbrough, MPH, FL EIS Fellow  
 Leah Eisenstein, MPH, Surveillance Epidemiologist  
 Aaron Kite-Powell, MS, Surveillance Epidemiologist  
 Lillian Stark, PhD, MPH, MS, Bureau of Laboratories-Tampa  
 Valerie Mock, Bureau of Laboratories-Jacksonville  
 Julian Everett, Influenza Coordinator  
 Subir Goyal, MBBS, FSU Intern



## IN THIS ISSUE:


- I. Summary
- II. FSPISN Influenza-like Illness Statewide Graph
- III. Florida Pneumonia and Influenza Mortality Surveillance
- IV. FDOH Laboratory Surveillance
- V. Laboratory and Influenza-like Illness Surveillance by Region
- VI. County Influenza Activity Map
- VII. Reports of Influenza or Influenza-like Illness (ILI) Outbreaks
- VIII. Notifiable Disease Reports: Influenza-associated deaths in children & post-influenza infection encephalitis
- IX. ESSENCE ILI syndromic surveillance summary

## I. SUMMARY

Influenza surveillance in Florida consists of seven surveillance components\*: 1) Florida Sentinel Physician Influenza Surveillance Network (FSPISN); 2) Florida Pneumonia & Influenza Mortality Surveillance System; 3) Bureau of Laboratories viral surveillance; 4) County influenza activity levels; 5) Notifiable Disease Reports: Influenza-associated deaths in children, post-influenza infection encephalitis and novel influenza cases; 6) Influenza or ILI outbreaks; 7) Syndromic surveillance.

During week 31 (08/02/09-08/08/09), the proportion of patient visits for influenza-like illness (ILI) as reported by the Florida Sentinel Physician Influenza Surveillance Network was 3.88 percent. This is above the state threshold for moderate activity of 2.98 percent. Seventy-seven (53%) of 144 ILI specimens tested by Bureau of Laboratories were positive for influenza. Three counties reported widespread activity and 12 counties reported localized activity. Twenty-four counties reported sporadic activity and 3 counties reported no activity. Twenty-five counties did not report. The graph below shows the progression of the 2007-08 & 2008-09 Florida influenza seasons as monitored by three\*\* of seven surveillance systems.

Each week an activity code for the state as a whole is reported to the Centers for Disease Control and Prevention (CDC). There are five possible categories: No Activity, Sporadic, Local, Regional, or Widespread. Regional activity has been reported in Florida for this reporting week (week 31). Florida meets the CDC regional activity definition which is: Outbreaks of influenza or increases in ILI cases and recent laboratory-confirmed influenza in at least two but less than half the regions of the state with recent laboratory evidence of influenza in those regions. The CDC report can be viewed at <http://www.cdc.gov/flu/weekly/usmap.htm>.




\*The purposes of these surveillance systems are to determine when and where influenza activity is occurring, to identify circulating viruses, to detect changes in the circulating influenza viruses, to track patterns of influenza-associated morbidity and mortality and estimate the overall impact of influenza in the state of Florida.

\*\*1) FSPISN, 2) State Laboratory Viral Surveillance, and 3) County Activity Levels.

## II. FSPISN INFLUENZA-LIKE ILLNESS STATEWIDE GRAPH

During week 31 3.88%\* of patient visits to Florida sentinel providers were due to ILI. This percentage is below the statewide threshold for moderate activity of 2.98%\*\*. The percentage of visits ranged from 0.00% in the Southwest to 6.18% in the Southeast region. \*FSPISN reporting is incomplete for this week (33% of providers reported). Numbers may change as more reports are received. Data displayed is weighted to the state population.




\*\*The 2008—09 threshold for moderate activity is calculated from FSPISN data. The threshold for moderate activity is the mean percentage of patient visits for ILI during influenza weeks for the previous three seasons plus two standard deviations. Only weeks with 10% or greater of laboratory specimens testing positive are included in the calculation. Due to wide variability in regional level data, it is not appropriate to apply the state baseline to regional data.

^ There was no week 53 during the 2006-07 and 2007-08 seasons; the week 53 data point for those seasons is an average of weeks 52 and 1.

## III. FLORIDA PNEUMONIA AND INFLUENZA MORTALITY SURVEILLANCE

During week 31, the total count of pneumonia and influenza (P&I) deaths reported was below the epidemic threshold. The expected number of deaths for week 31 was 99, and there were 77 observed deaths. This indicates that there were 0 excess deaths\*\*.

### Pneumonia and Influenza Mortality for 24 Florida Counties, Counts Model




\*\*Eighteen of 24 counties reported P&I deaths to create this trend graph. Please note: Data from Duval, Hillsborough, Lake, Orange, Pinellas, and Volusia were not available. 100% participation is required for this graph to provide an accurate representation. Graph created August 14, 2009 at 2:17pm ET.










# V. LABORATORY AND INFLUENZA-LIKE ILLNESS (ILI) SURVEILLANCE BY REGION

## Influenza Surveillance Regions



The table below shows the weighted ILI activity by region as reported by Florida sentinel physicians for the 2008-09 season. The graphs below include ILI activity as reported by sentinel physicians and FDOH laboratory data.

Week 31: FSPISN Weighted ILI Activity, by Region 2008-09 Season		
REGION	# and % of sentinel sites reporting	2008-09 ILI %
Centraleast	32% (10/31)	1.37%
Centralwest	51% (18/35)	2.11%
Northcentral	100% (2/2)	2.12%
Northeast	23% (5/22)	4.33%
Northwest	36% (5/14)	2.69%
Southeast	19% (3/16)	6.18%
Southwest	23% (3/13)	0.00%




- Influenza A specimens FDOH Laboratories
- Influenza B specimens FDOH Laboratories
- % of visits for ILI, reported by sentinel providers


## VI. COUNTY INFLUENZA ACTIVITY MAP

During week 31, three counties reported widespread activity. Twelve counties reported localized activity. Twenty-four counties reported sporadic activity and three counties reported no activity. Twenty-five counties did not report. *Please note: data reported from counties reporting after the deadline are recorded but may not be included in the activity map below.*

Activity Level	County
No Report	Bay, Calhoun, Charlotte, Citrus, Columbia, DeSoto, Dixie, Franklin, Gadsden, Gilchrist, Glades, Gulf, Hendry, Hillsborough, Holmes, Jefferson, Lee, Liberty, Madison, Pasco, Santa Rosa, Sarasota, Walton, Washington
No Activity	Baker, Putnam, Sumter
Sporadic	Alachua, Bradford, Broward, Clay, Duval, Escambia, Hardee, Hernando, Highlands, Jackson, Lafayette, Lake, Leon, Levy, Manatee, Marion, Monroe, Nassau, Okeechobee, Osceola, St. Johns, Suwannee, Union, Wakulla
Localized	Brevard, Flagler, Indian River, Martin, Nassau, Okaloosa, Palm Beach, Polk, St. Lucie, Seminole, Taylor, Volusia
Widespread	Collier, Miami-Dade, Orange

### Weekly County Influenza Activity (Week ending August 8, 2009 - Week 31) County influenza activity levels are reported by county health department epidemiologists





**Florida Department of Health**  
**Bureau of Epidemiology**

**Disclaimer:**  
This product is for reference purposes only and is not to be construed as a legal document. Any reliance on the information contained herein is at the user's own risk. The Florida Department of Health and its agents assume no responsibility for any use of the information contained herein or any loss resulting there from.  
Map printed August 12, 2009 at 8:45 am ET.

## COUNTY INFLUENZA ACTIVITY LEVEL DEFINITIONS

### 0 = No Activity:

Overall clinical activity remains low with no laboratory confirmed cases<sup>†</sup> in the county.

### 1 = Sporadic:

- And/or {
- Isolated cases of laboratory confirmed influenza<sup>†</sup> in the county.
  - An ILI<sup>§</sup> outbreak in a single setting<sup>‡</sup> in the county. (No detection of decreased ILI<sup>§</sup> activity by surveillance systems\*)

### 2 = Localized:

- And/or {
- ILI<sup>§</sup> activity detected by a *single* surveillance system\* within the county. (ILI<sup>§</sup> activity has not been detected by *multiple* ILI surveillance systems.)
  - Two or more outbreaks (ILI<sup>§</sup> or lab confirmed<sup>†</sup>) detected in a *single* setting<sup>‡</sup> in the county.

### AND

- Recent (within past three weeks) laboratory evidence<sup>†</sup> of influenza activity in the county.

### 3 = Widespread:

- And/or {
- An increase in ILI<sup>§</sup> activity detected in  $\geq 2$  surveillance systems in the county.
  - Two or more outbreaks ((ILI<sup>§</sup> or laboratory confirmed<sup>†</sup>) detected in *multiple* settings<sup>‡</sup> in the county.

**No Report:** (No report was received from the county at the time of publication)

<sup>†</sup> Laboratory confirmed case = case confirmed by rapid diagnostic test, antigen detection, culture, or PCR.

<sup>§</sup>ILI = Influenza-like illness, fever  $\geq 100^{\circ}\text{F}$  AND sore throat and/or cough *in the absence* of another known cause.

\*ILI surveillance system activity can be assessed using a variety of surveillance systems including sentinel providers, school/workplace absenteeism, long-term care facility (LTCF) surveillance, correctional institution surveillance, hospital emergency department surveillance and laboratory surveillance.

<sup>‡</sup>Settings include institutional settings (LTCFs, hospitals, prisons, schools, companies) & the community.

## VII. REPORTS OF INFLUENZA OR INFLUENZA-LIKE ILLNESS (ILI) OUTBREAKS

During week 31, there were over 15 new outbreaks and/or clusters of influenza A H1N1 (swine origin) reported in summer camps and other facilities throughout the state. Please see Epi Com for the latest information.

County Health Department epidemiologists should report Influenza and ILI outbreaks via EpiCom at: <https://fdens.com/vabtrs/GateStart.aspx> within the Influenza Forum.

**Total influenza or ILI outbreaks and/or clusters reported as of week 31 (08/08/09): 59**

## VIII. NOTIFIABLE DISEASE REPORTS: INFLUENZA-ASSOCIATED DEATHS AMONG CHILDREN (<18 YEARS) & POST-INFLUENZA INFECTION ENCEPHALITIS

As of the week ending August 8, 2009, there was one new influenza-associated death among those <18 years of age and/or post influenza infection encephalitis reported in the state of Florida. This was a reported death in a 3 year old from Miami-Dade that that died on June 27, 2009 (week 25).

Deaths that occurred during week 23 and week 25 were due to novel influenza A H1N1.

Reportable Disease	# of Cases 08-09 Influenza Season
Influenza-associated deaths among those <18 years of age	4*
Post-influenza infection encephalitis	0




\*Case reported during week 6 was reclassified as a suspect case. Please note that status of reported cases are subject to change upon receipt of additional information.

Influenza-associated deaths among those <18 years of age and/or post-influenza infection encephalitis are reportable; case report forms can be accessed at: [http://www.doh.state.fl.us/disease\\_ctrl/epi/topicscrforms.htm](http://www.doh.state.fl.us/disease_ctrl/epi/topicscrforms.htm).



## IX. SYNDROMIC SURVEILLANCE SUMMARY

Florida uses the Electronic Surveillance System for the Early Notification of Community-based Epidemics (ESSENCE) for syndromic surveillance, which currently collects data from 114\* hospitals. These data are processed into 11 different syndrome categories based on the patient's chief complaint. One of the categories is influenza-like illness (ILI), which is composed of chief complaints that include the words "influenza" or "flu", or either fever and cough or sore throat. The data are collected on a daily basis from participating hospital emergency departments (ED) across the state. Displayed below are the percentage of ILI visits to local EDs from 2006 to 2009 by week.



\*The total number of facilities participating in ESSENCE has increased steadily from 2006 to 2009. In 2007 ESSENCE was implemented as the state syndromic surveillance system. Please note that numbers may change as facility data is updated.

## X. SUMMARY OF WORLDWIDE, UNITED STATES, AND FLORIDA NOVEL INFLUENZA ACTIVITY

### Avian Influenza A (H5N1)

Since the outbreak activity of avian influenza A (H5N1) began at the end of December 2003, there have been a total of 436 confirmed human cases and 262 deaths. Cases and deaths have occurred in the following nations: Azerbaijan 8 cases and 5 deaths; Bangladesh 1 case and 0 deaths; Cambodia 8 cases and 7 deaths; China 38 cases and 25 deaths; Djibouti 1 case and 0 deaths; Egypt 81 cases and 27 deaths; Indonesia 141 cases and 115 deaths; Iraq 3 cases and 2 deaths; Lao People's Democratic Republic 2 cases and 2 deaths; Myanmar 1 case and 0 deaths; Nigeria 1 case and 1 death; Pakistan 3 cases and 1 death; Thailand 25 cases and 17 deaths; Turkey 12 cases and 4 deaths; and, Vietnam 111 cases and 56 deaths. For a complete analysis and summary of WHO confirmed human cases of H5N1 from 12/1/2003 to current, please visit:

[http://www.who.int/csr/disease/avian\\_influenza/ai\\_timeline/en/index.html](http://www.who.int/csr/disease/avian_influenza/ai_timeline/en/index.html)

Changes in the WHO case definition for human infection with avian influenza H5N1 can be found here:

[http://www.who.int/csr/disease/avian\\_influenza/guidelines/case\\_definition2006\\_08\\_29/en/index.html](http://www.who.int/csr/disease/avian_influenza/guidelines/case_definition2006_08_29/en/index.html)

During weeks 26-31, no new cases of Influenza A H5N1 were reported.

During week 25, 3 new cases of Influenza A (H5N1) were reported by the Ministry of Health of Egypt. The first case is a 1-year old male from Domiat Governorate. His symptoms started on June 1, 2009. He was admitted to hospital on June 2<sup>nd</sup>. The second case is a 4-year old female from Dakhliya Governorate. Her symptoms started on June 5, 2009. She was admitted to hospital on June 6<sup>th</sup>. Both patients received oseltamivir treatment, are have recovered. The third case is a 1-year old male from Kaleen District, Kefr El Sheikh Governorate. His symptoms started on June 15<sup>th</sup>, 2009. He was admitted to Kefr El Sheikh Fever Hospital on June 16, 2009, where he received oseltamivir treatment, and is in a stable condition. Investigations indicated that all three cases had close contact with dead and/or sick poultry.

Information about previously reported cases can be found at: [http://www.who.int/csr/disease/avian\\_influenza/updates/en/index.html](http://www.who.int/csr/disease/avian_influenza/updates/en/index.html)

### Influenza A (H1N1) - (Swine Origin) -Updated 08/12/09

Worldwide, a total of 177,457 confirmed human cases and 1,462 confirmed deaths due to influenza A (H1N1) have been reported, since the virus was first recognized in April 2009. The WHO is no longer collecting data on individual cases reported from around the world. As a result of this, the number of cases reported here is most likely an underestimate of the true burden of disease.

For a summary of the most up to date H1N1 information please visit:

<http://www.who.int/csr/disease/swineflu/en/index.html>

In the United States, a total of 50 states, the District of Columbia, Guam, Puerto Rico, and the U.S. Virgin Islands have reported confirmed cases of Influenza A (H1N1) swine origin. There have been a total of 43,771 cases and 302 deaths reported as of July 24, 2009.

July 24, 2009 is the last day that CDC is providing individual confirmed and probable cases of novel H1N1 influenza. CDC will report the total number of hospitalizations and deaths each week, and continue to use its traditional surveillance systems to track the progress of the novel H1N1 flu outbreak. For more information about CDC's novel H1N1 influenza surveillance system, see [Questions & Answers About CDC's Novel H1N1 Influenza Surveillance](#).

For the latest information about this rapidly evolving situation please visit: <http://www.cdc.gov/h1n1flu/>

Because the WHO, the CDC, and the Florida Department of Health are no longer collecting representative data on all H1N1 cases that are occurring, reporting the number of new cases likely under represents the true burden of disease. For that reason, case counts will no longer be reported in this publication.

FDOH novel influenza A H1N1 surveillance reports are updated weekly and available at [http://www.doh.state.fl.us/disease\\_ctrl/epi/swineflu/Reports/reports.htm](http://www.doh.state.fl.us/disease_ctrl/epi/swineflu/Reports/reports.htm)

For more information FDOH's novel influenza A H1N1 surveillance strategies please see [Novel Influenza A H1N1 Surveillance Strategies Questions and Answers](#) at [http://www.doh.state.fl.us/disease\\_ctrl/epi/swineflu/H1N1\\_Surv\\_QA.pdf](http://www.doh.state.fl.us/disease_ctrl/epi/swineflu/H1N1_Surv_QA.pdf)



## X. SUMMARY OF WORLDWIDE, UNITED STATES, AND FLORIDA NOVEL INFLUENZA ACTIVITY

The following tables present cases reported in Merlin for the dates and times listed.  
Please note that numbers are provisional and subject to change.

536 hospitalizations\* in those with laboratory-confirmed H1N1 have been reported as of 10:00 a.m. August 12, 2009  
22 hospitalizations\* were in newly reported cases (10:00 a.m. August 5, 2009 to 10:00 a.m. August 12, 2009)

**TABLE 2. Hospitalizations\* in all reported H1N1 cases by county, Florida (n=536)**

County	Frequency	Percent
Alachua	1	0.19
Baker	1	0.19
Bay	1	0.19
Brevard	1	0.19
Broward	60	11.19
Charlotte	2	0.37
Collier	3	0.56
Columbia	1	0.19
Dade	247	46.08
Duval	12	2.24
Escambia	3	0.56
Flagler	1	0.19
Gadsden	1	0.19
Hendry	1	0.19
Hernando	2	0.37
Highlands	4	0.75
Hillsborough	10	1.87
Lake	3	0.56
Lee	17	3.17
Levy	1	0.19
Manatee	4	0.75
Martin	2	0.37
Monroe	4	0.75
Okaloosa	1	0.19
Okeechobee	1	0.19
Orange	63	11.75
Osceola	1	0.19
Palm Beach	48	8.96
Pinellas	8	1.49
Polk	5	0.93
Santa Rosa	1	0.19
Sarasota	4	0.75
Seminole	12	2.24
St. Johns	3	0.56
Volusia	6	1.12
Walton	1	0.19

**TABLE 3. Hospitalizations\* in newly reported H1N1 cases by county, Florida (n=22)**

County	Frequency	Percent
Broward	4	18.18
Dade	4	18.18
Hillsborough	1	4.55
Lee	1	4.55
Martin	1	4.55
Orange	3	13.64
Palm Beach	3	13.64
Pinellas	2	9.09
Seminole	1	4.55
Volusia	2	9.09

\*Under the current surveillance strategy, case reporting is only required for confirmed or probable cases of novel Influenza A H1N1 in a) patients with life-threatening illness, b) pregnant women who are hospitalized, and c) deaths.

## X. SUMMARY OF WORLDWIDE, UNITED STATES, AND FLORIDA NOVEL INFLUENZA ACTIVITY

All deaths in reported laboratory-confirmed H1N1 cases are presented in the following tables. Note that Influenza A H1N1 may not necessarily be the attributable cause of death in all cases.

48 deaths in those with laboratory-confirmed H1N1 have been reported as of 10:00 a.m. August 12, 2009  
 7 deaths were newly reported (10:00 a.m. August 5, 2009 to 10:00 a.m. August 12, 2009)

**TABLE 4. Deaths in all reported H1N1 cases by county, Florida (n=48)**

County	Frequency	Percent
Broward	6	12.50
Dade	17	35.42
Duval	1	2.08
Hernando	1	2.08
Hillsborough	4	8.33
Lee	2	4.17
Monroe	1	2.08
Orange	3	6.25
Osceola	1	2.08
Palm Beach	4	8.33
Pinellas	1	2.08
Polk	1	2.08
Sarasota	2	4.17
Seminole	2	4.17
Volusia	1	2.08
Walton	1	2.08

**TABLE 5. Newly reported H1N1 deaths by county, Florida (n=7)**

County	Frequency	Percent
Broward	1	14.29
Hillsborough	1	14.29
Dade	2	28.57
Pinellas	1	14.29
Seminole	1	14.29
Walton	1	14.29

**TABLE 6. Deaths in all reported H1N1 cases by age, Florida (n=48)**

Age	Frequency	Percent
0-4	1	2.08
5-24	5	10.42
25-49	22	45.83
50-64	16	33.33
65+	4	8.33