



Summary

Week 41: October 9-15, 2016

State influenza and influenza-like illness (ILI) activity:

- In week 41, influenza activity remains at low levels across the state.
- Emergency department (ED) and urgent care center (UCC) visits for ILI remained low, are slightly below levels observed in previous seasons at this time.
- In week 40, the preliminary estimated number of deaths due to pneumonia and influenza (P&I) decreased and is below levels seen in previous seasons at this time.
- In week 41, the majority of counties reported “mild” or no influenza activity.
- **No influenza-associated pediatric deaths were reported in week 41.**
 - No influenza-associated pediatric deaths have been reported so far this season.
 - **Annual vaccination remains the best way to protect children against influenza infection. Now is the perfect time to get your annual vaccine.**
- In week 41, no outbreaks of influenza or ILI were reported.
- **In the last two weeks, the most common influenza subtype detected at the Bureau of Public Health Laboratories (BPHL) has transitioned from influenza B Yamagata lineage to influenza A (H3).** As Florida moves into the winter months, it is not uncommon for the predominantly circulating strain of influenza to change.

National influenza activity:

- **Influenza activity continues to circulate at low levels nationally. In recent weeks, influenza and ILI activity increased but remains below the national baseline.** The Centers for Disease Control and Prevention (CDC) has received reports of influenza outbreaks in different parts of the country.
- **In recent weeks, influenza A (H3) has been the most common subtype reported to the CDC by public health laboratories across the nation.**
- For the 2016-17 season, CDC recommends use of inactivated influenza vaccines (IIV) or recombinant influenza vaccines (RIV). **Live attenuated influenza vaccines (LAIV) should not be used during the 2016-17 influenza season.** This recommendation follows poor or relatively lower effectiveness of LAIV between 2013 and 2016.
 - To learn more, please visit: http://www.cdc.gov/mmwr/volumes/65/rr/rr6505a1.htm?s_cid=rr6505a1_w.
- There is increased risk for highly pathogenic avian influenza (HPAI) H5 virus identification in birds as we enter the fall migratory season. HPAI H5 has not been identified in Florida birds and would be expected to be seen in more northerly states first, but identifications are possible. To date, only one wild duck in Alaska has tested positive for HPAI H5 since November 2015. No human HPAI infections have been identified in Florida or other states.
 - To learn more about HPAI, please visit: www.floridahealth.gov/novelflu.

Weekly State Influenza Activity

Sporadic

For more information see page 2 ▶


A (H3)

For more information see page 6 ▶

Predominately Circulating Strain

Influenza and ILI Outbreaks Reported as of 10/15/2016


Outbreaks



For more information see page 5 ▶

County Influenza Activity

County Activity (N)



For more information see page 4 ▶

**Table of Contents
on the next page ▶**

Table of Contents

In This Issue 2

Statewide ILI Visits 2

Statewide ILI Outpatient Visits and P&I Deaths 3

County influenza and ILI Activity Maps 4

Influenza-Associated Pediatric Deaths 4

ILI Activity and Outbreaks by Setting 5

Respiratory Virus Surveillance 6, 11

Regional ILI Visits 7

Age Groups: ILI Visits and P&I Deaths 8

At-Risk Populations: ILI Visits 9

ILI Activity by Population and Setting Type 10


Respiratory Syncytial Virus Surveillance 11

Other Respiratory Virus Surveillance 12

Florida ILI Surveillance System Summary 13

Weekly State Influenza Activity Reporting

Below is the state influenza activity level reported to CDC each week since the 2012-13 influenza season. **Florida reported sporadic influenza activity for week 41.**



The graphic above shows how influenza activity in Florida can vary widely from season to season. This unpredictability underscores the importance of influenza surveillance in Florida.

Influenza surveillance goals:

- Influenza surveillance is conducted to detect changes in the influenza virus. These data are used to help determine the annual national vaccine composition and to prepare for potential epidemics or pandemics.
- Surveillance is also conducted to identify unusually severe presentations of influenza infection, detect outbreaks, and determine seasonal influenza trends in order to guide influenza prevention, particularly in high-risk populations like children, adults ≥65 years old, and pregnant women.
- See the back page of this report for more information on influenza surveillance systems used in Florida: Page 13 ►

Statewide ILI Visits

Influenza-like illness (ILI) is defined as a fever ≥100°F AND sore throat and/or cough *in the absence* of another known cause.

ED and UCC Visits for ILI by Flu Season

ED = emergency department, UCC = urgent care center, ILI = influenza-like illness




Figure 1 shows the percent of visits for ILI from ED and UCC chief complaint data for ESSENCE-FL participating facilities (n=274), week 40, 2013 to week 41, 2016.

In week 41, the percent of visits to EDs and UCCs for ILI decreased and is slightly below levels seen in previous seasons at this time.




Visits for ILI to Outpatient Providers by Flu Season

ILI = influenza-like illness

Figure 2 shows the percent of visits for ILI reported by ILINet outpatient providers statewide (n=43), week 40, 2013 to week 41, 2016.

In week 41, the percent of visits for ILI reported by ILINet outpatient providers decreased and is below levels seen in previous seasons at this time.



P&I Deaths* from Vital Statistics by Flu Season

P&I = pneumonia and influenza




Figure 3 shows P&I deaths* for all Florida counties from the Bureau of Vital Statistics, as reported into ESSENCE-FL, week 40, 2013 to week 40, 2016.

As of week 40 (ending October 8, 2016), 112 P&I deaths have been reported in the 2016-17 influenza season.

The number of P&I deaths decreased notably and is below levels seen in previous seasons at this time.

P&I Deaths*, Multi-Year Regression Model


P&I = pneumonia and influenza

Figure 4 shows the number of preliminary estimated P&I deaths* for all Florida counties, the number of deaths predicted using a multi-year regression model, and the upper bound of the 95% confidence interval for this prediction.

For week 40 (ending October 8, 2016):

144 preliminary estimated P&I deaths were reported.

The upper bound of the 95% confidence interval for prediction is 195 deaths, with no excess deaths.




* Current season P&I death counts are preliminary estimates, and may change as more data are received. The most recent data available are displayed here. Vital statistics death records received in ESSENCE-FL are considered to be complete through week 40, 2016.

County influenza activity data are reported by county health departments through EpiGateway on a weekly basis. Information is used to determine county activity and includes laboratory results, outbreak reports, and ILI activity. **The figures below reflect a county health department's assessment of influenza activity within their county. For week 41, 15 counties reported "increasing" activity, 48 counties reported activity at a "plateau," and four counties reported "decreasing" activity.**

County Influenza Activity


Map 1

County Influenza Activity Level for Week 20
Reported by 9:30 a.m. October 19, 2016



Map 2

County Influenza Activity Trend for Week 20
Reported by 9:30 a.m. October 19, 2016



As of 9:30 a.m. October 19, 2016, a total of 67 (100%) counties reported their weekly level of influenza activity. Please note that data reported after the deadline Tuesday at 5 p.m. are recorded but may not be included in the activity maps for this week.

Influenza-Associated Pediatric Deaths

Influenza-Associated Pediatric Deaths

FIGURES 5 - 7

Figures 5-7 show the number of pediatric deaths associated with influenza infection, week 40, 2012 to week 41, 2016.

No influenza-associated pediatric deaths were reported in week 41. No influenza-associated pediatric deaths have been reported so far this season.

While rare, Florida receives reports of influenza-associated pediatric deaths each season. **Most deaths occurred in unvaccinated children with underlying health conditions.**

Children, especially those with underlying health conditions, are at higher risk of severe outcomes from influenza infection. **Annual vaccination remains the best way to protect against the flu. Now is the perfect time to get vaccinated. CDC recommends vaccination as long as influenza viruses are circulating.** To learn more, please visit: www.cdc.gov/flu/protect/whoshouldvax.htm#annual-vaccination.

Figure 5: Influenza-Associated Pediatric Deaths by Vaccination Status




Figure 6: Influenza-Associated Pediatric Deaths by Medical History





Figure 7: Influenza-Associated Pediatric Deaths by Strain Type



Reported Influenza and ILI Outbreaks

ILI = influenza-like illness

The map to the right shows influenza and ILI outbreaks by county for week 41, 2016. **In week 41 (ending October 15, 2016), no outbreaks of influenza or ILI were reported.** No outbreaks of influenza and ILI have been reported into EpiCom or Merlin in the first two weeks of the 2016-17 season.

Map 3

Influenza and ILI Outbreaks by County
Week 40, 2016 through Week 41, 2016

Outbreaks




TABLE 1: Summary of Florida Influenza and ILI Outbreaks by Setting, Week 40 through Week 41, 2016

| Setting | Total | A (H3) | A 2009 (H1N1) | A Unsubtyped | A & B Unsubtyped | B Yamagata | B Victoria | B Unsubtyped | Influenza Unspecified | Other respiratory viruses | Currently unknown pathogen |
|---|----------|----------|---------------|--------------|------------------|------------|------------|--------------|-----------------------|---------------------------|----------------------------|
| Schools | - | - | - | - | - | - | - | - | - | - | - |
| Daycares | - | - | - | - | - | - | - | - | - | - | - |
| Jails & prisons | - | - | - | - | - | - | - | - | - | - | - |
| Mental health facilities | - | - | - | - | - | - | - | - | - | - | - |
| Nursing homes & long term care facilities | - | - | - | - | - | - | - | - | - | - | - |
| Health care facilities | - | - | - | - | - | - | - | - | - | - | - |
| Other | - | - | - | - | - | - | - | - | - | - | - |
| Total | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

Reported Influenza and ILI Outbreaks by Facility Type

ILI = influenza-like illness





Figure 8 shows the distribution of outbreaks by facility type and season.

In week 41, no outbreaks of influenza or ILI were reported.

In Florida, influenza activity often increases in children and then moves through other age groups. As such, it is expected that early season outbreaks will occur in facilities serving children, such as schools and daycares.

BPHL Viral Influenza Specimen Testing

BPHL = Bureau of Public Health Laboratories



Figures 9 and 10 use BPHL viral surveillance data.

Figure 9 shows the number of influenza-positive specimens, tested by subtype and lab event date*.

In the last two weeks, the most common influenza subtype detected at BPHL has transitioned from influenza B Yamagata lineage to influenza A (H3). As Florida moves into the winter months, it is not uncommon for the predominantly circulating strain of influenza to change.




Figure 10 shows the number of specimens tested by BPHL and the percent that were positive for influenza by lab event date*.

In recent weeks, the number of specimens tested for influenza and the percent of laboratory results testing positive for influenza has remained somewhat stable. Both indicators are similar to levels seen in previous seasons at this time.

**TABLE 2: Bureau of Public Health Laboratories (BPHL) Viral Surveillance by Lab Event Date*
Reported by 10:00 a.m. October 19, 2016**

| Influenza Type | Current Week 41 | Previous Week 40 | Current 2016-17 Season |
|---|-----------------|------------------|------------------------|
| Total Specimens Tested | 3 | 9 | 12 |
| Influenza positive specimens (% of total specimen tested) | 1 (33%) | 4 (44%) | 5 (42%) |
| Influenza A 2009 (H1N1) (% of influenza positives) | - | - | - |
| Influenza A (H3) (% of influenza positives) | - | 4 (100%) | 4 (80%) |
| Influenza A not yet subtyped (% of influenza positives) | - | - | - |
| Influenza A inconclusive** (% of influenza positives) | - | - | - |
| Influenza B Yamagata (% of influenza positives) | 1 (100%) | - | 1 (20%) |
| Influenza B Victoria (% of influenza positives) | - | - | - |
| Influenza B not yet subtyped (% of influenza positives) | - | - | - |

*"Lab event date" is defined as the earliest of the following dates associated with influenza testing at the laboratory: date specimen collected, date received by the laboratory, date reported or date inserted.

**Influenza A inconclusive test results are due to technical difficulties including an insufficient sample for testing or internal sample control failure and occur occasionally in routine laboratory testing.

ED and UCC Visits for ILI by Region

ED = emergency department, UCC = urgent care center, ILI = influenza-like illness

Figures 11-17 show the percent of visits for ILI from ED and UCC chief complaints for ESSENCE-FL participating facilities (n=274), by ESSENCE-FL Regional Domestic Security Task Force (RDSTF) regions (see map 4) from week 40, 2013 to week 41, 2016*. In week 41 (the second week of the 2016-17 flu season), the percent of ED and UCC visits for ILI remains low. In regions 1 and 6, slight increases occurred and ILI activity decreased in all other regions. The percent of ED and UCC visits for ILI is similar to or below levels seen in previous seasons in all regions at this time.

2016-17 2015-16 2014-15 2013-14

FIGURE 11: Region 1 (20 facilities)




FIGURE 12: Region 2 (10 facilities)




FIGURE 13: Region 3 (31 facilities)




FIGURE 14: Region 4 (49 facilities)




FIGURE 15: Region 5 (71 facilities)




FIGURE 16: Region 6 (30 facilities)






FIGURE 17: Region 7 (63 facilities)



MAP 4

Emergency Departments (EDs) and Urgent Care Centers (UCCs) Reporting Data to ESSENCE-FL by Regional Domestic Security Task Force (RDSTF), October 19, 2016 (n=274)



*There is no week 53 for the 2012-13, 2013-14, and 2015-16 seasons; the week 53 data point for those seasons is an average of weeks 52 and 1.


0 to 4 years old 5 to 24 years old 25 to 64 years old ≥65 years old

ED and UCC Visits for ILI by Age Group

ED = emergency department, UCC = urgent care center, ILI = influenza-like illness

Figure 18 shows the percent of visits for ILI from ED and UCC chief complaints by age group for ESSENCE-FL participating facilities (n=274), week 40, 2013 to week 41, 2016.

In week 41, ED and UCC visits for ILI decreased or remained the same in all age groups. Levels are similar to those seen in previous seasons in the 5-24, 25-64 and ≥65 age groups at this time. Levels are below those seen in previous seasons in the 0-4 age group at this time.



Visits to Outpatient Providers for ILI by Age Group*

ILI = influenza-like illness




Figure 19 shows the number of visits for ILI reported by ILINet outpatient providers statewide (n=43) by age group, week 40, 2013 to week 41, 2016.

In week 41, the number of visits for ILI increased slightly in the 5-24 and 25-64 age groups and decreased in in the 0-4 and ≥65 age groups. Levels are below those seen in previous seasons in all age groups at this time.


*Data presented here are counts, not proportions. This is because age group denominator data is not available through ILINet.

P&I Deaths* from Vital Statistics by Age Group

P&I = pneumonia and influenza

Figure 20 shows P&I deaths* for all Florida counties by age group, as reported into ESSENCE-FL, week 40, 2013 to week 40, 2016.

As of week 40 (ending October 8, 2016), the number of P&I deaths decreased or remained the same in all age groups. Levels are below those seen in previous seasons in all age groups at this time.



*Current season P&I death numbers are preliminary estimates, and may change as more data are received. The most recent data available are displayed here. Vital statistics death records received in ESSENCE-FL are currently considered to be complete through week 40, 2016.

ESSENCE-FL collects data daily from 274 EDs and UCCs. Data are processed into 11 different syndrome categories based on the patient's chief complaint. One of the categories is ILI, which is composed of chief complaints that include the words "influenza" or "flu," or complaints that contain "fever," "cough," and/or "sore throat." The Florida Department of Health uses ED and UCC chief complaint data to monitor influenza and ILI activity in a timely manner in groups at higher risk of severe health outcomes (such as hospitalization and death) from influenza infection. These at-risk groups include pregnant women, children ≤ 18 years old, and adults ≥ 65 years old.

— 2016-17
 — 2015-16
 — 2014-15
 — 2013-14


ED and UCC Visits for ILI by Pregnant Women

ED = emergency department, UCC = urgent care center, ILI = influenza-like illness

Pregnant women are at high risk for severe complications due to influenza infection.

Figure 21 shows the number of visits* to EDs and UCCs with chief complaints of influenza infection and pregnancy, as reported into ESSENCE-FL, week 40, 2013 to week 41, 2016.

In week 41, the number of visits to EDs and UCCs by pregnant women with mention of influenza decreased and is below levels seen in previous seasons at this time.



ED and UCC Visits for ILI by Children ≤ 18 Years Old

ED = emergency department, UCC = urgent care center, ILI = influenza-like illness




Figure 22 shows the percent of ILI visits among all ED and UCC visits for children ≤ 18 years old, as reported into ESSENCE-FL, week 40, 2013 to week 41, 2016.


In week 41, the percent of ILI visits among all ED and UCC visits for children ≤ 18 years old decreased and is below levels seen in previous seasons at this time.

ED and UCC Visits for ILI by Adults ≥ 65 Years Old

ED = emergency department, UCC = urgent care center, ILI = influenza-like illness

Figure 23 shows the percent of ILI visits among all ED and UCC visits for adults ≥ 65 years old, as reported into ESSENCE-FL, week 40, 2013 to week 41, 2016.

In week 41, the percent of ILI visits among all ED and UCC visits for adults ≥ 65 years old decreased and is slightly below levels seen in previous seasons at this time.



ILI Activity by Setting Type

ILI = influenza-like illness

County health departments are asked to evaluate influenza activity in certain settings within their county. The assessment scale for activity ranges from no or minimal activity to very high activity.


Figure 24 shows the results of the influenza activity assessment for week 41, 2016.

Counties that reported “not applicable” for the listed settings are excluded from the denominator in the calculations below.

ILI Activity Levels:

- No or very minimal activity
- Moderate activity
- High activity
- Very high activity

FIGURE 24



Settings for Children under 18

In elementary schools, 59 counties (91%) reported no or minimal influenza or ILI activity.

In daycare settings, 51 counties (86%) reported no or minimal influenza or ILI activity.

Settings for Adults over 65

In nursing homes, 54 counties (86%) reported no or minimal influenza or ILI activity.

In retirement homes, 44 counties (86%) reported no or minimal influenza or ILI activity.

Settings for Adults ages 18 to 65

In colleges, 36 of 43 counties (84%) reported no or minimal influenza or ILI activity.

In businesses, 44 counties (85%) reported no or minimal influenza or ILI activity.

In government offices, 49 counties (86%) reported no or minimal influenza or ILI activity.

Other Unique Settings

In jails and prisons, 55 counties (89%) reported no or minimal influenza or ILI activity.

In health care settings, including rehabilitation facilities and mental health facilities, 56 counties (88%) reported no or minimal influenza or ILI activity. One county (2%) reported moderate influenza or ILI activity.

RSV Activity Summary and Seasonality

RSV = respiratory syncytial virus

State respiratory syncytial virus (RSV) activity:


- In week 41, the percent of children under the age of five diagnosed with RSV at EDs and UCCs continued to increase and remains notably above levels observed in previous seasons at this time.
- To learn more about RSV in Florida, please visit: <http://www.floridahealth.gov/rsv>.

RSV Seasonality:

- RSV activity in Florida typically peaks in the months of November through January, though activity can vary dramatically by region. According to CDC, the start of RSV season is marked by the first two consecutive weeks during which the average percentage of specimens testing positive for RSV is at or above ten percent.
- Florida has established regular RSV seasons based on these thresholds.
 - Currently all regions are considered to be in RSV season (see map 4).
- Florida's RSV season is longer than the rest of the nation and has distinct regional seasonality. For more information on RSV seasonality in Florida, see the American Academy of Pediatrics' 2015 Red Book.

Map 4

Florida Respiratory Syncytial Virus (RSV) Regional Season Breakdown



RSV surveillance goals:

- A statewide RSV surveillance system was implemented in Florida to support clinical decision-making for prophylaxis of premature infants. The determination of unique seasonal and geographic trends of RSV activity has important implications as it relates to prescribing patterns for initiating prophylaxis to children at high-risk for RSV infection.
- See the back page of this report for more information on RSV surveillance systems used in Florida: Page 13 ►

ED and UCC Visits for RSV by Children <5

ED = emergency department, UCC = urgent care center, RSV = respiratory syncytial virus




Figure 25 shows the percent of visits to EDs and UCCs with discharge diagnoses that include RSV or RSV-associated illness, as reported by participating ESSENCE-FL facilities (n=274), week 30, 2013 to week 41, 2016.

In week 41, the percent of children presenting to the ED for care with RSV increased and remains notably above levels seen in previous seasons at this time.


— 2016-17 — 2014-15
— 2015-16 — 2013-14

Laboratory RSV Surveillance

RSV = respiratory syncytial virus

Figure 26 shows the percent of laboratory results testing positive for RSV, as reported by hospital laboratories (n=11), week 40, 2013 to week 41, 2016.

In week 41, the percent of specimens testing positive for RSV increased, and is above levels observed in previous two seasons at this time.



— 2016-17 — 2014-15
— 2015-16 — 2013-14

Other Respiratory Virus Surveillance

Statewide activity:

- In week 41, the percent of specimens testing positive for rhinovirus decreased notably and is below levels observed in previous seasons at this time.
- In week 41, the percent of specimens testing positive for adenovirus increased, but remains similar to levels observed in previous seasons at this time.
- In week 41, the percent of specimens testing positive for RSV continued to increase notably and is above levels observed in the last two seasons.

Enterovirus D68 (EV-D68) activity:

- In week 41, no new cases of EV-D68 were identified in Florida.
- Eight cases of EV-D68 have been identified in Florida since February 2016. These eight cases were identified in different regions of the state and represent the full spectrum of disease. These are the first identifications of EV-D68 in the United States since the fall of 2014.
- Six of these cases were identified as a result of Florida’s participation in the Influenza Incidence Surveillance Project (IISP).
 - To learn more about EV-D68, please visit: <http://www.floridahealth.gov/diseases-and-conditions/d68>.



Outbreaks:

- In week 41, no outbreaks of RSV, parainfluenza, MPV, adenovirus, or rhinovirus were reported.

Laboratory Viral Respiratory Surveillance

Figure 27 shows the percent of laboratory results testing positive for eight common respiratory viruses, as reported by hospital laboratories (n=11), week 40, 2013 to week 41, 2016.

In recent weeks, the percent of specimens testing positive for RSV increased notably, while rhinovirus activity appears to be declining.



Non-Influenza IISP Laboratory Outpatient Surveillance*

IISP = Influenza Incidence Surveillance Project





Figure 28 shows the number of specimens testing positive for 12 common respiratory viruses, as reported by BPHL and IISP outpatient providers statewide (n=6), week 40, 2015 to week 40, 2016.

In recent weeks, the most commonly identified viruses among IISP specimens were RSV and enterovirus.



*Data presented here are counts, not proportions. The most recent data available are displayed here. IISP laboratory data are currently considered to be complete through week 40, 2016.

Florida ILINet · Data source for figures: 2 and 19

- ILINet is a nationwide surveillance system composed of sentinel providers, predominately outpatient health care providers. Florida has 88 sentinel providers enrolled in ILINet who submit weekly ILI and total visit counts, as well as submit ILI specimens to the Bureau of Public Health Labs (BPHL) for confirmatory testing.

ESSENCE-FL Syndromic Surveillance and Vital Statistics Portal · Data source for figures 1, 3-7, 11-18, 20-23, 25; map 4

- Electronic Surveillance System for the Early Notification of Community-based Epidemics (ESSENCE-FL) measures trends in ILI visits from emergency departments (ED) and urgent care clinics (UCC) and influenza mortality by using death certificates from the Bureau of Vital Statistics. Participating EDs and UCCs (n=273) electronically transmit visit data into ESSENCE-FL daily or hourly.
- For statewide and regional data on influenza-like illness, visits are counted as ED or UCC visits to participating facilities that include influenza-like illness in patient chief complaints.
- For pneumonia and influenza (P&I) surveillance, death record literals are queried using a free-text query that searches for references to P&I on death certificates. Any mention of P&I in the death certificate literals, with certain exceptions, is counted as a P&I death.
- For respiratory syncytial virus (RSV) surveillance, visits are counted as ED or UCC visits to participating facilities for which RSV or RSV-associated illness is included in the discharge diagnosis.

County Influenza Activity in EpiGateway · Data source for figures 19, 24, and maps 1 and 2

- County health department (CHD) epidemiologists report their county's influenza and ILI surveillance data weekly into the EpiGateway website. Influenza activity is classified as: No Activity, Mild, Moderate or Elevated. Setting-specific influenza activity and influenza trend information is also reported. EpiGateway data provided by CHDs creates a county-by-county breakdown of influenza and ILI activity around the state.

Outbreak Reporting in EpiCom · Data source for figure 8, map 3, and table 1

- EpiCom tracks influenza and ILI outbreak investigations by county health departments (CHDs). Reports by CHDs include the type of respiratory disease causing the outbreak and settings where outbreaks are occurring. CHD epidemiologists report outbreaks of influenza and ILI into EpiCom, Florida's online communication setting.
- Outbreaks are defined as two or more cases of influenza or ILI in a specific setting.

Outbreak Reporting in Merlin · Data source for figure 8, map 3, and table 1

- Merlin tracks influenza and ILI outbreak investigations by CHDs. Reports by CHDs include the type of respiratory disease causing the outbreak and settings where outbreaks are occurring. CHD epidemiologists report outbreaks of influenza or ILI into Merlin, Florida's reportable disease surveillance system.
- Outbreaks are defined as two or more cases of influenza or ILI in a specific setting.

Bureau of Public Health Laboratories (BPHL) · Data source for figures 9, 10 and table 2

- BPHL performs confirmatory testing and subtyping on surveillance specimens from ILINet sentinel providers, outbreak investigations, patients with severe or unusual influenza presentations and medical examiners.
- For county-specific laboratory data, please refer to the Flu Lab Report in Merlin. For instructions on how to use the Flu Lab Report, please see the Guide to Flu Lab Report on the Bureau of Epidemiology website at www.floridahealth.gov/diseases-and-conditions/influenza/_documents/flulabreportguide.pdf.

Laboratory Viral Respiratory Surveillance- Data sources for figures 26-28

- The National Respiratory and Enteric Virus Surveillance System (NREVSS) and Electronic Laboratory Reporting (ELR) collect data from laboratories in Florida on a weekly basis and monitor temporal and geographic patterns of eight commonly circulating respiratory viruses. NREVSS data is collected by the Centers for Disease Control and Prevention (CDC) and electronic laboratory report (ELR) data is collected by the Florida Department of Health (DOH).

Case-Based Influenza Surveillance

- Influenza-associated pediatric deaths (reported into Merlin, DOH's reportable disease surveillance system).
- Influenza due to novel or pandemic strains (reported into Merlin)
- Deaths in children with laboratory-confirmed influenza infection and patients with influenza infection due to novel or pandemic strains are reportable in Florida. For more information about reportable diseases please visit www.Floridahealth.gov/diseasereporting.