

# HIV/AIDS Epidemiology Partnership 13

## Citrus, Lake, Marion and Sumter Counties

Excluding Dept. of Corrections

Created: 12/03/14

Revision: 10/14/15

Florida Department of Health  
HIV/AIDS Section  
Annual data trends as of 12/31/2014  
Living (Prevalence) data as of 06/30/2015



# HIV and AIDS Case Data

- ⦿ **AIDS Cases became reportable in Florida in 1981.**
- ⦿ **HIV (not AIDS) became reportable in Florida on July 1, 1997.**
- ⦿ **HIV Infection reporting represents newly Adult HIV Infection Cases, regardless of AIDS status at time of report, that were previously reported.**
- ⦿ **AIDS cases and HIV infection cases by year of report are NOT mutually exclusive and CANNOT be added together.**
- ⦿ **Frozen databases of year-end data are generated at the end of each calendar year. These are the same data used for Florida CHARTS and all grant-related data where annual data are included.**
- ⦿ **HIV prevalence data are generated later in the year, usually in July, when most of the “expected” death data are complete.**



# HIV and AIDS Case Data (con't)

- ⦿ **Adult cases represent ages 13 and older, pediatric cases are those under the age of 13. For data by year, the age is by age of diagnosis. For living data, the age is by current age at the end of the most recent calendar year, regardless of age at diagnosis.**
- ⦿ **Unless otherwise noted, whites are non-Hispanic and blacks are non-Hispanic.**
- ⦿ **Unless otherwise noted. Area and county data will exclude DOC cases.**


# Snapshot of Persons Reported with HIV Disease, 2014, Partnership 13

	HIV Infection and AIDS Cases Reported in 2014*		
	Adults (Age 13+)	Pediatrics (Age <13)	TOTAL
HIV Infection Cases	135	-	135
AIDS Cases	67	-	67
*HIV infection cases and AIDS cases by year of report are NOT mutually exclusive and CANNOT be added together.			
Total Population, 2014*	Cumulative HIV/AIDS Cases Reported 1981-2014		
	Adults (Age 13+)	Pediatrics (Age <13)	TOTAL
905,697	HIV (not AIDS) Cases**	5	679
	AIDS Cases	21	1,890
	Total	26	2,569
*2014 estimate is provisional	**HIV (not AIDS) cases were NOT reportable until 07/1997		
Persons Living with HIV Disease through 2014, as of 06/30/2015:		2,136	



# AIDS Cases and Rates\*

## By Year of Report, 2005-2014, Partnership 13




Enhanced laboratory reporting (ELR) laws in 2006 and the expansion of ELR in 2007 led to an artificial peak in newly reported cases of AIDS in 2008. This was followed by a general decline in reported cases through 2010. Another surge in the expansion of ELR in 2012 was followed by another increase in newly reported cases of AIDS in 2013. In 2014, AIDS cases dropped by 13% from the previous year. This is lower than the 15% decline observed by the state during the same time period.

\*Source: Population estimates are provided by Florida CHARTS as of 7/9/2015. Rates are expressed as per 100,000 population.




# Adult HIV Infection Cases, by Sex and Year of Report, 2005-2014, Partnership 13



Note: Recent trends in HIV transmission are best described by the HIV case data. The relative increases in male HIV infection cases might be attributed to proportional increases in HIV transmission among men who have sex with men (MSM), which may influence future AIDS trends. \*The male-to-female ratio is the number of cases among males divided by the number of cases among females.



# Adult AIDS Cases, by Sex and Year of Report, 2005-2014, Partnership 13




Note: AIDS cases tend to represent HIV transmission that occurred many years ago. The relative increases in males cases reflect the changing face of the AIDS epidemic over time. \*The male-to-female ratio is the number of cases among males divided by the number of cases among females.




# Adult HIV Infection and AIDS Cases by Sex, Reported in 2014, Partnership 13

**HIV Infection**  
**N=135**



**AIDS**  
**N=67**




■ Males  
■ Females

Note: Partnership 13's Adult Population is: 48% Male and 52% Female.




# Adult HIV Infection Cases by Race/Ethnicity and Year of Report, 2005-2014, Partnership 13



Note: HIV case reporting reflects more recent trends in the epidemic with respect to the distribution of cases by race/ethnicity. From 2005 to 2014, the proportion of HIV infection cases among whites increased by 15 percentage points. In contrast, the proportion of HIV infection cases among Hispanics and blacks decreased by 1 and 14 percentage points, respectively, over the same time period. Other races represent less than 4% of the cases and are not included.




# Adult Male HIV Infection Cases by Race/Ethnicity and Year of Report, 2005-2014, Partnership 13



Note: From 2005 to 2014, the proportion of HIV infection cases among Hispanic and black males decreased by 3 and 17 percentage points, respectively. In contrast, the proportion of HIV infection cases among white males increased by 18 percentage points during the same time period. Other races represent less than 5% of the cases and are not included.




# Adult Female HIV Infection Cases by Race/Ethnicity and Year of Report, 2005-2014, Partnership 13



Note: From 2005 to 2014, the proportion of adult HIV infection cases among white females decreased by 14 percentage points. In contrast, the proportion of HIV infection cases among black and Hispanic females increased by 11 and 7 percentage points, respectively, during the same time period. Other races represent less than 7% of the cases and are not included.



# Adult AIDS Cases by Race/Ethnicity and Year of Report, 2005-2014, Partnership 13




- Factors Affecting Disparities**
- Late diagnosis of HIV.
  - Access to/ acceptance of care.
  - Delayed prevention messages.
  - Stigma.
  - Non-HIV STD's in the community.
  - Prevalence of injection drug use.
  - Complex matrix of factors related to socioeconomic status.

**Note:** In 2014, blacks accounted for 42% of adult AIDS cases, but only 9% of the population. Over the past ten years, the proportion of black and white cases fluctuated over time, oftentimes, crossing paths. From 2005 to 2014, the proportion of adult AIDS cases among whites and Hispanics decreased by 1 and 11 percentage points, respectively. In contrast, the proportion of AIDS cases among whites increased by 11 percentage points during the same time period. Numerous disparities can affect the increases of HIV disease in a given population. Other races represent less than 5% of the cases and are not included.



# Adult HIV Infection Case Rates\* by Sex and Race/Ethnicity, Reported in 2014, Partnership 13




Note: Among black males, the HIV case rate is nearly 3 times higher than the rate among white males. Among black females, the HIV case rate is nearly 26-fold greater than the rate among white females. Among Hispanic males and females, the HIV case rate is higher than the rate among their white counterparts.

\*Source: Population estimates are provided by Florida CHARTS as of 7/9/2015.



# Adult AIDS Case Rates\* by Sex and Race/Ethnicity, Reported in 2014, Partnership 13




Note: Among black males, the AIDS case rate is nearly 6 times higher than the rate among white males. Among black females, the AIDS case rate is 24-fold greater than the rate among white females. Hispanic male and female rates are higher than the rates among their white counterparts.

\*Source: Partnership 13 population estimates are provided by Florida CHARTS as of 7/9/2015.




# Adult HIV and AIDS Cases Reported in 2014 and Population Data, by Race/Ethnicity, Partnership 13


**HIV Infection  
N=135**



**2014 Partnership 13  
Population Estimates\*  
N=792,229**



**AIDS  
N=67**



■ White
 ■ Black
 ■ Hispanic
 ■ Other\*\*


Note: In this snapshot for 2014, blacks are over-represented among the HIV and AIDS cases, accounting for 28% of adult HIV infection cases and 42% of adult AIDS cases, but only 9% of the adult population. A group is disproportionately impacted to the extent that the percentage of cases exceeds the percentage of the population.

\*Source: Population estimates are provided by Florida CHARTS as of 7/9/2015.

\*\*Other includes Asian/Pacific Islanders, Native Alaskans/American Indians and Multi-racial individuals.




# Adult HIV Infection Cases, by Age Group at Diagnosis, and Year of Report, 2005–2014, Partnership 13



Note: From 2005 to 2014, the proportion of adult HIV infection cases among those aged 50 or older increased by 18 percentage points.



# Adult HIV Infection Cases, by Sex and Age Group at Diagnosis, Reported in 2013, Partnership 13



Note: HIV infection cases tend to reflect more recent transmission than AIDS cases, and thus present a more current picture of the epidemic. With regard to the age group with the highest percent of HIV infection cases, recent estimates show that among males, 31% of HIV infection cases occur among those aged 50 or older, similarly among females, 38% of HIV infection cases occur among those in the 50 or older age group.




# Definitions of Mode of Exposure Categories

- ◆ **MSM** = Men who have sex with men or Male-to-male sexual contact with person with HIV/AIDS or known HIV risk
- ◆ **IDU** = Injection Drug User
- ◆ **MSM/IDU** = Men who have sex with men or Male-to-male sexual contact & Injection Drug User
- ◆ **Heterosexual** = Heterosexual contact with person with HIV/AIDS or known HIV risk
- ◆ **OTHER** = includes hemophilia, transfusion, perinatal, other pediatric risks and other confirmed risks.
- ◆ **NIR** = Cases reported with No Identified Risk
- ◆ **Redistribution of NIRs** = This illustrates the effect of statistically assigning (redistributing) the NIRs to recognized exposure (risk) categories by applying the proportions of historically reclassified NIRs to the unresolved NIRs.




# Adult Male HIV Infection Cases, by Mode of Exposure and Year of Report, 2005–2014, Partnership 13



Note: NIRs redistributed. Male-to-male sexual contact (MSM) remains as the primary mode of exposure among male HIV cases in Partnership 13, followed by heterosexual contact.



# Adult Female HIV Infection Cases by Exposure Category and Year of Report, 2005-2014, Partnership 13




Note: NIRs redistributed. The heterosexual risk continues to be the dominant mode of exposure among females.





# Adult Male HIV Infection and AIDS Cases, by Mode of Exposure, Reported in 2014, Partnership 13

**HIV Infection**  
**N=111**




**AIDS**  
**N=43**





**Note:** NIRs redistributed. Among the male HIV infection and AIDS cases reported for 2014, male-to-male sexual contact (MSM) was the most common risk factor (78% and 65% respectively) followed by cases with a heterosexual risk (12% for HIV and 28% for AIDS). The recent increase among MSM is indicated by the higher MSM among HIV infection cases compared to AIDS cases, as HIV infection cases tend to represent a more recent picture of the epidemic.

# Adult Female HIV Infection and AIDS Cases, by Mode of Exposure, Reported in 2014, Partnership 13

**HIV Infection**  
**N=24**



**AIDS**  
**N=24**




**Note: NIRs redistributed. Among the female HIV infection and AIDS cases reported for 2014, heterosexual contact was the highest risk (92% and 83% respectively).**

# Cases Living with HIV Disease

Unless otherwise noted, data in the following slides represent persons living with HIV/AIDS (PLWHAs), who were living in Florida (regardless where diagnosed) through the most recent calendar year. Living data are also referred to as prevalence cases or living with HIV disease.



# Adults Living with HIV Disease By Zip Code, Diagnosed through 2014, Partnership 13




NIRs are not redistributed.  
Excludes DOC, homeless, and cases with unknown zips.  
Data as of 06/30/2015





# Injection Drug Users (IDUs)\* Living with HIV Disease By Zip Code, Diagnosed through 2014, Partnership 13

**Presumed Living  
IDU HIV/AIDS Cases**




**N=342**

NIRs are not redistributed.  
Excludes DOC, homeless, and cases with unknown zips.  
\*Includes MSM/IDU cases.  
Data as of 06/30/2015




# Adult Heterosexuals Living with HIV Disease By Zip Code, Diagnosed through 2014, Partnership 13

## Presumed Living Heterosexual HIV/AIDS Cases




**N=706**



NIRs are not redistributed.  
Excludes DOC, homeless, and cases with unknown zips.  
Data as of 06/30/2015



# Adults Living with HIV Disease By Zip Code and Race/Ethnicity, Diagnosed through 2014, Partnership 13




Total includes all races, some which are not on map.  
Excludes DOC, homeless, and cases with unknown zips.  
Data as of 06/30/2015







# Adults Living with HIV Disease, by Sex and Race/Ethnicity Diagnosed through 2014, Partnership 13

**Males**  
N=1,406




**Females**  
N=728



**Note:** Among adults living with HIV disease, white males (55%) and black females are the most affected (53%).

\*Other includes Asian/Pacific Islanders, Native Alaskans/American Indians and Multi-racial individuals.

# Case Rates\* of Adults Living with HIV Disease, by Sex and Race/Ethnicity, Diagnosed through 2014, Partnership 13



Note: Among black males living with HIV disease Diagnosed through 2014, the case rate is 4 times higher than the rate among white males. Among black females living with HIV disease, the case rate is nearly 15 times higher than the rate among white females. The Hispanic male rate is 2 times higher and the Hispanic female rate is 3 times higher than the rate among their white counterparts. Data excludes Department of Corrections cases.

\*Source: Population estimates are provided by Florida CHARTS as of 7/9/2015.



\*\*Other includes Asian/Pacific Islanders, Native Alaskans/American Indians and Multi-racial individuals.



# Adults Living with HIV Disease, by Sex and Current Age Group, Diagnosed through 2014, Partnership 13

**Males**  
N=1,406

**Females**  
N=728




Note: Males living with HIV disease have a higher proportion of cases who are currently 40 years of age or older (80%), compared with females living with HIV disease (71%).




# Adult Males Living with HIV Disease by Race/Ethnicity and Mode of Exposure Diagnosed through 2014, Partnership 13


White Non-Hispanic,  
N=770



Black Non-Hispanic,  
N=390



Hispanic,  
N=213



■ MSM ■ IDU ■ MSM/IDU ■ Heterosexual ■ Other\*


Note: NIRs redistributed. Male-to-male sexual contact (MSM) represents the highest risk for all races. White males have the smallest proportion of heterosexual contact risk.

\* Other includes hemophilia, transfusion, perinatal, other pediatric risks and other confirmed risks.




# Adult Females Living with HIV Disease by Race/Ethnicity and Mode of Exposure Diagnosed through 2014, Partnership 13


White Non-Hispanic,  
N=236



Black Non-Hispanic,  
N=382



Hispanic,  
N=90




■ IDU ■ Heterosexual ■ Other\*

Note: NIRs redistributed. Heterosexual contact is the majority risk for all races. However, whites have the largest proportion of IDU risk.

\* Other includes hemophilia, transfusion, perinatal, other pediatric risks and other confirmed risks.



# Annual Prevalence of Adults Living with HIV Disease, 1995-2014, Partnership 13



As a result of declining deaths, annual HIV/AIDS diagnoses have exceeded deaths since 1995, and the number of persons reported with HIV/AIDS who are presumed to be alive have been increasing. Since 1995, the number of persons reported living with HIV/AIDS have increased over 640%. In 2014, the prevalence increased by 7.4% since the previous year.

Note: These data represent adults living with HIV disease diagnosed in Florida regardless of their current residence.




# **Partnership 13's Top-9 Priority Populations in 2014 for Primary and Secondary HIV Prevention Based on Persons Living with HIV Disease**

- 1. White Men who have sex with Men**
- 2. Black Heterosexual men and women**
- 3. Black Men who have sex with Men**
- 4. White Heterosexual men and women**
- 5. White Injection Drug User**
- 6. Hispanic Men who have sex with Men**
- 7. Hispanic Heterosexual men and women**
- 8. Black Injection Drug User**
- 9. Hispanic Injection Drug User**

This final ranking is a result of ranking 9 race/risk groups among those newly reported in eHARS with HIV disease from the 3 most recent years, plus ranking these same 9 race/risk groups from all persons who were reported and living with HIV disease in eHARS through the most recent calendar year. The two ranks were then weighted and combined resulting in the final rank.



## Number and Percentage of Persons Diagnosed and Living with HIV (PLWH) Engaged in Selected Stages of the Continuum of HIV Care Partnership 13 (excl. DOC), 2014



- 87% of those diagnosed with HIV in 2014 had documented HIV-related care within 3 months of diagnosis
- 81% of PLWH in care had a suppressed viral load in 2014

**(1) HIV Diagnosed:** Persons diagnosed and living with HIV (PLWH) in Florida through the end of 2014.

**(2) Ever in Care:** PLWH with at least 1 documented viral load (VL) or CD4 lab, medical visit or prescription since HIV diagnosis.


**(3) In Care:** PLWH with at least 1 documented VL or CD4 lab, medical visit or prescription in 2014.

**Retained in Care:** PLWH with 2 or more documented VL or CD4 labs, medical visits or prescriptions (at least 3 months apart) in 2014.

**(4) On ART:** Estimated PLWH on antiretroviral therapy (ART) in 2014 (estimated from 2013 FL MMP data).

**(5) Suppressed Viral Load:** PLWH with a suppressed VL (<200 copies/mL) on last VL in 2014.

## Number and Percentage of Persons Diagnosed and Living with HIV (PLWH) Engaged in Selected Stages of the Continuum of HIV Care Partnership 3/13 (excl. DOC), 2014



- 84% of those diagnosed with HIV in 2014 had documented HIV-related care within 3 months of diagnosis
- 80% of PLWH in care had a suppressed viral load in 2014

(1) **HIV Diagnosed:** Persons diagnosed and living with HIV (PLWH) in Florida through the end of 2014.

(2) **Ever in Care:** PLWH with at least 1 documented viral load (VL) or CD4 lab, medical visit or prescription since HIV diagnosis.


(3) **In Care:** PLWH with at least 1 documented VL or CD4 lab, medical visit or prescription in 2014.

**Retained in Care:** PLWH with 2 or more documented VL or CD4 labs, medical visits or prescriptions (at least 3 months apart) in 2014.

(4) **On ART:** Estimated PLWH on antiretroviral therapy (ART) in 2014 (estimated from 2013 FL MMP data).

(5) **Suppressed Viral Load:** PLWH with a suppressed VL (<200 copies/mL) on last VL in 2014.

# Resident Deaths due to HIV Disease by Year of Death, 1995-2014, Partnership 13



Race/Ethnicity	2014	
	No.	rate
White Male	11	3.3
White Female	4	1.1
Black Male	9	20.5
Black Female	5	11.5
Hispanic Male	-	0.0
Hispanic Female	-	0.0
Other*	-	0.0
<b>TOTAL</b>	<b>29</b>	<b>3.2</b>

Source: Florida Department of Health, Bureau of Vital Statistics, Death Certificates (as of 05/31/2015). Population data are provided by Florida CHARTS as of 7/9/2015.

\*Other includes Asian/Pacific Islanders, Native Alaskans/American Indians and Multi-racial individuals.



## Some Useful Links

- ⌘ CDC HIV/AIDS Surveillance Reports  
(State and Metro Data):  
<http://www.cdc.gov/hiv/stats/hasrlink.htm>
- ⌘ MMWR (Special Articles on Diseases, Including HIV/AIDS):  
<http://www.cdc.gov/mmwr/>
- ⌘ U.S. Census Data (Available by State, County):  
<http://www.census.gov>
- ⌘ Partnership 13 Dept. of Health, HIV/AIDS Section  
Website (Slide sets, Facts Sheets, Monthly Surveillance  
Report, Counseling & Testing Data, etc.):  
<http://www.floridahealth.gov/diseases-and-conditions/aids/surveillance/index.html>



**“The reason for collecting, analyzing and disseminating information on a disease is to control that disease. Collection and analysis should not be allowed to consume resources if action does not follow.”**

--Foege WH et al. Int. J of Epidemiology 1976; 5:29-37



# Partnership 13 Surveillance Contact

Richard Willis

Alachua County Health Department

Phone: 352-334-7968

Email: [Richard.Willis@flhealth.gov](mailto:Richard.Willis@flhealth.gov)

Gay Koehler-Sides, HIV/AIDS Program Coordinator

Phone: 352-334-7965

Email: [Gay.Koehler-Sides@flhealth.gov](mailto:Gay.Koehler-Sides@flhealth.gov)



For Florida HIV/AIDS Surveillance Data  
**Contact: (850) 245-4444**



**Lorene Maddox, MPH**

**Ext. 2613**

**Tracina Bush, BSW**

**Ext. 2612**

**Madgene Moise, MPH**

**Ext. 2373**

**Visit Florida's internet site for:**  
**Monthly Surveillance Reports**  
**Slide Sets and Fact Sheets**  
**Annual Reports and Epi Profiles**

**<http://www.floridahealth.gov/diseases-and-conditions/aids/surveillance/index.html>**

**Visit CDC's HIV/AIDS internet site for:**  
**Surveillance Reports, fact sheets and slide sets**

**<http://www.cdc.gov/hiv/topics/surveillance/resources/reports/index.htm>**