

Citrus County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Citrus County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	6.7	(3.5 - 9.8)	3.3	(3.0 - 3.5)	+		
Smoked cigars on one or more of the past 30 days	5.1	(2.8 - 7.3)	3.9	(3.6 - 4.2)			+
Used smokeless tobacco on one or more of the past 30 days	6.2	(3.5 - 8.9)	2.3	(2.1 - 2.6)	+		
Used any form of tobacco on one or more of the past 30 days	11.2	(7.1 - 15.4)	6.4	(6.0 - 6.8)	+		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	66.6	(60.4 - 72.8)	72.0	(71.2 - 72.8)			+
Exposed to secondhand smoke during the past 7 days	59.4	(53.7 - 65.0)	41.3	(40.4 - 42.2)	+		
Smoking is allowed in the home	18.9	(14.9 - 22.8)	9.9	(9.5 - 10.4)	+		
Taught about tobacco use in any classes this year	65.3	(59.3 - 71.3)	53.4	(52.0 - 54.7)	+		
Parent or guardian has talked about the dangers of tobacco in the past year	65.6	(60.4 - 70.8)	63.9	(63.1 - 64.6)			+
Served as an active member of SWAT	2.9	(1.1 - 4.6)	4.3	(3.9 - 4.7)			+
<u>Definitely did not think</u> that smokers have more friends	28.0	(22.8 - 33.1)	33.7	(32.9 - 34.5)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	70.9	(65.5 - 76.3)	73.7	(73.0 - 74.4)			+

Citrus County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Citrus County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	20.9	(16.9 - 24.9)	20.1	(19.5 - 20.7)			+
Current asthma, among all individuals	12.8	(9.3 - 16.3)	12.0	(11.5 - 12.5)			+
Asthma attack in past year, among individuals with lifetime asthma	16.5	(7.2 - 25.9)	18.1	(16.8 - 19.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	10.5	(7.3 - 13.8)	11.6	(11.0 - 12.1)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	15.2	(11.0 - 19.3)	16.5	(15.8 - 17.2)			+
Sufficient vigorous activity	77.5	(72.7 - 82.4)	70.1	(69.2 - 70.9)	+		
Sufficient moderate activity	25.0	(20.8 - 29.3)	24.8	(24.1 - 25.4)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	44.5	(39.1 - 49.8)	44.4	(43.6 - 45.2)			+
Described themselves as slightly or very overweight	32.0	(26.9 - 37.1)	30.2	(29.4 - 30.9)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	13.2	(9.7 - 16.8)	11.4	(10.8 - 12.0)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	19.2	(15.6 - 22.8)	17.8	(17.2 - 18.5)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	7.0	(4.5 - 9.6)	7.8	(7.3 - 8.3)			+
Purposely hurt themselves without wanting to die, during the past 12 months	13.1	(9.0 - 17.2)	12.6	(12.0 - 13.2)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	19.4	(15.0 - 23.7)	19.2	(18.5 - 19.8)			+

Citrus County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Citrus County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	17.8	(13.6 - 22.0)	10.1	(9.6 - 10.6)	+		
Smoked cigars on one or more of the past 30 days	13.4	(9.0 - 17.7)	11.4	(10.9 - 11.9)			+
Used smokeless tobacco on one or more of the past 30 days	12.9	(9.0 - 16.9)	5.6	(5.2 - 6.0)	+		
Used any form of tobacco on one or more of the past 30 days	27.5	(22.4 - 32.5)	17.9	(17.3 - 18.6)	+		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	45.9	(40.5 - 51.4)	58.4	(57.6 - 59.3)		+	
Exposed to secondhand smoke during the past 7 days	65.6	(60.3 - 70.9)	47.6	(46.7 - 48.4)	+		
Smoking is allowed in the home	18.1	(14.1 - 22.0)	10.0	(9.5 - 10.5)	+		
Taught about tobacco use in any classes this year	43.1	(35.6 - 50.6)	33.4	(32.4 - 34.4)	+		
Parent or guardian has talked about the dangers of tobacco in the past year	46.3	(41.1 - 51.6)	50.4	(49.6 - 51.2)			+
Served as an active member of SWAT	4.9	(2.6 - 7.1)	4.0	(3.7 - 4.3)			+
<u>Definitely did not think</u> that smokers have more friends	26.9	(22.4 - 31.4)	30.5	(29.8 - 31.2)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	73.3	(68.6 - 78.0)	71.7	(71.0 - 72.4)			+

Citrus County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Citrus County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	22.7	(18.3 - 27.2)	20.8	(20.2 - 21.5)			+
Current asthma, among all individuals	12.1	(8.8 - 15.4)	10.6	(10.1 - 11.1)			+
Asthma attack in past year, among individuals with lifetime asthma	18.4	(10.0 - 26.7)	16.3	(15.1 - 17.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	15.0	(11.6 - 18.4)	11.1	(10.5 - 11.6)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	14.5	(11.0 - 18.0)	14.3	(13.7 - 14.9)			+
Sufficient vigorous activity	63.8	(58.6 - 68.9)	62.7	(62.0 - 63.5)			+
Sufficient moderate activity	31.8	(26.3 - 37.4)	26.1	(25.4 - 26.7)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	44.2	(39.4 - 49.0)	42.3	(41.5 - 43.1)			+
Described themselves as slightly or very overweight	33.9	(28.9 - 39.0)	29.8	(29.0 - 30.5)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	13.1	(9.6 - 16.6)	11.2	(10.5 - 11.8)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	24.6	(21.1 - 28.1)	23.1	(22.4 - 23.8)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	8.0	(5.4 - 10.6)	6.9	(6.4 - 7.3)			+
Purposely hurt themselves without wanting to die, during the past 12 months	17.3	(13.6 - 20.9)	12.7	(12.1 - 13.2)	+		
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	25.1	(21.0 - 29.1)	24.5	(23.8 - 25.2)			+

Citrus County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Citrus County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	12.4	(9.6 - 15.3)	6.1	(5.8 - 6.4)	+		
Smoked cigars on one or more of the past 30 days	8.3	(5.8 - 10.9)	6.9	(6.6 - 7.2)			+
Used smokeless tobacco on one or more of the past 30 days	8.8	(6.3 - 11.3)	3.5	(3.2 - 3.7)	+		
Used any form of tobacco on one or more of the past 30 days	19.1	(15.5 - 22.7)	11.2	(10.8 - 11.7)	+		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	55.6	(50.9 - 60.3)	65.7	(65.1 - 66.3)		+	
Exposed to secondhand smoke during the past 7 days	63.4	(59.2 - 67.6)	44.1	(43.4 - 44.7)	+		
Smoking is allowed in the home	17.8	(14.9 - 20.6)	9.4	(9.0 - 9.8)	+		
Taught about tobacco use in any classes this year	54.0	(48.6 - 59.4)	43.2	(42.3 - 44.1)	+		
Parent or guardian has talked about the dangers of tobacco in the past year	56.0	(51.8 - 60.2)	57.4	(56.8 - 58.0)			+
Served as an active member of SWAT	3.4	(2.0 - 4.7)	3.9	(3.7 - 4.1)			+
<u>Definitely did not think</u> that smokers have more friends	27.5	(24.0 - 31.0)	32.0	(31.5 - 32.6)		+	
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	73.6	(70.1 - 77.1)	72.9	(72.4 - 73.4)			+

Citrus County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Citrus County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	22.3	(19.1 - 25.5)	20.5	(20.1 - 21.0)			+
Current asthma, among all individuals	12.7	(10.2 - 15.3)	11.3	(11.0 - 11.7)			+
Asthma attack in past year, among individuals with lifetime asthma	17.0	(10.7 - 23.3)	16.9	(16.0 - 17.8)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	13.1	(10.6 - 15.7)	11.1	(10.7 - 11.5)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	15.3	(12.5 - 18.2)	15.4	(15.0 - 15.9)			+
Sufficient vigorous activity	70.2	(66.1 - 74.3)	66.5	(65.9 - 67.1)			+
Sufficient moderate activity	28.2	(24.3 - 32.1)	25.3	(24.8 - 25.8)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	45.5	(42.0 - 49.0)	43.5	(42.9 - 44.1)			+
Described themselves as slightly or very overweight	33.9	(30.2 - 37.6)	30.0	(29.5 - 30.6)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	12.1	(9.6 - 14.5)	10.9	(10.4 - 11.3)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	22.0	(19.2 - 24.8)	20.2	(19.7 - 20.7)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	8.0	(6.1 - 10.0)	6.9	(6.6 - 7.2)			+
Purposely hurt themselves without wanting to die, during the past 12 months	15.8	(12.9 - 18.7)	12.4	(12.0 - 12.8)	+		
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	23.3	(20.2 - 26.5)	22.0	(21.5 - 22.5)			+