Seminole County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Seminole County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	**3.6**	(2.1 - 5.2)	**3.3**	(3.0 - 3.5)			+
Smoked cigars on one or more of the past 30 days	**5.7**	(3.8 - 7.5)	**3.9**	(3.6 - 4.2)			+
Used smokeless tobacco on one or more of the past 30 days	**3.0**	(1.5 - 4.5)	**2.3**	(2.1 - 2.6)			+
Used any form of tobacco on one or more of the past 30 days	**7.1**	(5.1 - 9.2)	**6.4**	(6.0 - 6.8)			+
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	**76.0**	(72.0 - 80.0)	**72.0**	(71.2 - 72.8)			+
Exposed to secondhand smoke during the past 7 days	**37.6**	(33.3 - 41.9)	**41.3**	(40.4 - 42.2)			+
Smoking is allowed in the home	**8.4**	(5.9 - 11.0)	**9.9**	(9.5 - 10.4)			+
Taught about tobacco use in any classes this year	**40.3**	(34.1 - 46.5)	**53.4**	(52.0 - 54.7)		+	
Parent or guardian has talked about the dangers of tobacco in the past year	**64.0**	(60.4 - 67.7)	**63.9**	(63.1 - 64.6)			+
Served as an active member of SWAT	**2.6**	(1.3 - 3.8)	**4.3**	(3.9 - 4.7)		+	
Definitely did not think that smokers have more friends	**36.5**	(32.8 - 40.1)	**33.7**	(32.9 - 34.5)			+
Definitely did not think that smoking helps young people “look cool” or “fit in”	**77.9**	(74.3 - 81.6)	**73.7**	(73.0 - 74.4)			+

Seminole County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Seminole County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	**19.2**	(16.3 - 22.1)	**20.1**	(19.5 - 20.7)			+
Current asthma, among all individuals	**12.6**	(10.2 - 14.9)	**12.0**	(11.5 - 12.5)			+
Asthma attack in past year, among individuals with lifetime asthma	**21.7**	(14.3 - 29.0)	**18.1**	(16.8 - 19.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	**6.5**	(4.4 - 8.6)	**11.6**	(11.0 - 12.1)		+	
Overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex)	**12.9**	(9.8 - 16.0)	**16.5**	(15.8 - 17.2)			+
Sufficient vigorous activity	**73.7**	(69.8 - 77.6)	**70.1**	(69.2 - 70.9)			+
Sufficient moderate activity	**27.2**	(23.8 - 30.6)	**24.8**	(24.1 - 25.4)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	**39.7**	(35.7 - 43.6)	**44.4**	(43.6 - 45.2)			+
Described themselves as slightly or very overweight	**29.3**	(25.5 -33.1)	**30.2**	(29.4 - 30.9)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	**5.0**	(3.0 - 7.1)	**11.4**	(10.8 - 12.0)		+	
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	**17.5**	(14.0- 20.9)	**17.8**	(17.2- 18.5)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	**7.8**	(5.6 - 10.0)	**7.8**	(7.3 - 8.3)			+
Purposely hurt themselves without wanting to die, during the past 12 months	**11.9**	(9.2 - 14.7)	**12.6**	(12.0 - 13.2)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	**20.1**	(16.7 - 23.5)	**19.2**	(18.5 - 19.8)			+

Seminole County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Seminole County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	**7.5**	(5.6 - 9.4)	**10.1**	(9.6 - 10.6)		+	
Smoked cigars on one or more of the past 30 days	**10.3**	(8.0 - 12.7)	**11.4**	(10.9 - 11.9)			+
Used smokeless tobacco on one or more of the past 30 days	**4.3**	(2.6 - 6.0)	**5.6**	(5.2 - 6.0)			+
Used any form of tobacco on one or more of the past 30 days	**15.8**	(13.1 - 18.5)	**17.9**	(17.3 - 18.6)			+
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	**58.6**	(54.4 - 62.8)	**58.4**	(57.6 - 59.3)			+
Exposed to secondhand smoke during the past 7 days	**42.1**	(38.2 - 46.1)	**47.6**	(46.7 - 48.4)		+	
Smoking is allowed in the home	**8.4**	(6.2 - 10.6)	**10.0**	(9.5 - 10.5)			+
Taught about tobacco use in any classes this year	**25.3**	(22.2 - 28.4)	**33.4**	(32.4 - 34.4)		+	
Parent or guardian has talked about the dangers of tobacco in the past year	**50.6**	(47.1 - 54.2)	**50.4**	(49.6 - 51.2)			+
Served as an active member of SWAT	**2.9**	(1.7 - 4.2)	**4.0**	(3.7 - 4.3)			+
Definitely did not think that smokers have more friends	**30.8**	(27.1 - 34.4)	**30.5**	(29.8 - 31.2)			+
Definitely did not think that smoking helps young people “look cool” or “fit in”	**71.6**	(68.4 - 74.8)	**71.7**	(71.0 - 72.4)			+

Seminole County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Seminole County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	**20.3**	(17.2 - 23.4)	**20.8**	(20.2 - 21.5)			+
Current asthma, among all individuals	**11.4**	(9.3 - 13.6)	**10.6**	(10.1 - 11.1)			+
Asthma attack in past year, among individuals with lifetime asthma	**17.6**	(12.1 - 23.2)	**16.3**	(15.1 - 17.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	**10.2**	(7.9 - 12.4)	**11.1**	(10.5 - 11.6)			+
Overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex)	**11.0**	(8.6 - 13.5)	**14.3**	(13.7 - 14.9)		+	
Sufficient vigorous activity	**66.7**	(63.6 - 69.9)	**62.7**	(62.0 - 63.5)	+		
Sufficient moderate activity	**27.2**	(24.0 - 30.3)	**26.1**	(25.4 - 26.7)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	**42.3**	(38.3 - 46.3)	**42.3**	(41.5 - 43.1)			+
Described themselves as slightly or very overweight	**28.1**	(24.6 - 31.6)	**29.8**	(29.0 - 30.5)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	**8.2**	(5.9 - 10.5)	**11.2**	(10.5 - 11.8)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	**20.5**	(17.6 - 23.3)	**23.1**	(22.4 - 23.8)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	**5.9**	(3.8 - 8.0)	**6.9**	(6.4 - 7.3)			+
Purposely hurt themselves without wanting to die, during the past 12 months	**13.7**	(10.7 - 16.8)	**12.7**	(12.1 - 13.2)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	**26.6**	(22.7 - 30.4)	**24.5**	(23.8 - 25.2)			+

Seminole County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Seminole County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	**5.3**	(4.0 - 6.6)	**6.1**	(5.8 - 6.4)			+
Smoked cigars on one or more of the past 30 days	**7.1**	(5.7 - 8.6)	**6.9**	(6.6 - 7.2)			+
Used smokeless tobacco on one or more of the past 30 days	**2.9**	(1.9 - 3.9)	**3.5**	(3.2 - 3.7)			+
Used any form of tobacco on one or more of the past 30 days	**10.6**	(8.9 - 12.3)	**11.2**	(10.8 - 11.7)			+
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	**68.0**	(64.8 - 71.2)	**65.7**	(65.1 - 66.3)			+
Exposed to secondhand smoke during the past 7 days	**39.1**	(36.1 - 42.1)	**44.1**	(43.4 - 44.7)		+	
Smoking is allowed in the home	**7.1**	(5.6 - 8.7)	**9.4**	(9.0 - 9.8)		+	
Taught about tobacco use in any classes this year	**31.1**	(27.7 - 34.6)	**43.2**	(42.3 - 44.1)		+	
Parent or guardian has talked about the dangers of tobacco in the past year	**57.5**	(54.8 - 60.2)	**57.4**	(56.8 - 58.0)			+
Served as an active member of SWAT	**2.4**	(1.6 - 3.3)	**3.9**	(3.7 - 4.1)		+	
Definitely did not think that smokers have more friends	**34.1**	(31.4 - 36.8)	**32.0**	(31.5 - 32.6)			+
Definitely did not think that smoking helps young people “look cool” or “fit in”	**75.1**	(72.7 - 77.6)	**72.9**	(72.4 - 73.4)			+

Seminole County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Seminole County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	**19.9**	(17.6 - 22.1)	**20.5**	(20.1 - 21.0)			+
Current asthma, among all individuals	**12.0**	(10.3 - 13.6)	**11.3**	(11.0 - 11.7)			+
Asthma attack in past year, among individuals with lifetime asthma	**19.3**	(14.6 - 24.0)	**16.9**	(16.0 - 17.8)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	**8.6**	(7.0 - 10.3)	**11.1**	(10.7 - 11.5)		+	
Overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex)	**11.5**	(9.6 - 13.4)	**15.4**	(15.0 - 15.9)		+	
Sufficient vigorous activity	**70.1**	(67.6 - 72.6)	**66.5**	(65.9 - 67.1)	+		
Sufficient moderate activity	**27.2**	(24.8 - 29.5)	**25.3**	(24.8 - 25.8)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	**41.6**	(38.7 - 44.5)	**43.5**	(42.9 - 44.1)			+
Described themselves as slightly or very overweight	**28.7**	(26.1 - 31.3)	**30.0**	(29.5 - 30.6)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	**6.1**	(4.5 - 7.7)	**10.9**	(10.4 - 11.3)		+	
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	**17.9**	(15.8 - 20.0)	**20.2**	(19.7 - 20.7)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	**6.0**	(4.5 - 7.5)	**6.9**	(6.6 - 7.2)			+
Purposely hurt themselves without wanting to die, during the past 12 months	**12.6**	(10.4 - 14.7)	**12.4**	(12.0 - 12.8)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	**23.2**	(20.4 - 25.9)	**22.0**	(21.5 - 22.5)			+