



Collier County: 2014 at a Glance

Data from the 2014 Florida Youth Tobacco Survey (FYTS)



Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Collier County %	95% CI	State %	95% CI	County Significantly Higher Than State	State Significantly Higher Than County	No Significant Difference
Smoked cigarettes on one or more of the past 30 days	4.2	(2.9-5.6)	4.3	(4.1-4.6)			+
Smoked cigars on one or more of the past 30 days	4.9	(3.3-6.6)	5.4	(5.2-5.7)			+
Used smokeless tobacco on one or more of the past 30 days	2.3	(1.4-3.1)	3.3	(3.1-3.5)			+
Used electronic cigarettes on one or more of the past 30 days	9.8	(7.7-12.0)	7.2	(6.8-7.5)	+		
Used hookah on one or more of the past 30 days	3.7	(2.3-5.0)	7.1	(6.7-7.5)		+	
Used any form of tobacco on one or more of the past 30 days*	7.8	(5.8-9.8)	9.2	(8.8-9.6)			+
Used any form of tobacco (excluding cigarettes) on one or more of the past 30 days	13.1	(10.1-16.0)	14.8	(14.2-15.3)			+
Thought using electronic cigarettes is less harmful than smoking cigarettes	73.3	(69.2-77.4)	69.4	(68.4-70.3)			+
Thought using hookah is less harmful than smoking cigarettes	42.0	(36.2-47.7)	52.8	(51.7-54.0)		+	
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	67.8	(64.4-71.1)	67.1	(66.4-67.7)			+
Exposed to secondhand smoke during the past seven days (in a room or car)	32.3	(29.2-35.4)	37.5	(36.9-38.2)		+	
Smoking is allowed in the home	4.7	(3.4-6.0)	8.4	(8.0-8.7)		+	

*This excludes specialty tobacco products, which include bidis, kreteks, pipe tobacco, snus, e-cigarettes, flavored cigarettes, flavored cigars, and flavored smokeless tobacco.



Collier County: 2014 at a Glance

Data from the 2014 Florida Youth Tobacco Survey (FYTS)



Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Collier County %	95% CI	State %	95% CI	County Significantly Higher Than State	State Significantly Higher Than County	No Significant Difference
Students who were taught about tobacco use in any classes this year	36.9	(32.2-41.6)	38.0	(37.1-39.0)			+
Parent or guardian has talked about the dangers of tobacco in the past year	63.1	(59.5-66.7)	62.4	(61.8-63.0)			+
Served as an active member of Students Working Against Tobacco (SWAT)	3.7	(2.6-4.7)	2.8	(2.7- 3.0)			+
Definitely did not think that smokers have more friends	28.4	(25.7-31.1)	28.2	(27.6-28.8)			+
Definitely did not think that smoking helps young people "look cool" or "fit in"	68.7	(65.8-71.6)	68.8	(68.2-69.3)			+
Lifetime asthma	16.2	(13.7-18.7)	20.8	(20.3-21.3)		+	
Current asthma	8.5	(6.4-10.5)	11.5	(11.1-11.9)		+	
Asthma attack in past year	21.1	(15.0-27.2)	17.7	(16.7-18.7)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	10.5	(8.5-12.4)	12.2	(11.8-12.7)			+
Overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex)	19.1	(16.8-21.5)	15.9	(15.4-16.4)	+		
Physically active for at least 60 minutes per day during the past seven days	22.7	(20.3-25.0)	22.7	(22.2-23.3)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	44.3	(40.7-47.9)	43.7	(43.1-44.4)			+
Described themselves as slightly or very overweight	31.4	(28.3-34.5)	30.7	(30.2-31.3)			+