



# Okaloosa County: 2014 at a Glance

Data from the 2014 Florida Youth Tobacco Survey (FYTS)



Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Okaloosa County %	95% CI	State %	95% CI	County Significantly Higher Than State	State Significantly Higher Than County	No Significant Difference
Smoked cigarettes on one or more of the past 30 days	<b>3.1</b>	(1.8-4.5)	<b>4.3</b>	(4.1-4.6)			+
Smoked cigars on one or more of the past 30 days	<b>5.2</b>	(3.0-7.5)	<b>5.4</b>	(5.2-5.7)			+
Used smokeless tobacco on one or more of the past 30 days	<b>4.5</b>	(2.5-6.5)	<b>3.3</b>	(3.1-3.5)			+
Used electronic cigarettes on one or more of the past 30 days	<b>8.0</b>	(5.3-10.6)	<b>7.2</b>	(6.8-7.5)			+
Used hookah on one or more of the past 30 days	<b>5.1</b>	(3.0-7.2)	<b>7.1</b>	(6.7-7.5)			+
Used any form of tobacco on one or more of the past 30 days*	<b>9.0</b>	(6.1-11.9)	<b>9.2</b>	(8.8-9.6)			+
Used any form of tobacco (excluding cigarettes) on one or more of the past 30 days	<b>14.4</b>	(10.9-18.0)	<b>14.8</b>	(14.2-15.3)			+
Thought using electronic cigarettes is less harmful than smoking cigarettes	<b>79.1</b>	(74.3-83.9)	<b>69.4</b>	(68.4-70.3)	+		
Thought using hookah is less harmful than smoking cigarettes	<b>46.6</b>	(39.1-54.2)	<b>52.8</b>	(51.7-54.0)			+
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	<b>67.2</b>	(62.8-71.5)	<b>67.1</b>	(66.4-67.7)			+
Exposed to secondhand smoke during the past seven days (in a room or car)	<b>39.6</b>	(34.9-44.3)	<b>37.5</b>	(36.9-38.2)			+
Smoking is allowed in the home	<b>7.3</b>	(5.0-9.6)	<b>8.4</b>	(8.0-8.7)			+

\*This excludes specialty tobacco products, which include bidis, kreteks, pipe tobacco, snus, e-cigarettes, flavored cigarettes, flavored cigars, and flavored smokeless tobacco.



# Okaloosa County: 2014 at a Glance

Data from the 2014 Florida Youth Tobacco Survey (FYTS)



Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Okaloosa County %	95% CI	State %	95% CI	County Significantly Higher Than State	State Significantly Higher Than County	No Significant Difference
Students who were taught about tobacco use in any classes this year	<b>35.8</b>	(30.2-41.4)	<b>38.0</b>	(37.1-39.0)			+
Parent or guardian has talked about the dangers of tobacco in the past year	<b>64.5</b>	(60.6-68.5)	<b>62.4</b>	(61.8-63.0)			+
Served as an active member of Students Working Against Tobacco (SWAT)	<b>1.9</b>	(0.7-3.1)	<b>2.8</b>	(2.7- 3.0)			+
Definitely did not think that smokers have more friends	<b>26.4</b>	(23.0-29.7)	<b>28.2</b>	(27.6-28.8)			+
Definitely did not think that smoking helps young people "look cool" or "fit in"	<b>72.1</b>	(68.4-75.8)	<b>68.8</b>	(68.2-69.3)			+
Lifetime asthma	<b>22.3</b>	(19.1-25.5)	<b>20.8</b>	(20.3-21.3)			+
Current asthma	<b>11.2</b>	(8.9-13.5)	<b>11.5</b>	(11.1-11.9)			+
Asthma attack in past year	<b>13.1</b>	(6.8-19.4)	<b>17.7</b>	(16.7-18.7)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	<b>9.9</b>	(7.3-12.5)	<b>12.2</b>	(11.8-12.7)			+
Overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex)	<b>13.0</b>	(10.3-15.6)	<b>15.9</b>	(15.4-16.4)			+
Physically active for at least 60 minutes per day during the past seven days	<b>21.1</b>	(18.1-24.2)	<b>22.7</b>	(22.2-23.3)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	<b>40.5</b>	(36.5-44.6)	<b>43.7</b>	(43.1-44.4)			+
Described themselves as slightly or very overweight	<b>28.0</b>	(24.2-31.7)	<b>30.7</b>	(30.2-31.3)			+