Bay County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Bay County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	**8.6**	(6.8 - 10.4)	**6.1**	(5.8 - 6.4)	+		
Smoked cigars on one or more of the past 30 days	**7.5**	(5.9 - 9.1)	**6.9**	(6.6 - 7.2)			+
Used smokeless tobacco on one or more of the past 30 days	**4.5**	(3.3 - 5.8)	**3.5**	(3.2 - 3.7)			+
Used any form of tobacco on one or more of the past 30 days	**14.5**	(12.3 - 16.7)	**11.2**	(10.8 - 11.7)	+		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	**61.6**	(58.3 - 65.0)	**65.7**	(65.1 - 66.3)		+	
Exposed to secondhand smoke during the past 7 days	**50.2**	(47.0 - 53.5)	**44.1**	(43.4 - 44.7)	+		
Smoking is allowed in the home	**14.6**	(12.6 - 16.5)	**9.4**	(9.0 - 9.8)	+		
Taught about tobacco use in any classes this year	**35.1**	(31.2 - 38.9)	**43.2**	(42.3 - 44.1)		+	
Parent or guardian has talked about the dangers of tobacco in the past year	**54.5**	(51.7 - 57.2)	**57.4**	(56.8 - 58.0)			+
Served as an active member of SWAT	**7.2**	(5.7 - 8.6)	**3.9**	(3.7 - 4.1)	+		
Definitely did not think that smokers have more friends	**31.0**	(28.2 - 33.8)	**32.0**	(31.5 - 32.6)			+
Definitely did not think that smoking helps young people “look cool” or “fit in”	**73.4**	(70.9 - 75.8)	**72.9**	(72.4 - 73.4)			+

Bay County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Bay County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	**19.3**	(17.5 - 21.2)	**20.5**	(20.1 - 21.0)			+
Current asthma, among all individuals	**11.1**	(9.6 - 12.6)	**11.3**	(11.0 - 11.7)			+
Asthma attack in past year, among individuals with lifetime asthma	**21.2**	(16.7 - 25.7)	**16.9**	(16.0 - 17.8)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	**10.8**	(8.9 - 12.6)	**11.1**	(10.7 - 11.5)			+
Overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex)	**15.3**	(13.1 - 17.4)	**15.4**	(15.0 - 15.9)			+
Sufficient vigorous activity	**66.2**	(63.8 - 68.6)	**66.5**	(65.9 - 67.1)			+
Sufficient moderate activity	**28.1**	(25.7 - 30.5)	**25.3**	(24.8 - 25.8)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	**43.3**	(40.5 - 46.2)	**43.5**	(42.9 - 44.1)			+
Described themselves as slightly or very overweight	**29.8**	(27.3 - 32.3)	**30.0**	(29.5 - 30.6)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	**10.2**	(8.4 - 12.0)	**10.9**	(10.4 - 11.3)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	**21.0**	(19.0 - 23.0)	**20.2**	(19.7 - 20.7)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	**6.1**	(4.8 - 7.4)	**6.9**	(6.6 - 7.2)			+
Purposely hurt themselves without wanting to die, during the past 12 months	**13.6**	(11.6 - 15.6)	**12.4**	(12.0 - 12.8)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	**21.8**	(19.4 - 24.2)	**22.0**	(21.5 - 22.5)			+