

Calhoun County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Calhoun County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	6.5	(3.3 - 9.7)	6.1	(5.8 - 6.4)			+
Smoked cigars on one or more of the past 30 days	6.2	(3.6 - 8.9)	6.9	(6.6 - 7.2)			+
Used smokeless tobacco on one or more of the past 30 days	4.8	(2.6 - 6.9)	3.5	(3.2 - 3.7)			+
Used any form of tobacco on one or more of the past 30 days	11.4	(7.4 - 15.3)	11.2	(10.8 - 11.7)			+
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	63.5	(55.5 - 71.4)	65.7	(65.1 - 66.3)			+
Exposed to secondhand smoke during the past 7 days	54.4	(47.9 - 61.0)	44.1	(43.4 - 44.7)	+		
Smoking is allowed in the home	17.4	(12.6 - 22.2)	9.4	(9.0 - 9.8)	+		
Taught about tobacco use in any classes this year	62.2	(55.6 - 68.7)	43.2	(42.3 - 44.1)	+		
Parent or guardian has talked about the dangers of tobacco in the past year	52.0	(45.5 - 58.4)	57.4	(56.8 - 58.0)			+
Served as an active member of SWAT	21.3	(16.5 - 26.1)	3.9	(3.7 - 4.1)	+		
<u>Definitely did not think</u> that smokers have more friends	37.6	(31.7 - 43.6)	32.0	(31.5 - 32.6)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	77.3	(72.0 - 82.7)	72.9	(72.4 - 73.4)			+

Calhoun County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Calhoun County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	19.5	(14.8 - 24.2)	20.5	(20.1 - 21.0)			+
Current asthma, among all individuals	11.9	(8.5 - 15.3)	11.3	(11.0 - 11.7)			+
Asthma attack in past year, among individuals with lifetime asthma	21.6	(10.4 - 32.8)	16.9	(16.0 - 17.8)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	18.8	(14.5 - 23.1)	11.1	(10.7 - 11.5)	+		
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	16.4	(12.6 - 20.2)	15.4	(15.0 - 15.9)			+
Sufficient vigorous activity	71.1	(65.8 - 76.4)	66.5	(65.9 - 67.1)			+
Sufficient moderate activity	27.6	(22.9 - 32.3)	25.3	(24.8 - 25.8)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	50.0	(45.7 - 54.3)	43.5	(42.9 - 44.1)	+		
Described themselves as slightly or very overweight	34.3	(30.3 - 38.3)	30.0	(29.5 - 30.6)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	10.5	(6.8 - 14.3)	10.9	(10.4 - 11.3)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	18.3	(13.9 - 22.7)	20.2	(19.7 - 20.7)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	5.6	(2.9 - 8.3)	6.9	(6.6 - 7.2)			+
Purposely hurt themselves without wanting to die, during the past 12 months	10.5	(6.3 - 14.8)	12.4	(12.0 - 12.8)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	20.1	(15.4 - 24.8)	22.0	(21.5 - 22.5)			+