

Clay County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Clay County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	8.0	(6.2 - 9.8)	6.1	(5.8 - 6.4)			+
Smoked cigars on one or more of the past 30 days	8.0	(6.3 - 9.8)	6.9	(6.6 - 7.2)			+
Used smokeless tobacco on one or more of the past 30 days	5.1	(3.4 - 6.7)	3.5	(3.2 - 3.7)			+
Used any form of tobacco on one or more of the past 30 days	14.7	(12.1 - 17.4)	11.2	(10.8 - 11.7)	+		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	64.0	(61.1 - 67.0)	65.7	(65.1 - 66.3)			+
Exposed to secondhand smoke during the past 7 days	51.5	(48.3 - 54.7)	44.1	(43.4 - 44.7)	+		
Smoking is allowed in the home	11.5	(9.7 - 13.4)	9.4	(9.0 - 9.8)			+
Taught about tobacco use in any classes this year	43.9	(39.3 - 48.4)	43.2	(42.3 - 44.1)			+
Parent or guardian has talked about the dangers of tobacco in the past year	55.0	(52.0 - 58.1)	57.4	(56.8 - 58.0)			+
Served as an active member of SWAT	4.3	(3.1 - 5.4)	3.9	(3.7 - 4.1)			+
<u>Definitely did not think</u> that smokers have more friends	30.8	(28.1 - 33.5)	32.0	(31.5 - 32.6)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	75.9	(73.7 - 78.2)	72.9	(72.4 - 73.4)	+		

Clay County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Clay County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	19.8	(17.9 - 21.8)	20.5	(20.1 - 21.0)			+
Current asthma, among all individuals	11.3	(9.6 - 13.0)	11.3	(11.0 - 11.7)			+
Asthma attack in past year, among individuals with lifetime asthma	20.1	(15.0 - 25.2)	16.9	(16.0 - 17.8)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	9.4	(7.8 - 10.9)	11.1	(10.7 - 11.5)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	13.6	(11.8 - 15.5)	15.4	(15.0 - 15.9)			+
Sufficient vigorous activity	72.2	(69.6 - 74.9)	66.5	(65.9 - 67.1)	+		
Sufficient moderate activity	28.7	(26.2 - 31.1)	25.3	(24.8 - 25.8)	+		
Exercised to lose weight or to keep from gaining weight during the past 30 days	41.3	(38.6 - 43.9)	43.5	(42.9 - 44.1)			+
Described themselves as slightly or very overweight	28.9	(26.4 - 31.3)	30.0	(29.5 - 30.6)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	8.9	(7.2 - 10.7)	10.9	(10.4 - 11.3)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	17.9	(15.9 - 20.0)	20.2	(19.7 - 20.7)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	5.2	(3.9 - 6.5)	6.9	(6.6 - 7.2)		+	
Purposely hurt themselves without wanting to die, during the past 12 months	12.9	(11.3 - 14.6)	12.4	(12.0 - 12.8)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	21.3	(19.1 - 23.5)	22.0	(21.5 - 22.5)			+