

Flagler County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Flagler County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	9.7	(7.0 - 12.4)	6.1	(5.8 - 6.4)	+		
Smoked cigars on one or more of the past 30 days	10.2	(7.7 - 12.7)	6.9	(6.6 - 7.2)	+		
Used smokeless tobacco on one or more of the past 30 days	4.5	(3.0 - 6.1)	3.5	(3.2 - 3.7)			+
Used any form of tobacco on one or more of the past 30 days	16.1	(12.7 - 19.5)	11.2	(10.8 - 11.7)	+		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	62.1	(57.8 - 66.5)	65.7	(65.1 - 66.3)			+
Exposed to secondhand smoke during the past 7 days	54.4	(50.1 - 58.7)	44.1	(43.4 - 44.7)	+		
Smoking is allowed in the home	12.0	(9.4 - 14.6)	9.4	(9.0 - 9.8)			+
Taught about tobacco use in any classes this year	35.1	(29.3 - 40.8)	43.2	(42.3 - 44.1)		+	
Parent or guardian has talked about the dangers of tobacco in the past year	55.2	(51.3 - 59.1)	57.4	(56.8 - 58.0)			+
Served as an active member of SWAT	2.4	(1.4 - 3.5)	3.9	(3.7 - 4.1)		+	
<u>Definitely did not think</u> that smokers have more friends	27.4	(24.2 - 30.5)	32.0	(31.5 - 32.6)		+	
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	68.5	(65.3 - 71.8)	72.9	(72.4 - 73.4)		+	

Flagler County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Flagler County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	23.0	(20.0 - 26.0)	20.5	(20.1 - 21.0)			+
Current asthma, among all individuals	14.1	(11.8 - 16.5)	11.3	(11.0 - 11.7)	+		
Asthma attack in past year, among individuals with lifetime asthma	18.0	(12.2 - 23.7)	16.9	(16.0 - 17.8)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	10.2	(7.9 - 12.5)	11.1	(10.7 - 11.5)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	15.7	(13.1 - 18.3)	15.4	(15.0 - 15.9)			+
Sufficient vigorous activity	67.8	(64.0 - 71.7)	66.5	(65.9 - 67.1)			+
Sufficient moderate activity	27.8	(24.4 - 31.2)	25.3	(24.8 - 25.8)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	46.0	(42.5 - 49.6)	43.5	(42.9 - 44.1)			+
Described themselves as slightly or very overweight	30.8	(27.6 - 34.0)	30.0	(29.5 - 30.6)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	11.4	(9.2 - 13.6)	10.9	(10.4 - 11.3)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	24.7	(21.1 - 28.3)	20.2	(19.7 - 20.7)	+		
Did not go to school on one or more of the past 30 days because they felt unsafe	6.0	(4.1 - 7.8)	6.9	(6.6 - 7.2)			+
Purposely hurt themselves without wanting to die, during the past 12 months	16.0	(13.1 - 19.0)	12.4	(12.0 - 12.8)	+		
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	26.2	(22.8 - 29.6)	22.0	(21.5 - 22.5)	+		