

Glades County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Glades County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	8.4	(4.1 - 12.7)	6.1	(5.8 - 6.4)			+
Smoked cigars on one or more of the past 30 days	9.1	(5.2 - 13.0)	6.9	(6.6 - 7.2)			+
Used smokeless tobacco on one or more of the past 30 days	6.2	(2.6 - 9.7)	3.5	(3.2 - 3.7)			+
Used any form of tobacco on one or more of the past 30 days	16.6	(11.6 - 21.7)	11.2	(10.8 - 11.7)			+
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	57.0	(50.2 - 63.8)	65.7	(65.1 - 66.3)		+	
Exposed to secondhand smoke during the past 7 days	47.3	(40.0 - 54.6)	44.1	(43.4 - 44.7)			+
Smoking is allowed in the home	22.4	(16.3 - 28.6)	9.4	(9.0 - 9.8)	+		
Taught about tobacco use in any classes this year	51.0	(40.4 - 61.6)	43.2	(42.3 - 44.1)			+
Parent or guardian has talked about the dangers of tobacco in the past year	55.8	(47.0 - 64.5)	57.4	(56.8 - 58.0)			+
Served as an active member of SWAT	19.7	(13.3 - 26.2)	3.9	(3.7 - 4.1)	+		
<u>Definitely did not think</u> that smokers have more friends	32.9	(26.5 - 39.2)	32.0	(31.5 - 32.6)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	66.1	(59.8 - 72.4)	72.9	(72.4 - 73.4)			+

Glades County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Glades County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	19.5	(15.1 - 23.9)	20.5	(20.1 - 21.0)			+
Current asthma, among all individuals	11.6	(7.1 - 16.0)	11.3	(11.0 - 11.7)			+
Asthma attack in past year, among individuals with lifetime asthma	16.5	(4.3 - 28.8)	16.9	(16.0 - 17.8)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	16.9	(10.9 - 23.0)	11.1	(10.7 - 11.5)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	19.4	(13.6 - 25.2)	15.4	(15.0 - 15.9)			+
Sufficient vigorous activity	70.6	(62.1 - 79.1)	66.5	(65.9 - 67.1)			+
Sufficient moderate activity	27.1	(20.4 - 33.8)	25.3	(24.8 - 25.8)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	46.4	(37.6 - 55.2)	43.5	(42.9 - 44.1)			+
Described themselves as slightly or very overweight	38.8	(32.2 - 45.5)	30.0	(29.5 - 30.6)	+		
Never or rarely wore a seat belt when riding in a car driven by someone else	20.9	(14.9 - 26.9)	10.9	(10.4 - 11.3)	+		
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	22.3	(17.7 - 26.9)	20.2	(19.7 - 20.7)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	6.8	(3.4 - 10.2)	6.9	(6.6 - 7.2)			+
Purposely hurt themselves without wanting to die, during the past 12 months	17.6	(13.0 - 22.3)	12.4	(12.0 - 12.8)	+		
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	28.6	(22.0 - 35.3)	22.0	(21.5 - 22.5)			+