

Santa Rosa County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Santa Rosa County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	9.1	(6.9 - 11.4)	6.1	(5.8 - 6.4)	+		
Smoked cigars on one or more of the past 30 days	7.6	(5.3 - 9.9)	6.9	(6.6 - 7.2)			+
Used smokeless tobacco on one or more of the past 30 days	6.0	(4.1 - 7.8)	3.5	(3.2 - 3.7)	+		
Used any form of tobacco on one or more of the past 30 days	15.2	(12.2 - 18.1)	11.2	(10.8 - 11.7)	+		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	65.9	(61.9 - 69.9)	65.7	(65.1 - 66.3)			+
Exposed to secondhand smoke during the past 7 days	45.6	(42.3 - 49.0)	44.1	(43.4 - 44.7)			+
Smoking is allowed in the home	9.4	(7.5 - 11.3)	9.4	(9.0 - 9.8)			+
Taught about tobacco use in any classes this year	58.1	(53.2 - 63.0)	43.2	(42.3 - 44.1)	+		
Parent or guardian has talked about the dangers of tobacco in the past year	59.1	(56.0 - 62.3)	57.4	(56.8 - 58.0)			+
Served as an active member of SWAT	4.2	(2.8 - 5.5)	3.9	(3.7 - 4.1)			+
<u>Definitely did not think</u> that smokers have more friends	34.5	(31.0 - 38.0)	32.0	(31.5 - 32.6)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	74.5	(71.4 - 77.6)	72.9	(72.4 - 73.4)			+

Santa Rosa County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Santa Rosa County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	22.0	(19.3 - 24.8)	20.5	(20.1 - 21.0)			+
Current asthma, among all individuals	13.6	(11.5 - 15.7)	11.3	(11.0 - 11.7)			+
Asthma attack in past year, among individuals with lifetime asthma	21.9	(17.0 - 26.8)	16.9	(16.0 - 17.8)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	10.5	(8.5 - 12.5)	11.1	(10.7 - 11.5)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	15.8	(13.1 - 18.6)	15.4	(15.0 - 15.9)			+
Sufficient vigorous activity	68.2	(65.0 - 71.3)	66.5	(65.9 - 67.1)			+
Sufficient moderate activity	29.1	(26.0 - 32.2)	25.3	(24.8 - 25.8)	+		
Exercised to lose weight or to keep from gaining weight during the past 30 days	43.0	(39.9 - 46.1)	43.5	(42.9 - 44.1)			+
Described themselves as slightly or very overweight	28.1	(25.3 - 31.0)	30.0	(29.5 - 30.6)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	9.3	(6.9 - 11.6)	10.9	(10.4 - 11.3)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	18.4	(15.8 - 20.9)	20.2	(19.7 - 20.7)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	4.7	(3.2 - 6.3)	6.9	(6.6 - 7.2)		+	
Purposely hurt themselves without wanting to die, during the past 12 months	12.5	(10.6 - 14.5)	12.4	(12.0 - 12.8)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	18.6	(16.0 - 21.2)	22.0	(21.5 - 22.5)		+	