

Holmes County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Holmes County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	6.4	(2.9 - 10.0)	3.3	(3.0 - 3.5)			+
Smoked cigars on one or more of the past 30 days	4.1	(2.1 - 6.1)	3.9	(3.6 - 4.2)			+
Used smokeless tobacco on one or more of the past 30 days	8.9	(5.3 - 12.5)	2.3	(2.1 - 2.6)	+		
Used any form of tobacco on one or more of the past 30 days	13.4	(9.3 - 17.5)	6.4	(6.0 - 6.8)	+		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	69.6	(62.8 - 76.3)	72.0	(71.2 - 72.8)			+
Exposed to secondhand smoke during the past 7 days	52.0	(46.4 - 57.7)	41.3	(40.4 - 42.2)	+		
Smoking is allowed in the home	24.1	(18.3 - 30.0)	9.9	(9.5 - 10.4)	+		
Taught about tobacco use in any classes this year	80.3	(71.8 - 88.7)	53.4	(52.0 - 54.7)	+		
Parent or guardian has talked about the dangers of tobacco in the past year	62.1	(56.6 - 67.6)	63.9	(63.1 - 64.6)			+
Served as an active member of SWAT	14.2	(9.2 - 19.1)	4.3	(3.9 - 4.7)	+		
<u>Definitely did not think</u> that smokers have more friends	35.8	(29.6 - 42.0)	33.7	(32.9 - 34.5)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	76.0	(71.0 - 81.1)	73.7	(73.0 - 74.4)			+

Holmes County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Holmes County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	19.1	(13.7 - 24.6)	20.1	(19.5 - 20.7)			+
Current asthma, among all individuals	12.8	(8.3 - 17.3)	12.0	(11.5 - 12.5)			+
Asthma attack in past year, among individuals with lifetime asthma	11.8	(3.2 - 20.3)	18.1	(16.8 - 19.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	20.3	(15.3 - 25.3)	11.6	(11.0 - 12.1)	+		
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	15.1	(10.8 - 19.4)	16.5	(15.8 - 17.2)			+
Sufficient vigorous activity	70.1	(65.0 - 75.2)	70.1	(69.2 - 70.9)			+
Sufficient moderate activity	22.5	(18.7 - 26.2)	24.8	(24.1 - 25.4)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	48.2	(41.7 - 54.7)	44.4	(43.6 - 45.2)			+
Described themselves as slightly or very overweight	35.6	(30.2 - 41.0)	30.2	(29.4 - 30.9)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	12.9	(8.3 - 17.6)	11.4	(10.8 - 12.0)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	15.6	(11.6 - 19.6)	17.8	(17.2 - 18.5)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	5.0	(1.6 - 8.4)	7.8	(7.3 - 8.3)			+
Purposely hurt themselves without wanting to die, during the past 12 months	9.3	(5.7 - 12.9)	12.6	(12.0 - 13.2)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	20.6	(15.9 - 25.4)	19.2	(18.5 - 19.8)			+

Holmes County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Holmes County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	14.7	(9.8 - 19.6)	10.1	(9.6 - 10.6)			+
Smoked cigars on one or more of the past 30 days	10.9	(6.8 - 15.1)	11.4	(10.9 - 11.9)			+
Used smokeless tobacco on one or more of the past 30 days	14.3	(9.4 - 19.1)	5.6	(5.2 - 6.0)	+		
Used any form of tobacco on one or more of the past 30 days	25.1	(19.0 - 31.1)	17.9	(17.3 - 18.6)	+		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	54.2	(46.2 - 62.3)	58.4	(57.6 - 59.3)			+
Exposed to secondhand smoke during the past 7 days	56.4	(50.3 - 62.5)	47.6	(46.7 - 48.4)	+		
Smoking is allowed in the home	17.9	(12.9 - 22.9)	10.0	(9.5 - 10.5)	+		
Taught about tobacco use in any classes this year	43.8	(36.3 - 51.4)	33.4	(32.4 - 34.4)	+		
Parent or guardian has talked about the dangers of tobacco in the past year	43.4	(37.5 - 49.3)	50.4	(49.6 - 51.2)		+	
Served as an active member of SWAT	16.2	(11.3 - 21.1)	4.0	(3.7 - 4.3)	+		
<u>Definitely did not think</u> that smokers have more friends	33.4	(27.0 - 39.7)	30.5	(29.8 - 31.2)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	68.8	(62.0 - 75.6)	71.7	(71.0 - 72.4)			+

Holmes County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Holmes County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	18.7	(14.4 - 23.1)	20.8	(20.2 - 21.5)			+
Current asthma, among all individuals	7.7	(4.8 - 10.7)	10.6	(10.1 - 11.1)			+
Asthma attack in past year, among individuals with lifetime asthma	20.0	(10.0 - 30.0)	16.3	(15.1 - 17.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	14.5	(9.9 - 19.1)	11.1	(10.5 - 11.6)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	14.4	(9.8 - 19.0)	14.3	(13.7 - 14.9)			+
Sufficient vigorous activity	63.3	(56.6 - 69.9)	62.7	(62.0 - 63.5)			+
Sufficient moderate activity	23.8	(18.4 - 29.2)	26.1	(25.4 - 26.7)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	43.1	(36.9 - 49.2)	42.3	(41.5 - 43.1)			+
Described themselves as slightly or very overweight	32.0	(27.0 - 37.0)	29.8	(29.0 - 30.5)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	14.4	(10.4 - 18.4)	11.2	(10.5 - 11.8)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	21.4	(16.9 - 25.8)	23.1	(22.4 - 23.8)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	5.6	(2.5 - 8.8)	6.9	(6.4 - 7.3)			+
Purposely hurt themselves without wanting to die, during the past 12 months	14.5	(9.6 - 19.4)	12.7	(12.1 - 13.2)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	22.6	(17.4 - 27.9)	24.5	(23.8 - 25.2)			+

Holmes County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Holmes County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	9.3	(6.5 - 12.1)	6.1	(5.8 - 6.4)	+		
Smoked cigars on one or more of the past 30 days	6.5	(4.5 - 8.6)	6.9	(6.6 - 7.2)			+
Used smokeless tobacco on one or more of the past 30 days	9.4	(6.8 - 11.9)	3.5	(3.2 - 3.7)	+		
Used any form of tobacco on one or more of the past 30 days	17.1	(13.5 - 20.8)	11.2	(10.8 - 11.7)	+		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	62.9	(57.3 - 68.5)	65.7	(65.1 - 66.3)			+
Exposed to secondhand smoke during the past 7 days	54.7	(50.3 - 59.1)	44.1	(43.4 - 44.7)	+		
Smoking is allowed in the home	21.5	(17.6 - 25.5)	9.4	(9.0 - 9.8)	+		
Taught about tobacco use in any classes this year	63.6	(56.4 - 70.8)	43.2	(42.3 - 44.1)	+		
Parent or guardian has talked about the dangers of tobacco in the past year	54.0	(49.5 - 58.5)	57.4	(56.8 - 58.0)			+
Served as an active member of SWAT	15.3	(11.5 - 19.2)	3.9	(3.7 - 4.1)	+		
<u>Definitely did not think</u> that smokers have more friends	33.2	(28.7 - 37.7)	32.0	(31.5 - 32.6)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	72.7	(68.3 - 77.1)	72.9	(72.4 - 73.4)			+

Holmes County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Holmes County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	19.0	(15.6 - 22.5)	20.5	(20.1 - 21.0)			+
Current asthma, among all individuals	10.2	(7.5 - 12.9)	11.3	(11.0 - 11.7)			+
Asthma attack in past year, among individuals with lifetime asthma	17.1	(10.0 - 24.2)	16.9	(16.0 - 17.8)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	17.5	(13.9 - 21.2)	11.1	(10.7 - 11.5)	+		
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	14.7	(11.4 - 18.0)	15.4	(15.0 - 15.9)			+
Sufficient vigorous activity	67.3	(62.8 - 71.8)	66.5	(65.9 - 67.1)			+
Sufficient moderate activity	22.8	(19.3 - 26.3)	25.3	(24.8 - 25.8)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	46.6	(41.8 - 51.4)	43.5	(42.9 - 44.1)			+
Described themselves as slightly or very overweight	33.9	(29.9 - 37.8)	30.0	(29.5 - 30.6)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	13.3	(10.3 - 16.3)	10.9	(10.4 - 11.3)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	18.4	(15.4 - 21.5)	20.2	(19.7 - 20.7)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	4.6	(2.4 - 6.8)	6.9	(6.6 - 7.2)			+
Purposely hurt themselves without wanting to die, during the past 12 months	11.8	(8.7 - 15.0)	12.4	(12.0 - 12.8)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	21.8	(18.2 - 25.5)	22.0	(21.5 - 22.5)			+