



# Escambia County: 2014 at a Glance

Data from the 2014 Florida Youth Tobacco Survey (FYTS)



Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Escambia County %	95% CI	State %	95% CI	County Significantly Higher Than State	State Significantly Higher Than County	No Significant Difference
Smoked cigarettes on one or more of the past 30 days	5.7	(4.4-7.1)	4.3	(4.1-4.6)			+
Smoked cigars on one or more of the past 30 days	6.6	(4.9-8.3)	5.4	(5.2-5.7)			+
Used smokeless tobacco on one or more of the past 30 days	4.2	(2.9-5.4)	3.3	(3.1-3.5)			+
Used electronic cigarettes on one or more of the past 30 days	5.9	(4.3-7.6)	7.2	(6.8-7.5)			+
Used hookah on one or more of the past 30 days	4.1	(2.8-5.4)	7.1	(6.7-7.5)		+	
Used any form of tobacco on one or more of the past 30 days*	11.8	(9.7-13.9)	9.2	(8.8-9.6)	+		
Used any form of tobacco (excluding cigarettes) on one or more of the past 30 days	13.9	(11.5-16.2)	14.8	(14.2-15.3)			+
Thought using electronic cigarettes is less harmful than smoking cigarettes	67.3	(62.8-71.8)	69.4	(68.4-70.3)			+
Thought using hookah is less harmful than smoking cigarettes	39.4	(33.7-45.0)	52.8	(51.7-54.0)		+	
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	65.1	(62.3-67.8)	67.1	(66.4-67.7)			+
Exposed to secondhand smoke during the past seven days (in a room or car)	45.3	(41.8-48.7)	37.5	(36.9-38.2)	+		
Smoking is allowed in the home	14.3	(12.2-16.5)	8.4	(8.0-8.7)	+		

\*This excludes specialty tobacco products, which include bidis, kreteks, pipe tobacco, snus, e-cigarettes, flavored cigarettes, flavored cigars, and flavored smokeless tobacco.



# Escambia County: 2014 at a Glance

Data from the 2014 Florida Youth Tobacco Survey (FYTS)



Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Escambia County %	95% CI	State %	95% CI	County Significantly Higher Than State	State Significantly Higher Than County	No Significant Difference
Students who were taught about tobacco use in any classes this year	<b>34.0</b>	(30.4-37.6)	<b>38.0</b>	(37.1-39.0)			+
Parent or guardian has talked about the dangers of tobacco in the past year	<b>59.0</b>	(55.8-62.1)	<b>62.4</b>	(61.8-63.0)			+
Served as an active member of Students Working Against Tobacco (SWAT)	<b>2.4</b>	(1.6-3.2)	<b>2.8</b>	(2.7- 3.0)			+
Definitely did not think that smokers have more friends	<b>28.2</b>	(25.5-31.0)	<b>28.2</b>	(27.6-28.8)			+
Definitely did not think that smoking helps young people "look cool" or "fit in"	<b>70.2</b>	(67.8-72.7)	<b>68.8</b>	(68.2-69.3)			+
Lifetime asthma	<b>23.6</b>	(21.3-25.9)	<b>20.8</b>	(20.3-21.3)			+
Current asthma	<b>13.3</b>	(11.3-15.3)	<b>11.5</b>	(11.1-11.9)			+
Asthma attack in past year	<b>18.9</b>	(14.8-23.0)	<b>17.7</b>	(16.7-18.7)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	<b>13.7</b>	(11.3-16.0)	<b>12.2</b>	(11.8-12.7)			+
Overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex)	<b>17.0</b>	(14.5-19.4)	<b>15.9</b>	(15.4-16.4)			+
Physically active for at least 60 minutes per day during the past seven days	<b>23.8</b>	(21.2-26.5)	<b>22.7</b>	(22.2-23.3)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	<b>42.9</b>	(40.4-45.4)	<b>43.7</b>	(43.1-44.4)			+
Described themselves as slightly or very overweight	<b>31.0</b>	(28.4-33.5)	<b>30.7</b>	(30.2-31.3)			+