### SEASONAL HIGH WATER TABLE INDICATORS NON-HYDRIC APRIL 2015

David Hammonds Environmental Consultant Florida Department of Health

Division of Disease Control and Health Protection

IOTIC

<section-header><list-item><list-item><list-item><list-item><list-item><list-item><list-item>



### Soil Textures and Redox Features

- Sandy Soil- any soil with a texture of Loamy Fine Sand (LFS) or more coarse
- Loamy/Clayey Soil- means any soil with a texture of Loamy Very Fine Sand (LVFS) <u>or finer</u>

Division of Disease Control and He

IOTIC IE AI T

HEALT

Sandy	Textured	Soils
-------	----------	-------

- Very coarse sand (VCOS)
- Coarse sand (COS)
- Sand (S)
- Fine sand (FS)
- Very fine sand (VFS)
- Loamy coarse sand (LCOS)
- Loamy sand (LS)
- Loamy fine sand (LFS)
   Division of Disease Control and Health Pro
  To crotest: Control and Health Pro
  To crotest: Control and method the health of all propile in Florida through the



### Loamy/Clayey Textured Soils

- Silt loam (SIL)
- Silt (SI)
- Sandy clay loam (SCL)
- Clay loam (CL)
- Silty clay loam (SICL)
- Sandy clay (SC)
- Silty clay (SICL)

### **Seasonal High Water Tables**

Division of Disease Control and Health Pr

• Seasonal High Water Table (Seasonal High Saturation in USDA NRCS terms) is the highest level to a zone of saturation in the soil in most years. Normally persists for several weeks and normally occurs during the time of the year when the most rain falls (June-September in FL)

Division of Disease Control and Health Protect

8 HEALT



### **Apparent/Perched Water Table**

- Apparent: a thick zone of free water in the soil. It is indicated by the level at which water stands in an uncased borehole after adequate time is allowed for adjustment in the surrounding soil
- Perched: Water standing above an unsaturated zone. In some places, an upper, or perched, water table is separated from a lower one by a dry zone

Division of Disease Control and Health Pro



### **Redoximorphic Reactions**

- Also known as REDOX, which is an Oxidation/Reduction reaction.
- This is a process in which one or more substances are changed into others
- Change in color could be due to presence of water which creates the redox reaction

Division of Disease Control and H

### Redoximorphic (Redox) Features

- Caused by presence of specific minerals and their reaction to water over a specific minimum time frame
- Used to predict SHWT
- Requires certain colors, amounts, and contrast when compared with surrounding soils

Division of Disease Control and Health P

### **Redoximorphic (Redox) Features**

- Redox <u>Concentrations</u>: areas of apparent <u>accumulation</u> (more color) of Fe-Mn (Iron-Manganese) oxides, resulting in splotches of higher chroma
- Redox <u>Depletions</u>: <u>removal</u> of Fe or Mn from the soil resulting in a removal of color, leaving grayer soils

Division of Disease Control and Health Pro

14 HEALT



# Non-hydric Soil Indicator Usage The non-hydric indicators are used beginning at 12" below natural grade by below natural grade From top of natural grade to 12" the hydric soil indicators are used The hydric soil indicators will be discussed in the next presentation

IOTIC IE AI T

HEALTH





### **Non-Hydric Soil Indicators**

- Different in sandy soils than in loamy/clayey (finer textured) soils
- Must be used beginning at 12" below <u>natural soil surface</u>, cannot be used within 12" of natural soil surface
- Where part of natural soil has been removed <u>must take that into account</u>

Division of Disease Control and Health Pr

### Volume of Redoximorphic Features

The quantity (volume) of the redoximorphic features in the soil sample is important for the determination of the estimated seasonal high water tables

Division of Disease Control and Health Protect

Florid: HEALT

Florida HEALTH

### Non-hydric soils

- Redoximorphic features must be at least COMMON, but can also be MANY
- They <u>CANNOT</u> be FEW
- Need to know the required volume that redox features must occupy to count as "common" or "many"

Division of Disease Control and Health Protec and improve the health of all people in Florida through integral

### Quantity of Redoximorphic Features

Division of Disease Control and Health Prote

- Few -- less than 2% (<2%)
- Common -- 2 to 20% (2-20%)
- Many -- more than 20% (>20%)



### Redox Features in Sandy <u>Non-hydric soils</u>

- Mottles must meet following to be a Redoximorphic (redox) feature:
- Hue: 2.5YR through 10YR
- Value: ≥5
- Chroma: ≥6
- Must be at least common (2%) or many (>20%) in volume

Division of Disease Control and Health Prot

• And....

### Redox Features in Sandy Non-hydric soils

• Must have diffuse boundaries and have distinct/prominent contrast with matrix

Division of Disease Control and He

• Matrix chroma ≥3 ; any value

### NODULES AND CONCRETIONS:

- Nodules are cemented or hardened plinthite
- Concretions are similar except for the presence of visible, concentric layers around a point or line
- Both have sharp boundaries and except as specifically noted for specific HSI, do not count as a redoximorphic feature

Division of Disease Control and Health Pr

IOTIC











### Redox Features in Loamy/Clayey, Non-hydric soils

- Hue: ANY (non-gley hues)
- Value: ≥5
- Chroma: ≤2 from 12" to 1 meter (39.37")
- Chroma: ≤3 deeper than one meter
- <u>Can be diffuse mottles or be the matrix</u> <u>color</u>

Division of Disease Control and H

HEALTH















- The depth at which these low chroma colors are encountered is the estimated depth of the SHWT
- Exception to use of this color pattern as an <u>indicator:</u> where low chroma colors directly underlie a dark topsoil layer, the SHWT is at, near, or above the soil surface (this could be a hydric soil indicator)

ase Control and He

35 HEALT

Division of Dise



# Gley Charts- SHWT Indicators for <u>ALL</u> soil textures

- Hue: ANY on the Gley Charts
- Value: ≥4
- Chroma: ANY (Gley chromas all ≤2)
- Can be matrix color (commonly is)
- NOTE: These types of colors found in very wet mineral soils







13




### Contemporary vs. Relict Features

- Contemporary Soil morphological features that reflect <u>current</u> hydrologic conditions of saturation and anaerobiosis. These <u>are</u> used to determine SHWT
- Relict Soil morphological features that reflect <u>past</u> hydrologic conditions of saturation and anaerobiosis. These <u>are not</u> used to determine SHWT

Division of Disease Control and Health Protect

41 HEALT



A COUPLE OF "SPECIAL CASES" WITHIN THE LOAMY/CLAYEY GROUPING

Division of Disease Control and Health Pr

lorid

Florid: HEALT

45 HEALTH

Redoximorphic Features in Shubuta, Cowarts, Esto and Nankin Soil Series found in the Florida Panhandle

> According to the USDA NRCS, these soils were formed in loamy, clayey or loamy/clayey sediments

> > Division of Disease Control and Health Pro

- Many of the gray colors in these soils are thought to originate from parent material
- Are not a result of saturation (water movement)
- The gray colors normally mixed with redder and yellow-red and brown are not indicative of a SHWT as the color boundaries are sharp, not diffuse
- Diffuse boundaries would indicate SHWT

Division of Disease Control and Health Pro

### Shubuta, Cowarts, Esto and Nankin Soil Series

- Shubuta is a well drained soil (SHWT generally >6 feet)
- Cowarts is well drained or moderately well drained with a SHWT routinely >6 feet

Division of Disease Control and H

```
Shubuta, Cowarts, Esto and
Nankin Soil Series
Esto is a well drained soil with SHWT
routinely >6 feet
Nankin is a well drained soil with
rodeximerable indicators parmally
```

redoximorphic indicators normally occurring below 40 inches

Division of Dise

0/2015

ase Control and He

### USDA Drainage Classes (agricultural)

6 Classes common in Florida	<u>SHWT</u>			
Very Poorly Drained Poorly Drained Somewhat Poorly Drained Moderately Well Drained Well Drained	0 - 24 <u>above</u> 0 -18" below 12-30" below 24-48" below 60" or more			
Excessively Drained	>72"			
Division of Disease Control and Health Protection To protect, promote and improve the health of all people in Fordia through integrated state, county, and community efforts.				





### SOIL REMOVAL/ADDITIONS

 Where the natural soil has been altered by the action of humans (maybe animals) via soil removal and/or placement of fill material, all SHWT indicators must be judged using natural soil surface criteria accounting for what was added or removed (or moved around by animals)

Division of Disease Control and Health Prot

50 HEALTH

h of all pe

Top









# What is texture of fill and how long has it been on site?

- Try and determine the amount of time that the fill material has been on site
- May be able to use Google Earth's historic imagery feature to look back to about 1994
- Another site: Florida Aerial Photography at http://ufdc.ufl.edu/aerials

Division of Disease Control and Health Pr

54 HEALTH

h of all pe

To pro

# What is texture of fill and how long has it been on site?

- Some filled areas are much more recent
- Filled areas present additional problems
- It is advisable to conduct <u>many</u> more soil profiles in the area of the drainfield to properly view the possibly very different and complex soil properties

Division of Disease Control and Health Protection protect, promote and improve the health of all people in Florida through integrated state, county, and comm

Florid HEALT

### Anthropogenic Soils and SHWT

- Many areas in Florida have been filled for decades
- This occurred over large areas during the 1950's through the 1970's, especially in SW Florida
- One of the reasons that they were filled is because they were <u>very wet</u> and unbuildable

Division of Disease Control and Health Protecti



### Redox Features in Fill Materials

- Use the information found in Hydric Soils Technical Note #5
- Sandy Soils: 3-5 years for contemporary features to form, if all "ingredients" are present for feature formation

Division of Disease Control and Health Pr

• Loamy/Clayey soils: 8-10 years for contemporary features to form, if all "ingredients" are present for feature formation

To prot

### Redox Feature Formation in Fill Material

- Sandy Soils: 3-5 years for contemporary features to form, if all "ingredients" are present for feature formation
- Loamy/Clayey soils: 8-10 years for contemporary features to form, if all "ingredients" are present for feature formation

Division of Dise

### **Fill Materials**

ase Control and Health Pro

- Ensure the feature used to determine the SHWT was not transported with the fill material
- Feature must form based on the seasonal high water table <u>at the site</u>

Division of Disease Control and H

• It is common to misread indicators in fill

### \*\*\*NOTE\*\*\*

- A USDA NRCS soil scientist may be hesitant or possibly unwilling to make a determination regarding the estimated seasonal high water table based on the soil morphology in anthropogenic soils
- CHD personnel still required to determine the SHWT
- The CHD must always have a SHWT determination for OSTDS permitting





# <section-header><list-item><list-item><list-item><list-item><list-item>

### Stripped Matrix (Stripping)

- Used in <u>Sandy soils only (except for</u> <u>LVFS)</u>
- The following definition has been adjusted to allow for <u>non-hydric soils</u> (compare to later definition found in HSI presentation)

Division of Disease Control and Health Pro

IOTIC IE AI T

65 HEALT

- Fe/Mn oxides and/or organic matter have been stripped from the matrix
- Exposes primary base color of soil materials (e.g. values 5-6)
- Stripped areas form a *faint, diffuse splotchy* pattern of two or more colors

Division of Disease Control and Health Pro

 Stripped areas are ≥10% of the volume, rounded and approximately 1- 3 cm (0.5-1 inch) in diameter































# Seasonal High Water Table indicators in Organic Soils

- Muck muck at surface, SHWT is at or above the natural soil surface
- Mucky Mineral SHWT is at 0 6 inches below the natural soil surface
- Peat SHWT is at 0 6 inches from the natural soil surface
- See Hydric Soil Presentation for information

  Division of Disease Control and Health Protection

  To protect, protect health of a protect in Floats through theogened date, county, and community effort.

74 HEALTH

Florida HEALTH

What if there are really no redox features at all?

Barring anything else-

Division of Disease Control and Health Protection

To protect, pro



### Where no other indicators:

- The following moist colors of surface horizons (the A horizon) can be correlated with SHWT's:
- Black--0 to 9 inches (within 12")
- Very Dark Gray--about 12 inches (9-15")
- Dark Gray--greater than 15 inches
- DIG LOTS OF HOLES IF NECESSARY AND BE OBSERVANT !!!!

   Division of Disease Control and Health Production

77 HEALT

Florida HEALTH

n of all p

## Some things that do not indicate the SHWT

- Lamellae Some sandy soils contain a subsoil accumulation of clay that is distributed in fine lateral bands called lamellae
- Lamellae usually appear in contrasting colors, and are not indicative of wetness (unless the lamellae are low chroma colors)

Division of Disease Control and Health Pr













### Plinthite

- Iron-rich, highly weathered mixture of clay, quartz and other minerals
- Occurs commonly as red mottles that can be removed from the soil in one piece
- Usually platy, polygonal or reticulate patterns

Division of Disease Control and H























30

### Spodics and SHWTs

- Immokalee Soil Spodic is below 30", about 10-50" thick, <u>SHWT within 10" of</u> <u>soil surface</u>
- Pottsburg Soil Spodic is below 50" and about 24" thick, <u>SHWT is within 10" of</u> <u>soil surface</u>

Division of Disease Control and H

### **CONCLUSIONS:**

- A spodic layer <u>does not</u> have a direct relationship to the SHWT, it occurs <u>due to</u> a fluctuating water table
- The spodic layer *is not* a SHWT indicator
- The SHWT can be above, within OR below the spodic layer
- High chroma colors within the spodic layer <u>is not</u> a SHWT indicator

Division of Disease Control and Health Pr

92 HEALT

93 HEALTH

# There are approximately 80 differentiated spodosols in Florida

Only about 3 of them have seasonal high water tables found normally <u>below</u> the spodic layer

Division of Disease Control and Health Pro



 _
_

















 Bh1--20 to 24 inches; black (N 2/0) sand; weak coarse subangular blocky structure; many fine and medium roots; <u>sand grains</u> <u>coated with organic matter except for</u> <u>common fine pockets of uncoated sand</u> <u>grains</u>; very strongly acid; clear wavy boundary. (2 to 13 inches thick)

Division of Disease Control and Health F

33

 Bh2--24 to 32 inches; dark reddish brown (5YR 2/2) sand; common coarse faint vertical tongues of very dark brown (10YR 2/2) weak coarse subangular blocky structure; many fine and medium roots; <u>sand grains coated with organic</u> <u>matter</u>; very strongly acid; clear smooth boundary. (0 to 23 inches thick)

Division of Disease Control and Health Prote

IOTIC IE AI T

101 HEALT

 Bh3--32 to 36 inches; dark reddish brown (5YR 2/2) sand; weak fine granular structure; very friable; few fine roots; <u>sand grains coated with organic matter</u>; strongly acid; clear wavy boundary. (0 to 16 inches thick)

Division of Disease Control and Health Pro

A word about Spodic Soils What if you don't see redox features??

### Other indicators of the SHWT

- Thickness and color of the surface layer ("A" horizon), and stripped matrix, if present
- Note that high chroma colors <u>WITHIN</u> the spodic IS NOT a SHWT indicator. Stripped matrix can occur within the spodic and can be used

Division of Disease Control and Health Pr

IE AL T

105 HEALTH

```
Spodosol with SHWT below the Spodic Layer – KUREB Series
```

- Kureb –Rapid permeability. Depth to seasonal high water table is more than 6 feet during most of the year
- A--0 to 3 inches; dark gray (10YR 4/1) sand; single grained; loose; organic matter and quartz grains have salt and pepper appearance; many fine and large roots; neutral; clear wavy boundary (2 to 5 inches thick)

Division of Disease Control and Health Prot

- E--3 to 26 inches; light gray (10YR 7/1) sand; single grained; loose few large roots; neutral; clear irregular boundary. (4 to 45 inches thick)
- C/Bh--26 to 51 inches; brownish yellow (10YR 6/6) sand; single grained; loose; few tongues of light gray (10YR 7/1) extend from above horizon; dark brown (7.5YR 3/4) and few bands and bodies (Bh) of dark reddish brown (5YR 3/2); (continued next slide)

Division of Disease Control and Health Protect

To protect, pr

 (C/Bh continued) bands are intermittent at horizon contact and vertically along walls of tongues; many clean and coated sand grains; neutral; gradual wavy boundary. (4 to 46 inches thick)

 C--51 to 89 inches; pale brown (10YR 6/3) sand; single grained; loose, slightly acid

		-
Division of Disease Control and Health Protection		
To protect, promote and improve the health of all people in Florida through integrated state, county, and commu	nity efforts.	clorid a
4/29/2015	106	HEALTH









