HIV/AIDS Epidemiology Partnership 2a

Bay, Calhoun, Gulf, Holmes, Jackson and Washington Counties

Excluding Dept. of Corrections

Created: 12/03/14

Revision: 10/14/15

Florida Department of Health
HIV/AIDS Section
Annual data trends as of 12/31/2014
Living (Prevalence) data as of 06/30/2015

HIV and AIDS Case Data

- AIDS Cases became reportable in Florida in 1981.
- HIV (not AIDS) became reportable in Florida on July 1, 1997.
- HIV Infection reporting represents newly Adult HIV Infection Cases, regardless of AIDS status at time of report, that were previously reported.
- AIDS cases and HIV infection cases by year of report are NOT mutually exclusive and CANNOT be added together.
- Frozen databases of year-end data are generated at the end of each calendar year. These are the same data used for Florida CHARTS and all grant-related data where annual data are included.
- HIV prevalence data are generated later in the year, usually in July, when most of the “expected” death data are complete.

HIV and AIDS Case Data (con't)

- Adult cases represent ages 13 and older, pediatric cases are those under the age of 13. For data by year, the age is by age of diagnosis. For living data, the age is by current age at the end of the most recent calendar year, regardless of age at diagnosis.
- Unless otherwise noted, whites are non-Hispanic and blacks are non-Hispanic.
- Unless otherwise noted. Area and county data will exclude DOC cases.

Snapshot of Persons Reported with HIV Disease, 2014, Partnership 2a

Total Population, 2014*

298,037

*2014 estimate is provisional

HIV Infection and AIDS Cases Reported in 2014*

	Adults (Age 13+)	Pediatrics (Age <13)	TOTAL
HIV Infection Cases	**49**	**0**	49
AIDS Cases	20	0	20

*HIV infection cases and AIDS cases by year of report are NOT mutually exclusive and CANNOT be added together.

Cumulative HIV/AIDS Cases Reported 1981-2014

	Adults (Age 13+)	Pediatrics (Age<13)	TOTAL
HIV (not AIDS) Cases**	**292**	**0**	292
AIDS Cases	748	1	749
Total	1040	1	1041

**HIV (not AIDS) cases were NOT reportable until 07/1997

Persons Living with HIV Disease through 2014, as of 06/30/2015: 867

HIV Infection Cases and Rates*, by Year of Report, 2005-2014, Partnership 2a



Year	Rate
05	17.1
06	17.2
07	11.4
08	14.6
09	11.9
10	11.9
11	8.8
12	10.8
13	17.2
14	16.4

Note: Enhanced laboratory reporting (ELR) laws in 2006 and the expansion of ELR in 2007 led to an artificial peak in newly reported cases of HIV infection in 2008. This was followed by a general decline in reported cases through 2012. Another surge in the expansion of ELR in 2012 was followed by another increase in newly reported cases of HIV infection in 2013. In 2014, HIV infection cases decreased by 4% from the previous year. In contrast, the state observed a 12% incline in HIV infection cases during the same time period.

***Source: Population estimates are provided by Florida CHARTS as of 7/9/2015. Rates are expressed as per 100,000 population.**

AIDS Cases and Rates*
by Year of Report, 2005-2014, Partnership 2a



Year	Rate
05	12.8
06	10.2
07	7.2
08	7.1
09	10.9
10	7.1
11	5.4
12	11.2
13	7.8
14	6.7

Electronic laboratory reporting delays in late 2007 decreased cases in that year. The expansion of electronic lab reporting increased the timeliness of reporting, which contributed to the artificial spike in 2009 followed by the artificial dip in 2010. Another surge in the expansion of ELR in 2012 was followed by another increase in newly reported cases of AIDS. AIDS cases in 2014 dropped by 13% from the previous year. This is lower than the 15% decline observed by the state during the same time period.

***Source: Population estimates are provided by Florida CHARTS as of 7/9/2015. Rates are expressed as per 100,000 population.**

Adult HIV Infection Cases, by Sex and Year of Report, 2005-2014, Partnership 2a



Note: Recent trends in HIV transmission are best described by the HIV case data. The relative increases in male HIV infection cases might be attributed to proportional increases in HIV transmission among men who have sex with men (MSM), which may influence future AIDS trends. *The male-to-female ratio is the number of cases among males divided by the number of cases among females.

Adult AIDS Cases, by Sex and Year of Report, 2005-2014, Partnership 2a



Note: AIDS cases tend to represent HIV transmission that occurred many years ago. The relative increases in male cases reflect the changing face of the AIDS epidemic over time. *The male-to-female ratio is the number of cases among males divided by the number of cases among females.

Adult AIDS and HIV Infection Cases by Sex, Reported in 2014, Partnership 2a



Note: Partnership 2a's Adult Population is: 52% Male and 48% Female.

Adult HIV Infection Cases by Race/Ethnicity and Year of Report, 2005-2014, Partnership 2a



Note: HIV case reporting, implemented in mid-1997, reflects more recent trends in the epidemic with respect to the distribution of cases by race/ethnicity. Over the past ten years, the proportion of HIV infection cases among blacks and whites fluctuated over time, oftentimes crossing paths. From 2005 to 2014, the proportion of HIV infection cases among Hispanics decreased by 9 percentage points. In contrast, the proportion of HIV infection cases among blacks and whites increased by 3 and 6 percentage points, respectively, during the same time period. Other races represent less than 4% of the cases and are not included.

Florida HEALTH

Adult AIDS Cases by Race/Ethnicity and Year of Report, 2005-2014, Partnership 2a



Factors Affecting Disparities

-Late diagnosis of HIV.

-Access to/ acceptance of care.

-Delayed prevention messages.

-Stigma.

-Non-HIV STD's in the community.

-Prevalence of injection drug use.

-Complex matrix of factors related to socioeconomic status

Note: In 2014, blacks accounted for 35% of adult AIDS cases, but only 14% of the population. Although the proportion of AIDS cases among blacks and whites fluctuated over time, oftentimes crossing paths, the proportion of cases among blacks decreased by 22 percentage points from 2005 to 2014. Likewise, the proportion of AIDS cases among Hispanics decreased 9 percentage points; whereas cases among whites increased by 34 percentage points during the same time period. Other races represent less than 6% of the cases and are not included.

Florida HEALTH

Adult Male HIV Infection Cases by Race/Ethnicity and Year of Report, 2005-2014, Partnership 2a



Note: Over the years, the proportion of male HIV infection cases among whites and blacks has shifted up and down, oftentimes crossing paths. White men continue to represent the majority (>50%) of male HIV infection cases for most of the years. From 2005 to 2014, the proportion of HIV cases among white and black males increased by 4 and 7 percentage points, respectively. In contrast, the proportion of HIV cases among Hispanic males decreased by 11 percentage points during the same time period. Other races represent less than 5% of the cases and are not included.

Adult Female HIV Infection Cases by Race/Ethnicity and Year of Report, 2005-2014, Partnership 2a



Note: HIV case disparities are more evident among women than men. For the past ten years, black women represented 50% or more of the cases for most years. From 2005 to 2014, the proportion of HIV infection cases among both black and white females decreased by 4 percentage points, respectively, whereas HIV cases among Hispanic females increased by 9 percentage points during the same time period. Other races represent less than 5% of the cases and are not included.

Adult HIV Infection Case Rates* by Sex and Race/Ethnicity, Reported in 2014, Partnership 2a



Rate Ratios:

MALES

Black:White, 3.1:1

Hispanic:White, 0.6:1

FEMALES

Black:White, 9.9:1

Hispanic:White, 21.0:1

Note: Among black males, the HIV infection case rate is 3 times higher than the rate among white males. Among black females, the HIV case rate is nearly 10 times higher than the rate among white females. Among Hispanic females, the HIV case rate is 21-fold greater than the rate among white females. In contrast, the HIV case rate is lower among Hispanic males than the rate among white males.

***Source: Population estimates are provided by Florida CHARTS as of 7/9/2015.**

Florida HEALTH

Adult AIDS Case Rates* by Sex and Race/Ethnicity, Reported in 2014, Partnership 2a



Rate Ratios:

MALES
Black:White, 2.4:1
Hispanic:White, 0.0:1

FEMALES
Black:White, 4.4:1
Hispanic:White, 0.0:1

Note: Among black males, the AIDS case rate is 2 times higher than the rate among white males. Similarly, among black females, the AIDS case rate is 4 times higher than the rate among white females.
***Source: Partnership 2a Population estimates are provided by Florida CHARTS as of 7/9/2015.**

Adult HIV and AIDS Cases Reported in 2014 and Population Data, by Race/Ethnicity, Partnership 2a



Note: In this snapshot for 2014, blacks are over-represented among the HIV and AIDS cases, accounting for 39% of adult HIV cases and 35% of adult AIDS cases, but only 14% of the adult population. A group is disproportionately impacted to the extent that the percentage of cases exceeds the percentage of the population.
***Source: Population estimates are provided by Florida CHARTS as of 7/9/2015.**
****Other includes Asian/Pacific Islanders, Native Alaskans/American Indians and mixed races.**

Adult HIV Infection Cases, by Age Group at Diagnosis, and Year of Report, 2005–2014, Partnership 2a



Note: From 2005 to 2014, the proportion of adult HIV infection cases among those aged 20-29 and those aged 40-49 increased by 6 and 3 percentage points, respectively.

Adult HIV Infection Cases, by Sex and Age Group at Diagnosis, Reported in 2014, Partnership 2a



Note: HIV infection cases tend to reflect more recent transmission than AIDS cases, and thus present a more current picture of the epidemic. With regard to the age group with the highest percent of HIV infection cases, recent estimates show that among males, 33% of HIV infection cases occur among those aged 30-39, similarly among females, 86% of HIV infection cases occur among those aged 40-49.

Definitions of Mode of Exposure Categories

- **MSM =** Men who have sex with men or Male-to-male sexual contact with person with HIV/AIDS or known HIV risk
- **IDU =** Injection Drug User
- **MSM/IDU =** Men who have sex with men or Male-to-male sexual contact & Injection Drug User
- **Heterosexual =** Heterosexual contact with person with HIV/AIDS or known HIV risk
- **OTHER** = includes hemophilia, transfusion, perinatal, other pediatric risks and other confirmed risks.
- **NIR =** Cases reported with No Identified Risk
- **Redistribution of NIRs =** This illustrates the effect of statistically assigning (redistributing) the NIRs to recognized exposure (risk) categories by applying the proportions of historically reclassified NIRs to the unresolved NIRs.

Florida HEALTH

Adult Male HIV Infection Cases, by Mode of Exposure and Year of Report, 2005–2014, Partnership 2a



Note: NIRs redistributed. Among male HIV cases in Partnership 2a, men who have sex with men (MSM) remains as the primary mode of exposure for most years.

Adult Female HIV Infection Cases by Exposure Category and Year of Report, 2005-2014, Partnership 2a



Note: NIRs redistributed. The heterosexual risk continues to be the dominant mode of exposure among females.

Adult Male HIV Infection and AIDS Cases, by Mode of Exposure, Reported in 2014, Partnership 2a



Note: NIRs redistributed. Among the male HIV infection and AIDS cases reported in 2014, men who have sex with men (MSM) was the most common risk factor (81% for HIV and 73% for AIDS), followed by cases with a heterosexual risk (17% for HIV and 14% for AIDS).

Adult Female HIV Infection and AIDS Cases, by Mode of Exposure, Reported in 2014, Partnership 2a



Note: NIRs redistributed. Among the female HIV infection and AIDS cases reported for 2014, heterosexual contact was the highest risk (100% for HIV and 100% for AIDS).

Cases Living with HIV Disease

Unless otherwise noted, data in the following slides represent persons living with HIV/AIDS (PLWHAs), who were living in Florida (regardless where diagnosed) through the most recent calendar year. Living data are also referred to as prevalence cases or living with HIV disease.

Florida HEALTH

Adults Living with HIV Disease
By Zip Code, Diagnosed through 2014, Partnership 2a

Total Adult Living HIV/AIDS Cases

- 0
- 1 - 5
- 5 - 10
- Over 10



N= 866

NIRs are not redistributed.
Excludes DOC, homeless, and cases with unknown zips.
Data as of 06/30/2015

Men who have Sex with Men (MSM)*
Living with HIV Disease
By Zip Code, Diagnosed through 2014, Partnership 2a

Presumed Living MSM HIV/AIDS Cases

- 0
- 1 - 5
- 5 - 10
- Over 10



N=438

NIRs are not redistributed.
Excludes DOC, homeless, and cases with unknown zips.
***Includes MSM/IDU cases.**
Data as of 06/30/2015

Injection Drug Users (IDUs)*
Living with HIV Disease
By Zip Code, Diagnosed through 2014, Partnership 2a

Presumed Living IDU HIV/AIDS Cases

- 0
- 1 - 5
- 5 - 10
- Over 10



N=131

NIRs are not redistributed.
Excludes DOC, homeless, and cases with unknown zips.
***Includes MSM/IDU cases.**
Data as of 06/30/2015

Florida HEALTH

Adult Heterosexuals Living with HIV Disease
By Zip Code, Diagnosed through 2014, Partnership 2a

Presumed Living Heterosexual HIV/AIDS Cases

- 0
- 1 - 5
- 5 - 10
- Over 10



N=336

NIRs are not redistributed.
Excludes DOC, homeless, and cases with unknown zips.
Data as of 06/30/2015

Adults Living with HIV Disease By Zip Code and Race/Ethnicity, Diagnosed through 2014, Partnership 2a

1 Dot = 3 cases
Dots are randomly placed within zip codes.

- **Hispanic**
- **Black, not-Hispanic**
- **White, not-Hispanic**

N=849



Total includes all races, some which are not on map.
Excludes DOC, homeless, and cases with unknown zips.
Data as of 06/30/2015

Adults Living with HIV Disease By Zip Code and Sex, Diagnosed through 2014, Partnership 2a

1 Dot = 3 cases
Dots are randomly placed within zip codes.

- **Male**
- **Female**

N=866



Excludes DOC, homeless, and cases with unknown zips.
Data as of 06/30/2015

Adults Living with HIV Disease, by Sex and Race/Ethnicity Diagnosed through 2014, Partnership 2a



Note: Among adult males living with HIV disease, whites represent the race most affected (47%). Among adult females, blacks represent the race most affected (61%).
*Other includes Asian/Pacific Islanders, Native Alaskans/American Indians and Multi-racial individuals.

Case Rates* of Adults Living with HIV Disease, by Sex and Race/Ethnicity, Diagnosed through 2014, Partnership 2a



RATE RATIOS:

MALES
Blacks:Whites, 4.5:1
Hispanics:Whites, 2.2:1

FEMALES
Black:Whites, 13.4:1
Hispanics:Whites, 4.5:1

Note: Among black males living with HIV disease Diagnosed through 2014, the case rate is nearly 5 times higher than the rate among white males. Among black females living with HIV disease, the case rate is 13 times higher than the rate among white females. Hispanic male and female rates are higher than the rates among their white counterparts. Data excludes Department of Corrections cases.

***Source: Population estimates are provided by Florida CHARTS as of 7/9/2015**

****Other includes Asian/Pacific Islanders, Native Alaskans/American Indians and Multi-racial individuals.**

Adults Living with HIV Disease, by Sex and Current Age Group, Diagnosed through 2014, Partnership 2a

Males
N=665



Females
N=202



Note: Males living with HIV disease have a higher proportion of cases who are currently 40 years of age or older (69%), compared with females who are currently 40 years of age or older living with HIV disease (63%).

Adult Males Living with HIV Disease by Race/Ethnicity and Mode of Exposure Diagnosed through 2014, Partnership 2a



Note: NIRs redistributed. Male to male sexual contact (MSM) represents the highest risk for all races. White males have the smallest proportion of heterosexual contact risk.
* Other includes hemophilia, transfusion, perinatal, other pediatric risks and other confirmed risks.

Adult Females Living with HIV Disease by Race/Ethnicity and Mode of Exposure Diagnosed through 2014, Partnership 2a



Note: Heterosexual contact is the majority risk for all races. However, Hispanics have the highest proportion of IDU risk.

*** Other includes hemophilia, transfusion, perinatal, other pediatric risks and other confirmed risks.**

Annual Prevalence of Adults Living with HIV Disease, 1995-2014, Partnership 2a



Note: As a result of declining deaths, annual HIV/AIDS diagnoses have exceeded deaths since 1995, and the number of persons reported with HIV/AIDS who are presumed to be alive have been increasing. Since 1995, the number of persons reported living with HIV/AIDS have increased over 650%. In 2014, the prevalence increased by 7% from the previous year.
Note: These data represent adults living with HIV disease diagnosed in Florida regardless of their current residence.

Partnership 2a'sTop-9 Priority Populations in 2014 for Primary and Secondary HIV Prevention Based on Persons Living with HIV Disease

1. White Men who have sex with Men
2. Black Heterosexual men and women
3. Black Men who have sex with Men
4. White Heterosexual men and women
5. White Injection Drug User
6. Hispanic Heterosexual men and women
7. Hispanic Men who have sex with Men
8. Black Injection Drug User
9. Hispanic Injection Drug User

This final ranking is a result of ranking 9 race/risk groups among those newly reported in eHARS with HIV disease in Partnership 2a from the 3 most recent years, plus ranking these same 9 race/risk groups from all persons who were reported and living with HIV disease in eHARS in Partnership 1 through the most recent calendar year. The two ranks were then weighted and combined resulting in the final rank.

Florida HEALTH

Number and Percentage of Persons Diagnosed and Living with HIV (PLWH) Engaged in Selected Stages of the Continuum of HIV Care Partnership 2a (excl. DOC), 2014



- 82% of those diagnosed with HIV in 2014 had documented HIV-related care within 3 months of diagnosis
- 82% of PLWH in care had a suppressed viral load in 2014

(1) **HIV Diagnosed:** Persons diagnosed and living with HIV (PLWH) in Florida through the end of 2014.
(2) **Ever in Care**: PLWH with at least 1 documented viral load (VL) or CD4 lab, medical visit or prescription since HIV diagnosis.
(3) **In Care**: PLWH with at least 1 documented VL or CD4 lab, medical visit or prescription in 2014.
Retained in Care: PLWH with 2 or more documented VL or CD4 labs, medical visits or prescriptions (at least 3 months apart) in 2014.
(4) **On ART**: Estimated PLWH on antiretroviral therapy (ART) in 2014 (estimated from 2013 FL MMP data).
(5) **Suppressed Viral Load**: PLWH with a suppressed VL (<200 copies/mL) on last VL in 2014.

Florida HEALTH

Resident Deaths due to HIV Disease by Year of Death, 1995-2014, Partnership 2a



Race/Ethnicity	2014	
	No.	rate
White Male	6	5.2
White Female	-	0.0
Black Male	4	16.7
Black Female	2	11.1
Hispanic Male	-	0.0
Hispanic Female	-	0.0
Other*	-	0.0
TOTAL	12	4.0

Source: Florida Department of Health, Bureau of Vital Statistics, Death Certificates (as of 05/16/2014). Population data are provided by Florida CHARTS as of 7/9/2015.

* Other includes Asian/Pacific Islanders, Native Alaskans/American Indians and Multi-racial individuals.

Some Useful Links

- CDC HIV/AIDS Surveillance Reports
  (State and Metro Data):
  http://www.cdc.gov/hiv/stats/hasrlink.htm

- MMWR (Special Articles on Diseases, Including HIV/AIDS):
  http://www.cdc.gov/mmwr/

- U.S. Census Data (Available by State, County):
  http://www.census.gov

- Partnership 2a, Dept. of Health, HIV/AIDS & Hepatitis Section Website (Slide sets, Facts Sheets, Monthly Surveillance Report, Counseling & Testing Data, etc.):
  http://www.floridahealth.gov/diseases-and-conditions/aids/surveillance/index.html

Florida HEALTH

"The reason for collecting, analyzing and disseminating information on a disease is to control that disease. Collection and analysis should not be allowed to consume resources if action does not follow."

--Foege WH et al. Int. J of Epidemiology 1976; 5:29-37

Partnership 2a Surveillance Contact

Darrell Barfield
Leon County Health Department
Phone: 850-606-8263
Email: Darrell.Barfield@flhealth.gov

Rick Vitale, HIV/AIDS Program Coordinator
Phone: 850-872-4455 ext. 1184
Email: Rick.Vitale@flhealth.gov

For Florida HIV/AIDS Surveillance Data

Contact: (850) 245-4444

Lorene Maddox, MPH	**Ext. 2613**
Tracina Bush, BSW	**Ext. 2612**
Madgene Moise, MPH	**Ext. 2373**

Visit Florida's internet site for:

Monthly Surveillance Reports

Slide Sets and Fact Sheets

Annual Reports and Epi Profiles

http://www.floridahealth.gov/diseases-and-conditions/aids/surveillance/index.html

Visit CDC's HIV/AIDS internet site for:

Surveillance Reports, fact sheets and slide sets

http://www.cdc.gov/hiv/topics/surveillance/resources/reports/index.htm