

INFANT MORTALITY AND LOW BIRTH WEIGHT RATES COMPARED TO EXPECTED RATES BY HEALTHY START COALITION AREA 2005

**By: Daniel Thompson, M.P.H.; Carol Graham, Ph.D.; Cheryl Clark, M.P.H., R.H.I.A.;
Angel Watson, M.P.H., R.H.I.A.**

**Florida Department of Health, Division of Family Health Services
Bureau of Family and Community Health**

October 30, 2006

Introduction

Infant mortality and birth weight statistics are used extensively in public health. These statistics are especially useful because of their relevance as maternal and child health indicators and because of their ease of availability and relatively high level of completeness.

The purpose of this analysis is to identify geographic areas in the state where low birth weight (LBW) rates and infant mortality (IM) rates are statistically, significantly higher than would be expected considering the unique demographics of each area. These areas should then be the focus of further, more detailed analyses to determine the reasons for the high rates and to develop intervention strategies for improving the outcomes.

IM and LBW rates vary in relation to the demographic characteristics and the variation in rates across the Healthy Start coalition areas is due in part to the unique demographic characteristics of the local populations. In this analysis, adjustments are made to account for the differences in demographic characteristics.

Three demographic variables are used in calculating the adjusted and expected statistics. These are maternal race, marital status, and education. These variables are used because they are known to be associated with risk of LBW and IM, and because public health interventions are not designed to influence these characteristics in the prenatal or infancy period. In an analysis (data not shown) of Florida resident births in 2001, linked to infant deaths, risk of infant death was found to be 133 percent (133%) higher for maternal race Black, 89 percent (89%) higher for unmarried maternal marital status, and 41 percent (41%) higher for maternal education less than high school. In the same analysis, risk of LBW was found to be 82 percent (82%) higher for maternal race Black, 44 percent (44%) higher for unmarried maternal marital status, and 22 percent (22%) higher for maternal education less than high school. These results were all statistically significant at the 0.05 alpha level. Maternal characteristics such as maternal age and smoking status are not used in the adjustment because there are public health efforts directed at changing these factors and adjusting for them would eliminate differences due to these factors. For example, if an area has an actual LBW percentage significantly lower than the expected LBW percentage, the difference could be due to the extraordinary success of a smoking cessation program in the area. If adjustments were made for smoking status, this difference would not be apparent. Maternal age can be influenced by reducing teen births, and by the same logic, adjustments are not made for maternal age.

IM and LBW rates also reflect random variation. In this analysis, statistical methods are used to separate the random variation from the non-random variation, so rates that are significantly high are most likely a result of non-random influences. Likewise, rates that are higher than expected,

but not significantly high, are likely to be the result of random variation and are said to be within the range of normal variation.

Methods

The data used in this analysis were extracted from the birth records for residents of Florida born in calendar years 2004 and 2005. Births were classified as LBW if the birth weight on the birth record was in the range of 1 to 2499 grams. Three demographic variables were used in this analysis: mother’s race, marital status, and education. These are recorded on the birth record, and for the purposes of this analysis, two categories were used for each variable. Mother’s race was classified as Black or non-Black, marital status was classified as married or not married, and mother’s education was classified as 12th grade or higher completed or less than 12th grade completed. The three variables were then used to classify the births into eight mutually exclusive categories. Birth records with unknown values for any of the three variables were placed in a ninth category. There were roughly 3000 birth records in the ninth category (about 1.3% of the resident births). The nine categories are as follows:

<u>Mother’s Category</u>	<u>Mother’s Race</u>	<u>Mother’s Marital Status</u>	<u>Education</u>
1	Non-Black	Married	High School or More
2	Non-Black	Married	Less than High School
3	Non-Black	Not Married	High School or More
4	Non-Black	Not Married	Less than High School
5	Black	Married	High School or More
6	Black	Married	Less than High School
7	Black	Not Married	High School or More
8	Black	Not Married	Less than High School
9*	Unknown	Unknown	Unknown

* This includes records with unknown values in any of the three categories.

Calculating Expected Rates:

Using this classification, the category-specific rates were calculated from the 2003 (the latest year for complete matched birth and infant death data) statewide totals, and these rates were used with the 2004 births in each Healthy Start coalition area to calculate the expected LBW births and infant deaths. In this way, the expected statistics are adjusted for the three demographic characteristics and then used to calculate the adjusted rates. The term for this adjustment technique is “indirect adjustment.”

In March of 2004, the recording of maternal race on the birth record was changed so that more than one race can be selected. For the purposes of this analysis, births where the only maternal race recorded was Black were classified as Black and all others were classified as non-Black. There were 49,258 births with maternal race Black and 47,957 (97.4%) of these recorded no other race for maternal race.

For example, if an area existed where all the births were in category 1, then the expected statistics for the area would be the same as the statewide statistics for category 1. Another area

might have had births that were all in category 8. For this area, the expected statistics would be the same as the statewide statistics for category 8. These two hypothetical areas would have different expected statistics because they have populations with different demographic characteristics. If both areas had actual rates equal to the expected rates, they would be considered equal regarding the rates. Stated differently, both areas are doing equally well at preventing IM and LBW, considering their different demographic characteristics.

The correlation between actual IM and LBW across the areas was also assessed. The normal approximation to the binomial distribution formulas were used for statistical testing in areas where the number of infant deaths or low birth weight infants were above 50. When these were 50 or below, the Poisson formula was used.

Results

The results of this analysis are shown in the following tables where actual statistics are compared to expected statistics. The expected statistics are adjusted for the demographic characteristics in each area, as described above. Areas with statistically, significantly high actual statistics are indicated in the tables with an “H” and “L” indicates significantly low actual statistics.

There is a borderline statistically significant correlation between areas with high LBW percentages and areas with high infant death rates. This means areas with high LBW percentages tend to have high infant death rates, and areas with low LBW percentages tend to have low infant death rates. The correlation coefficient based on the ranks of the p values across coalition areas is 0.33 with an associated p value of 0.06.

Discussion

This analysis should be considered a preliminary step in the continuing endeavor to reduce risk of infant death and low birth weight in Florida. The rationale is to use the results of this analysis to focus further analysis and efforts on the areas where the risks are significantly high. Since adjustments were used to account for the differing demographic composition in each area, further analysis would focus on other factors such as smoking rates and mother’s age at birth. Unique factors in each county contribute to infant deaths and low birth weight. Local area analysis of factors associated with these outcomes should be undertaken to better understand the reasons for higher than expected rates. The process becomes much more complicated at this point, and a separate analysis should be done for each area of concern. Finally, although demographic adjustment is useful for analyzing additional influencing variables, it remains critical to continue efforts to address issues such as racial disparity in health outcomes.

**2005 FLORIDA ACTUAL INFANT DEATH RATES PER 1000 BIRTHS
COMPARED TO EXPECTED¹ RATES PER 1000 BIRTHS**

	2005		2005		H=Actual Rate Signif.Higher ²	L=Actual Rate Signif.Lower ²
	Expected ¹	Actual	Expected ¹	Actual		
Healthy Start Coalition	2005 Births	2005 Infant Deaths	2005 Infant Deaths	2005 Expected Infant Death Rate Per 1000 Births	2005 Actual Infant Death Rate Per 1000 Births	
Bay, Franklin, Gulf Healthy Start Coalition	2,640	17.7	16	6.70	6.06	
Broward Healthy Start Coalition, Inc.	23,127	202.6	144	8.76	6.23	L
Capital Area Healthy Start Coalition, Inc.	3,393	26.9	28	7.93	8.25	
Central Healthy Start, Inc.	6,241	41.8	50	6.70	8.01	
Charlotte County Healthy Start Coalition, Inc.	1,084	6.4	8	5.90	7.38	
Chipola Healthy Start Coalition, Inc.	1,329	9.1	16	6.85	12.04	H
Desoto	505	3.6	2	7.13	3.96	
Escambia County Healthy Start Coalition, Inc.	4,237	32.0	30	7.55	7.08	
Florida Keys Healthy Start Coalition, Inc.	768	4.7	1	6.12	1.30	
Gadsden Citizens for Healthy Babies Inc.	741	7.6	10	10.26	13.50	
Miami-Dade	32,365	237.3	176	7.33	5.44	L
Healthy Start Community Coalition of Okaloosa and Walton Counties, Inc.	3,342	20.1	20	6.01	5.98	
Healthy Start of North Central Florida, Inc.	10,123	71.9	91	7.10	8.99	H
Healthy Start Coalition of Sarasota County, Inc.	2,997	17.8	16	5.94	5.34	
Healthy Start Coalition of Hardee / Highlands / Polk Counties, Inc.	9,181	65.0	73	7.08	7.95	
Healthy Start Coalition of Hillsborough County, Inc.	16,753	118.1	149	7.05	8.89	H
Healthy Start Coalition of Jefferson / Madison / Taylor Counties, Inc.	664	5.6	6	8.43	9.04	
Healthy Start Coalition of Manatee County, Inc.	3,809	25.8	26	6.77	6.83	
Maternal Child Family Health Alliance of Palm Beach County, Inc.	15,160	111.9	96	7.38	6.33	
Healthy Start Coalition of Pasco County, Inc.	4,753	28.1	34	5.91	7.15	
Healthy Start Coalition of Pinellas County, Inc.	9,065	61.7	77	6.81	8.49	H
Healthy Start Coalition of Santa Rosa County, Inc.	1,746	9.3	9	5.33	5.15	
Healthy Start Coalition of Southwest Florida, Inc.	11,607	77.4	61	6.67	5.26	L
Healthy Start Coalition of St. Lucie County, Inc.	3,003	22.0	19	7.33	6.33	
Indian River County Healthy Start Coalition, Inc.	1,360	9.0	7	6.62	5.15	
Martin County Healthy Start Coalition, Inc.	1,340	8.5	11	6.34	8.21	
Northeast Florida Healthy Start Coalition, Inc.	18,154	134.8	189	7.43	10.41	H
Okeechobee County Family Health / Healthy Start Coalition, Inc.	587	3.9	1	6.64	1.70	
Orange County Healthy Start Coalition, Inc.	16,556	120.2	128	7.26	7.73	
Prenatal and Infant Health Care Coalition of Brevard County, Inc.	5,387	34.5	37	6.40	6.87	
Seminole County Healthy Start Coalition, Inc.	4,786	29.8	37	6.23	7.73	
The Healthy Start Prenatal & Infant Coalition of Flagler and Volusia Counties, Inc.	5,782	37.5	28	6.49	4.84	
The Healthy Start Coalition of Osceola County, Inc.	3,593	22.4	29	6.23	8.07	
TOTAL	226,178	1,625	1,625	7.18	7.18	

¹ The expected number of infant deaths is calculated based on the maternal race, marital status and education characteristics of the births in each county

² The significance level used is .05

**2005 FLORIDA ACTUAL LOW BIRTH WEIGHT¹ PERCENTAGES
COMPARED TO EXPECTED² PERCENTAGES**

	2005 Births	2005	2005	2005	2005	H=Actual Rate Signif.Higher ³ L=Actual Rate Signif.Lower ³ Than Expected
		Expected ¹ LBW Births	Actual LBW Births	Expected LBW Percent	Actual LBW Percent	
Healthy Start Coalition						
Bay, Franklin, Gulf Healthy Start Coalition	2,640	220.0	238	8.33%	9.02%	
Broward Healthy Start Coalition, Inc.	23,127	2202.2	2,142	9.52%	9.26%	
Capital Area Healthy Start Coalition, Inc.	3,393	321.2	326	9.47%	9.61%	
Central Healthy Start, Inc.	6,241	514.7	486	8.25%	7.79%	
Charlotte County Healthy Start Coalition, Inc.	1,084	85.0	70	7.84%	6.46%	L
Chipola Healthy Start Coalition, Inc.	1,329	113.9	137	8.57%	10.31%	H
Desoto	505	43.0	36	8.51%	7.13%	
Escambia County Healthy Start Coalition, Inc.	4,237	389.4	431	9.19%	10.17%	H
Florida Keys Healthy Start Coalition, Inc.	768	61.3	61	7.98%	7.94%	
Gadsden Citizens for Healthy Babies Inc.	741	83.1	95	11.21%	12.82%	
Miami-Dade	32,365	2893.6	2,918	8.94%	9.02%	
Healthy Start Community Coalition of Okaloosa and Walton Counties, Inc.	3,342	266.2	270	7.97%	8.08%	
Healthy Start of North Central Florida, Inc.	10,123	889.7	919	8.79%	9.08%	
Healthy Start Coalition of Sarasota County, Inc.	2,997	237.1	225	7.91%	7.51%	
Healthy Start Coalition of Hardee / Highlands / Polk Counties, Inc.	9,181	797.9	774	8.69%	8.43%	
Healthy Start Coalition of Hillsborough County, Inc	16,753	1449.4	1,504	8.65%	8.98%	
Healthy Start Coalition of Jefferson / Madison / Taylor Counties, Inc.	664	65.1	63	9.80%	9.49%	
Healthy Start Coalition of Manatee County, Inc.	3,809	321.4	291	8.44%	7.64%	L
Maternal Child Family Health Alliance of Palm Beach County, Inc.	15,160	1358.2	1,395	8.96%	9.20%	
Healthy Start Coalition of Pasco County, Inc.	4,753	368.3	389	7.75%	8.18%	
Healthy Start Coalition of Pinellas County, Inc.	9,065	764.4	763	8.43%	8.42%	
Healthy Start Coalition of Santa Rosa County, Inc	1,746	131.2	129	7.51%	7.39%	
Healthy Start Coalition of Southwest Florida, Inc.	11,607	970.2	891	8.36%	7.68%	L
Healthy Start Coalition of St. Lucie County, Inc.	3,003	269.0	240	8.96%	7.99%	L
Indian River County Healthy Start Coalition, Inc.	1,360	113.8	107	8.37%	7.87%	
Martin County Healthy Start Coalition, Inc.	1,340	108.7	117	8.11%	8.73%	
Northeast Florida Healthy Start Coalition, Inc.	18,154	1632.6	1,637	8.99%	9.02%	
Okeechobee County Family Health / Healthy Start Coalition, Inc.	587	48.1	53	8.19%	9.03%	
Orange County Healthy Start Coalition, Inc.	16,556	1476.5	1,476	8.92%	8.92%	
Prenatal and Infant Health Care Coalition of Brevard County, Inc.	5,387	444.3	484	8.25%	8.98%	H
Seminole County Healthy Start Coalition, Inc.	4,786	389.5	365	8.14%	7.63%	
The Healthy Start Prenatal & Infant Coalition of Flagler and Volusia Counties, Inc.	5,782	479.7	464	8.30%	8.02%	
The Healthy Start Coalition of Osceola County, Inc.	3,593	290.5	303	8.09%	8.43%	
TOTAL	226,178	19,799	19,799	8.75%	8.75%	

¹ LBW = Low birth Weight, defined as birth weight below 2500 grams.

² The expected number of infant deaths is calculated based on the maternal race, marital status and education characteristics of the births in each county

³ The significance level used is .05