

Pertussis Surveillance

September 2018

Key Points

 23 cases

 No outbreaks


 Average of 3 contacts per case

 <1 year olds had highest incidence


 61% cases not up-to-date/unknown immunizations




The number of pertussis cases reported in September decreased from previous months and remained below the previous 5-year average. In general, more pertussis cases are reported during the summer months.




The 23 pertussis cases in September were reported among the 11 counties outlined in black. During the previous 3 months (June through August), the average county rate has varied throughout the state.



From January 1, 2018 through September 30, 2018, 251 pertussis cases were reported in 32 counties. Since 2014, the number of pertussis cases reported annually decreased. Pertussis is cyclic in nature with peaks in disease every 3-5 years. Pertussis cases last peaked between 2013 and 2014. Thus far in 2018, it appears case counts will remain consistent with those seen during non-peak years.



In September, 8 (35%) of 23 total cases were associated with transmission within households and no cases were outbreak-associated. For most pertussis cases, exposure to other known cases is never identified, and they are not able to be linked to outbreaks.



No pertussis outbreaks were reported in September.

In 2018, a total of 7 pertussis outbreaks have been reported. Outbreak settings include school (3 outbreaks), daycare (2 outbreaks), work place (1 outbreak), and extended family (1 outbreak).





For each case reported in September, there was an average of 3 contacts for whom antibiotics were recommended to prevent illness. For those diagnosed with pertussis, antibiotics can shorten the amount of time they are contagious to others. Antibiotics can also be used to prevent illness in those who have been exposed to someone with pertussis while they are contagious.


23 cases



69 contacts



In September, the rate was highest among infants <1 year old at 1.8 cases per 100,000 population, which is consistent with previous months. Infants experience the greatest burden of pertussis infections, not only in number of cases but also in severity. Infants <2 months old are too young to receive vaccinations against pertussis, which is why vaccination of grandparents, parents, siblings, and other age groups is so important to help prevent infection in infants.




Vaccination is the best way to prevent pertussis infections. In September, more than half of individuals reported with pertussis had not received the recommended number of pertussis vaccinations for their age or had unknown vaccination status. Vaccination against pertussis is important for infants, children, teenagers, and adults. Pregnant women should get vaccinated during the third trimester of each pregnancy to protect their babies.

Never vaccinated | Under-vaccinated | Too young for vaccination | Up-to-date on vaccinations | Unknown vaccination status





Thus far in 2018, over half of cases in infants and children aged 6 months to 5 years were not up-to-date on their pertussis vaccinations. **In general, those who have received at least 1 pertussis vaccination have less severe outcomes than those who have never been vaccinated.** Over half of adults ≥ 19 years old had unknown vaccination status. See the last page for links to vaccination schedules recommended by the Centers for Disease Control and Prevention.



National activity

The number of pertussis cases gradually increased since the 1980s, peaking in 2012 at levels not seen since the 1950s. Since 2012, the number of pertussis cases started gradually decreasing. Pertussis incidence has remained highest among infants < 1 year old and lowest among adults ≥ 20 years old since the 1990s.

Pertussis surveillance goals

- Identify cases to limit transmission in settings with infants or others who may transmit pertussis to infants
- Identify and prevent outbreaks
- Identify contacts of cases and recommend appropriate prevention measures, including exclusion, antibiotic prophylaxis, and immunization, and to monitor the effectiveness of immunization programs and vaccines

To learn more about pertussis, please visit www.floridahealth.gov/pertussis. For more information on the data sources used in Florida for pertussis surveillance, see the last page.

Vaccine-Preventable Diseases Surveillance System Summary

Case Data

- Current case data are preliminary and will change as new information is gathered. The most recent data available are displayed in this report.
- Pertussis, varicella, and hepatitis A are reportable diseases in Florida. Case information is documented by county health department (CHD) epidemiologists in Merlin, Florida's reportable disease surveillance system.
- Only Florida residents are included in case counts, but contact investigations are conducted for all exposed individuals.
 - Pertussis, varicella, and hepatitis A case counts include both confirmed and probable cases, while measles case counts include only confirmed cases.
- CHD epidemiologists also report outbreaks of pertussis, varicella, and hepatitis A into Merlin.
 - Household-associated cases are defined as ≥ 2 cases exposed within the same household.
 - Pertussis and mumps outbreaks are defined as ≥ 2 cases associated with a specific setting outside of a household.
 - Varicella outbreaks are defined as ≥ 5 cases associated with a specific setting outside of a household.
 - Measles outbreaks are defined as any person acquiring measles while in Florida.
- For more information about reportable diseases, please visit www.Floridahealth.gov/diseasereporting.
- For more information about Florida's guides to surveillance and investigation, including disease-specific surveillance case definitions, please visit www.floridahealth.gov/gsi.

Population Data

- Population data used to calculate incidence rates are from FLHealthCHARTS (Community Health Assessment Resource Tool Set).
- For more information about FLHealthCHARTS, please visit www.flhealthcharts.com.

Vaccination Data

- Vaccination data for identified cases are from Merlin, as documented by CHD epidemiologists.
- Vaccination status is determined using the Advisory Committee on Immunization Practices Recommended Immunization Schedule for Children and Adolescents Aged 18 Years or Younger, 2018.
- For more information about immunization schedules, please visit www.cdc.gov/vaccines/schedules/index.html.
- Individuals are considered up-to-date on vaccinations if they have received the recommended number of doses of vaccine for a particular disease for their age at the time of their illness onset. Individuals are considered under-vaccinated if they have received at least one but not all doses of vaccine recommended for a particular disease for their age at the time of their illness onset.
- For a full text version of a new study on pertussis vaccination, please visit www.cidid.org/publications-1/2018/3/29/the-impact-of-past-vaccination-coverage-and-immunity-on-pertussis-resurgence.