

Vibrio parahaemolyticus

Vibrios are gram-negative bacteria that are found in estuarine and marine environments¹. They colonize filter feeding animals such as shellfish and can also be found free-living in seawater. The optimal growing temperature is between 68-95°F (20-35°C), but it can grow at temperatures up to 105°F (41°C). In addition, the bacterium is halophilic². Given their distribution, *Vibrio parahaemolyticus* (*V. parahaemolyticus*) infections occur in states nationwide including those surrounding the Gulf of Mexico³. The bacteria are highly susceptible to pH, cooking, and freezing and can be killed by common disinfectants such as bleach or alcohol².

People become infected with the bacteria in several ways. One way is by consumption of raw or undercooked seafood or cross-contamination of food with contaminated water. In healthy individuals, the bacteria can cause symptoms of gastroenteritis that generally lasts about three days¹. However, in those who are immunocompromised the bacteria can infect the blood causing septicemia that can cause severe or deadly infections in other parts of the body. Septicemia can be characterized by fever and chills, occasionally accompanied by vomiting, diarrhea, abdominal pain, and pain in the extremities. Those at highest risk for developing septicemia are those who are immunocompromised or have other chronic health problems, such as high serum iron levels or liver disease. Bloodstream infections and death are uncommon and usually occur in those with underlying conditions¹. Another way that people can be infected with the bacteria is through an open wound exposure to seawater. Wound infections can be characterized by inflammation at the wound site, which can progress to cellulitis, bullous lesions, and necrosis. Infections can become systemic which can lead to additional symptoms of chills, fever, hypotension, and altered mental status². The health status of individuals can determine the severity of illness, as death or amputations can be a result of wound infections. People with underlying medical conditions may be at increased risk of infection and serious complications¹.

In Florida, under Florida Administrative Code 61C-4.010(8), all public food service establishments serving raw oysters shall display, on menus, placards, or other effective means, the following notice: “Consumer Information: There is risk associated with consuming raw oysters. If you have chronic illness of the liver, stomach or blood or have immune disorders, you are at greater risk of serious illness from raw oysters and should eat oysters fully cooked. If unsure of your risk, consult a physician.”

Infections from *Vibrio* bacteria that cause cholera and vibriosis have been reportable diseases in Florida since 1981. Since 2007, the states of Florida, Texas and California are consistently responsible for reporting the most cases of vibriosis³. *V. parahaemolyticus* is a type of vibriosis that has been an interest of the Florida Department of Agriculture and Consumer Services due to its potential to be linked to locally harvested shellfish. Single cases of *V. parahaemolyticus* are investigated as if they are outbreaks to determine exposure source. From 2000 to 2017, there has been an average of 31.05 *V. parahaemolyticus* cases reported each year in Florida with a range of 13 – 55 cases. Most cases are reported during the months of May through October. Between 2004 and 2017, there were 2,057 cases of vibriosis reported in Florida and *V. parahaemolyticus* accounted for 477 (23.2%) of those cases. During that time frame, there were 159 deaths reported due to vibriosis and *V. parahaemolyticus* infections accounted for 8 (5%) of those cases. For *V. parahaemolyticus* cases, seafood exposures accounted for 35.2% of cases, wound infections accounted for 37.9% of reported cases and the remaining 26.9% had exposures that could not be classified due to multiple factors (unable to interview, had both exposures to seawater and raw seafood). Cases were white (79.2%), non-Hispanic (83.8%) males (68.8%). Data collected from 2008 to 2017 indicated that 13% of reported cases had heart disease, 13% had diabetes, and 6% had renal disease (systematic collection of risk factors was not collected prior to 2008)⁴.

1. *Vibrio* Species Causing Vibriosis. Centers for Disease Control and Prevention. Page last updated: July 16, 2018. Page accessed on July 19, 2018 <https://www.cdc.gov/vibrio/index.html>
2. Bad Bug Book: Foodborne Pathogenic Microorganisms and Natural Toxins Handbook. Food and

Drug Administration. 2012. Pages: 29-32

3. COVIS Annual Summary. Centers for Disease Control and Prevention. Page last updated: May 12, 2016. Page accessed on July 19, 2018 <https://www.cdc.gov/vibrio/surveillance.html>
4. Florida Department of Health, Bureau of Epidemiology. Florida's Reportable Disease Surveillance System – Merlin.

Updated July 2018