

What Americans believe about opioid prescription painkiller use

Presented by:

Deborah A.P. Hersman, president and CEO, National Safety Council

Dr. Donald Teater, M.D., medical advisor, National Safety Council





Deborah A.P. Hersman

President & CEO,

National Safety Council







Dr. Donald Teater, M.D

Medical Advisor,
Prescription Drug Overdose Initiatives,
National Safety Council





Who is NSC?


Our Mission:

The National Safety Council saves lives by preventing injuries and deaths at work, in homes and communities, and on the roads through leadership, research, education and advocacy.



Drug overdose trends

Number of Poisoning Related Deaths in U.S





National Poll

Methodology:

- Conducted an online survey with a nationally representative sample balanced to U.S. Census figures for gender, age geographic region, and ethnicity.
- To qualify for the study, respondents had to be 18 or older.
- Fieldwork was completed between January 30th thru February 5th, 2015.


Key Takeaways:

- 1. Americans don't know their painkillers contain opioids, or that it is a felony to share them.
- 2. Opioid users are unconcerned about addiction, but most have reason to worry.
- 3. Opioid users overestimate the benefits of opioids and underestimate the risks of addiction or death.





Nearly 30% of respondents say they have taken or been prescribed an opioid painkiller in the past three years.






Actual usage is much higher when users are told which painkillers contain opioids.

Names of Common Opioid Painkillers

- Codeine
- Fentanyl (Actiq, Duragesic, Fentora)
- Hydrocodone (Lorcet, Lortab, Norco, Vicodin)
- Hydrocodone and Acetaminophen (Lorcet, Lortab, Norco, Vicodin)
- Hydromorphone (Dilaudid, Exalgo)

- Meperidine (Demerol)
- Methadone (Dolophine, Methadose)
- Morphine (Avinza, Kadian, MS Contin, Ora-Morph SR)
- Oxycodone (OxyContin, Oxyfast, Percocet, Roxicodone)
- Oxycodone and Acetaminophen (Percocet)
- Oxycodone and Naloxone (Targiniq ER)



Source: NSC Rx Study – In the past three years have you...


© 2015 National Safety Council – Opioid Painkiller Media Briefing



Nearly 70% of opioid painkiller users do not know sharing painkillers is a felony

 Majority of users are likely to believe that sharing their prescriptions with family/friends is perfectly legal or slightly risky

<u>Attitude Towards Sharing Medications with Family/Friends – Among Users</u>





Source: NSC Rx Study - Which of the following best describes the act of sharing a prescription painkiller with a friend or family member? (Among users –n =427)



Nine in 10 opioid painkiller users are unconcerned about addiction

Specific Concerns about Opioid Prescription







Most users should be concerned about addiction. Nearly 60% have at least one addiction risk factor...

...and opioid users admit these pills are more addictive than other medications.

Personal or Family History

43% 22% 26% 16% 16% Alcoholism Depression Use of Physical, Do not have history with psychiatric mental or medication sexual abuse any of these

Agreement Statement*

"Opioid prescriptions are more addictive than other types of prescriptions"





Source: NSC Rx Study


- Which of the following were you most concerned with before taking the prescription? (Among users -n =427)
- Please indicate your agreement with each of the following statements about opioid painkillers as a way of dealing with severe pain.




Users overestimate the benefits of taking opioid painkillers.

Opinions of Using Opioids as Pain Solution

%Somewhat/Strongly Agree











Americans know opioid painkillers are addictive – but they do not understand how addictive.





Only one in five Americans consider prescription pain medication to be a serious safety threat.





NSC resources

- Visit nsc.org/rxpainkillers for up-to-date information
- Report: Prescription Nation
- Report: The Proactive Role Employers Can Take
- Report: Evidence for the Efficacy of Pain Medications
- Report: The Psychological and Physical Side Effects of Pain Medications

Tips for the Public:

- Talk to your doctor about alternatives.
- Take opioid painkillers for the shortest length of time possible, and at the lowest dose.
- Understand the risks, especially the risk of addiction.
- Never share your medication, even with friends or family.
- Dispose of unused drugs properly. Never keep them or save them for later.
- Store opioid painkillers in a locked box or a locking cabinet.



Subject matter experts



Deborah A.P. Hersman president & CEO



Dr. Donald Teater, M.D. medical advisor

Contact information

(630) 775-2226 or email media@nsc.org





Questions?

