

Franklin County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Franklin County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	16.3	(10.4 - 22.3)	6.1	(5.8 - 6.4)	+		
Smoked cigars on one or more of the past 30 days	19.2	(12.0 - 26.5)	6.9	(6.6 - 7.2)	+		
Used smokeless tobacco on one or more of the past 30 days	11.3	(5.6 - 17.1)	3.5	(3.2 - 3.7)	+		
Used any form of tobacco on one or more of the past 30 days	30.5	(20.2 - 40.8)	11.2	(10.8 - 11.7)	+		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	44.9	(35.7 - 54.0)	65.7	(65.1 - 66.3)		+	
Exposed to secondhand smoke during the past 7 days	66.2	(57.5 - 75.0)	44.1	(43.4 - 44.7)	+		
Smoking is allowed in the home	26.2	(18.8 - 33.5)	9.4	(9.0 - 9.8)	+		
Taught about tobacco use in any classes this year	52.9	(39.7 - 66.0)	43.2	(42.3 - 44.1)			+
Parent or guardian has talked about the dangers of tobacco in the past year	60.4	(51.3 - 69.4)	57.4	(56.8 - 58.0)			+
Served as an active member of SWAT	24.1	(13.8 - 34.4)	3.9	(3.7 - 4.1)	+		
<u>Definitely did not think</u> that smokers have more friends	30.9	(23.4 - 38.4)	32.0	(31.5 - 32.6)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	65.8	(58.4 - 73.2)	72.9	(72.4 - 73.4)			+

Franklin County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Franklin County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	27.1	(20.5 - 33.7)	20.5	(20.1 - 21.0)			+
Current asthma, among all individuals	14.4	(9.6 - 19.2)	11.3	(11.0 - 11.7)			+
Asthma attack in past year, among individuals with lifetime asthma	19.8	(6.7 - 32.9)	16.9	(16.0 - 17.8)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	15.5	(9.4 - 21.6)	11.1	(10.7 - 11.5)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	16.5	(9.8 - 23.3)	15.4	(15.0 - 15.9)			+
Sufficient vigorous activity	60.0	(52.7 - 67.3)	66.5	(65.9 - 67.1)			+
Sufficient moderate activity	27.7	(21.1 - 34.4)	25.3	(24.8 - 25.8)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	49.4	(41.9 - 56.9)	43.5	(42.9 - 44.1)			+
Described themselves as slightly or very overweight	37.8	(29.7 - 45.8)	30.0	(29.5 - 30.6)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	19.6	(12.5 - 26.8)	10.9	(10.4 - 11.3)	+		
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	30.6	(23.2 - 37.9)	20.2	(19.7 - 20.7)	+		
Did not go to school on one or more of the past 30 days because they felt unsafe	10.8	(5.7 - 15.8)	6.9	(6.6 - 7.2)			+
Purposely hurt themselves without wanting to die, during the past 12 months	20.8	(14.0 - 27.6)	12.4	(12.0 - 12.8)	+		
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	32.9	(26.0 - 39.7)	22.0	(21.5 - 22.5)	+		