

Hillsborough County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Hillsborough County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	4.5	(3.3 - 5.7)	6.1	(5.8 - 6.4)		+	
Smoked cigars on one or more of the past 30 days	5.8	(4.6 - 7.0)	6.9	(6.6 - 7.2)			+
Used smokeless tobacco on one or more of the past 30 days	2.7	(1.7 - 3.7)	3.5	(3.2 - 3.7)			+
Used any form of tobacco on one or more of the past 30 days	8.9	(7.2 - 10.5)	11.2	(10.8 - 11.7)		+	
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	68.9	(66.0 - 71.7)	65.7	(65.1 - 66.3)			+
Exposed to secondhand smoke during the past 7 days	41.1	(38.2 - 44.1)	44.1	(43.4 - 44.7)			+
Smoking is allowed in the home	9.4	(7.9 - 11.0)	9.4	(9.0 - 9.8)			+
Taught about tobacco use in any classes this year	44.3	(40.2 - 48.5)	43.2	(42.3 - 44.1)			+
Parent or guardian has talked about the dangers of tobacco in the past year	53.8	(51.2 - 56.5)	57.4	(56.8 - 58.0)		+	
Served as an active member of SWAT	4.0	(2.9 - 5.1)	3.9	(3.7 - 4.1)			+
<u>Definitely did not think</u> that smokers have more friends	33.3	(30.8 - 35.9)	32.0	(31.5 - 32.6)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	72.5	(70.0 - 75.0)	72.9	(72.4 - 73.4)			+

Hillsborough County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Hillsborough County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	22.9	(20.7 - 25.1)	20.5	(20.1 - 21.0)			+
Current asthma, among all individuals	12.1	(10.3 - 13.8)	11.3	(11.0 - 11.7)			+
Asthma attack in past year, among individuals with lifetime asthma	19.5	(15.6 - 23.5)	16.9	(16.0 - 17.8)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	12.1	(10.2 - 13.9)	11.1	(10.7 - 11.5)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	15.8	(13.7 - 18.0)	15.4	(15.0 - 15.9)			+
Sufficient vigorous activity	66.2	(63.4 - 69.1)	66.5	(65.9 - 67.1)			+
Sufficient moderate activity	27.0	(24.9 - 29.2)	25.3	(24.8 - 25.8)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	43.7	(41.0 - 46.4)	43.5	(42.9 - 44.1)			+
Described themselves as slightly or very overweight	29.2	(26.8 - 31.6)	30.0	(29.5 - 30.6)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	9.2	(7.2 - 11.2)	10.9	(10.4 - 11.3)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	18.5	(16.2 - 20.7)	20.2	(19.7 - 20.7)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	9.6	(8.0 - 11.2)	6.9	(6.6 - 7.2)	+		
Purposely hurt themselves without wanting to die, during the past 12 months	12.5	(10.7 - 14.3)	12.4	(12.0 - 12.8)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	20.7	(18.5 - 23.0)	22.0	(21.5 - 22.5)			+