Jackson County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Jackson County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	**10.6**	(8.0 - 13.1)	**6.1**	(5.8 - 6.4)	+		
Smoked cigars on one or more of the past 30 days	**7.7**	(5.4 - 9.9)	**6.9**	(6.6 - 7.2)			+
Used smokeless tobacco on one or more of the past 30 days	**8.6**	(6.0 - 11.2)	**3.5**	(3.2 - 3.7)	+		
Used any form of tobacco on one or more of the past 30 days	**18.1**	(14.2 - 22.0)	**11.2**	(10.8 - 11.7)	+		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	**60.8**	(56.3 - 65.2)	**65.7**	(65.1 - 66.3)			+
Exposed to secondhand smoke during the past 7 days	**54.4**	(50.5 - 58.3)	**44.1**	(43.4 - 44.7)	+		
Smoking is allowed in the home	**14.5**	(11.9 - 17.1)	**9.4**	(9.0 - 9.8)	+		
Taught about tobacco use in any classes this year	**55.5**	(48.8 - 62.3)	**43.2**	(42.3 - 44.1)	+		
Parent or guardian has talked about the dangers of tobacco in the past year	**53.6**	(50.4 - 56.8)	**57.4**	(56.8 - 58.0)		+	
Served as an active member of SWAT	**12.3**	(9.7 - 14.9)	**3.9**	(3.7 - 4.1)	+		
Definitely did not think that smokers have more friends	**33.7**	(29.5 - 37.8)	**32.0**	(31.5 - 32.6)			+
Definitely did not think that smoking helps young people “look cool” or “fit in”	**72.6**	(69.1 - 76.1)	**72.9**	(72.4 - 73.4)			+

Jackson County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Jackson County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	**22.3**	(19.5 - 25.1)	**20.5**	(20.1 - 21.0)			+
Current asthma, among all individuals	**11.3**	(9.3 - 13.4)	**11.3**	(11.0 - 11.7)			+
Asthma attack in past year, among individuals with lifetime asthma	**12.8**	(8.4 - 17.3)	**16.9**	(16.0 - 17.8)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	**15.3**	(12.9 - 17.7)	**11.1**	(10.7 - 11.5)	+		
Overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex)	**17.4**	(15.2 - 19.6)	**15.4**	(15.0 - 15.9)			+
Sufficient vigorous activity	**66.0**	(62.7 - 69.2)	**66.5**	(65.9 - 67.1)			+
Sufficient moderate activity	**23.6**	(21.1 - 26.2)	**25.3**	(24.8 - 25.8)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	**46.8**	(43.4 - 50.1)	**43.5**	(42.9 - 44.1)			+
Described themselves as slightly or very overweight	**31.2**	(28.1 - 34.2)	**30.0**	(29.5 - 30.6)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	**15.5**	(12.6 - 18.3)	**10.9**	(10.4 - 11.3)	+		
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	**20.7**	(18.2 - 23.1)	**20.2**	(19.7 - 20.7)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	**5.2**	(3.7 - 6.7)	**6.9**	(6.6 - 7.2)			+
Purposely hurt themselves without wanting to die, during the past 12 months	**11.7**	(9.6 - 13.9)	**12.4**	(12.0 - 12.8)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	**21.8**	(19.0 - 24.5)	**22.0**	(21.5 - 22.5)			+