

Martin County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Martin County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	4.6	(3.1 - 6.0)	6.1	(5.8 - 6.4)			+
Smoked cigars on one or more of the past 30 days	4.3	(2.9 - 5.6)	6.9	(6.6 - 7.2)		+	
Used smokeless tobacco on one or more of the past 30 days	3.0	(1.9 - 4.0)	3.5	(3.2 - 3.7)			+
Used any form of tobacco on one or more of the past 30 days	8.3	(6.3 - 10.3)	11.2	(10.8 - 11.7)		+	
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	64.3	(60.6 - 67.9)	65.7	(65.1 - 66.3)			+
Exposed to secondhand smoke during the past 7 days	45.3	(41.6 - 49.0)	44.1	(43.4 - 44.7)			+
Smoking is allowed in the home	6.8	(5.2 - 8.4)	9.4	(9.0 - 9.8)		+	
Taught about tobacco use in any classes this year	52.9	(47.8 - 58.0)	43.2	(42.3 - 44.1)	+		
Parent or guardian has talked about the dangers of tobacco in the past year	58.8	(55.3 - 62.4)	57.4	(56.8 - 58.0)			+
Served as an active member of SWAT	4.5	(3.1 - 6.0)	3.9	(3.7 - 4.1)			+
<u>Definitely did not think</u> that smokers have more friends	43.3	(39.5 - 47.2)	32.0	(31.5 - 32.6)	+		
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	75.5	(72.7 - 78.4)	72.9	(72.4 - 73.4)			+

Martin County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Martin County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	16.6	(14.1 - 19.1)	20.5	(20.1 - 21.0)		+	
Current asthma, among all individuals	9.8	(8.0 - 11.6)	11.3	(11.0 - 11.7)			+
Asthma attack in past year, among individuals with lifetime asthma	17.6	(11.6 - 23.5)	16.9	(16.0 - 17.8)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	11.2	(8.8 - 13.6)	11.1	(10.7 - 11.5)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	12.9	(10.4 - 15.5)	15.4	(15.0 - 15.9)			+
Sufficient vigorous activity	70.7	(67.4 - 74.0)	66.5	(65.9 - 67.1)	+		
Sufficient moderate activity	29.5	(26.7 - 32.4)	25.3	(24.8 - 25.8)	+		
Exercised to lose weight or to keep from gaining weight during the past 30 days	40.5	(37.2 - 43.9)	43.5	(42.9 - 44.1)			+
Described themselves as slightly or very overweight	28.0	(25.1 - 31.0)	30.0	(29.5 - 30.6)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	7.3	(5.4 - 9.1)	10.9	(10.4 - 11.3)		+	
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	18.2	(15.6 - 20.9)	20.2	(19.7 - 20.7)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	4.7	(3.3 - 6.2)	6.9	(6.6 - 7.2)		+	
Purposely hurt themselves without wanting to die, during the past 12 months	8.8	(7.0 - 10.6)	12.4	(12.0 - 12.8)		+	
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	17.6	(15.1 - 20.1)	22.0	(21.5 - 22.5)		+	