

Okaloosa County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Okaloosa County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	6.4	(4.8 - 8.0)	6.1	(5.8 - 6.4)			+
Smoked cigars on one or more of the past 30 days	5.8	(4.1 - 7.5)	6.9	(6.6 - 7.2)			+
Used smokeless tobacco on one or more of the past 30 days	4.1	(2.7 - 5.4)	3.5	(3.2 - 3.7)			+
Used any form of tobacco on one or more of the past 30 days	11.3	(8.9 - 13.7)	11.2	(10.8 - 11.7)			+
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	68.1	(64.9 - 71.2)	65.7	(65.1 - 66.3)			+
Exposed to secondhand smoke during the past 7 days	46.4	(43.0 - 49.8)	44.1	(43.4 - 44.7)			+
Smoking is allowed in the home	9.3	(7.3 - 11.3)	9.4	(9.0 - 9.8)			+
Taught about tobacco use in any classes this year	45.1	(40.5 - 49.7)	43.2	(42.3 - 44.1)			+
Parent or guardian has talked about the dangers of tobacco in the past year	58.3	(55.1 - 61.4)	57.4	(56.8 - 58.0)			+
Served as an active member of SWAT	2.5	(1.5 - 3.5)	3.9	(3.7 - 4.1)		+	
<u>Definitely did not think</u> that smokers have more friends	35.4	(32.6 - 38.1)	32.0	(31.5 - 32.6)	+		
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	78.7	(76.2 - 81.2)	72.9	(72.4 - 73.4)	+		

Okaloosa County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Okaloosa County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	21.2	(18.9 - 23.4)	20.5	(20.1 - 21.0)			+
Current asthma, among all individuals	13.4	(11.4 - 15.4)	11.3	(11.0 - 11.7)			+
Asthma attack in past year, among individuals with lifetime asthma	19.2	(14.2 - 24.3)	16.9	(16.0 - 17.8)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	11.0	(9.0 - 12.9)	11.1	(10.7 - 11.5)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	11.9	(9.7 - 14.1)	15.4	(15.0 - 15.9)		+	
Sufficient vigorous activity	68.9	(65.6 - 72.1)	66.5	(65.9 - 67.1)			+
Sufficient moderate activity	26.5	(23.8 - 29.2)	25.3	(24.8 - 25.8)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	41.1	(37.7 - 44.6)	43.5	(42.9 - 44.1)			+
Described themselves as slightly or very overweight	29.1	(26.3 - 31.9)	30.0	(29.5 - 30.6)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	7.3	(5.6 - 9.1)	10.9	(10.4 - 11.3)		+	
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	16.9	(14.1 - 19.7)	20.2	(19.7 - 20.7)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	4.9	(3.4 - 6.3)	6.9	(6.6 - 7.2)		+	
Purposely hurt themselves without wanting to die, during the past 12 months	12.7	(10.6 - 14.8)	12.4	(12.0 - 12.8)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	22.6	(19.6 - 25.6)	22.0	(21.5 - 22.5)			+