

Seminole County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Seminole County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	5.3	(4.0 - 6.6)	6.1	(5.8 - 6.4)			+
Smoked cigars on one or more of the past 30 days	7.1	(5.7 - 8.6)	6.9	(6.6 - 7.2)			+
Used smokeless tobacco on one or more of the past 30 days	2.9	(1.9 - 3.9)	3.5	(3.2 - 3.7)			+
Used any form of tobacco on one or more of the past 30 days	10.6	(8.9 - 12.3)	11.2	(10.8 - 11.7)			+
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	68.0	(64.8 - 71.2)	65.7	(65.1 - 66.3)			+
Exposed to secondhand smoke during the past 7 days	39.1	(36.1 - 42.1)	44.1	(43.4 - 44.7)		+	
Smoking is allowed in the home	7.1	(5.6 - 8.7)	9.4	(9.0 - 9.8)		+	
Taught about tobacco use in any classes this year	31.1	(27.7 - 34.6)	43.2	(42.3 - 44.1)		+	
Parent or guardian has talked about the dangers of tobacco in the past year	57.5	(54.8 - 60.2)	57.4	(56.8 - 58.0)			+
Served as an active member of SWAT	2.4	(1.6 - 3.3)	3.9	(3.7 - 4.1)		+	
<u>Definitely did not think</u> that smokers have more friends	34.1	(31.4 - 36.8)	32.0	(31.5 - 32.6)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	75.1	(72.7 - 77.6)	72.9	(72.4 - 73.4)			+

Seminole County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Seminole County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	19.9	(17.6 - 22.1)	20.5	(20.1 - 21.0)			+
Current asthma, among all individuals	12.0	(10.3 - 13.6)	11.3	(11.0 - 11.7)			+
Asthma attack in past year, among individuals with lifetime asthma	19.3	(14.6 - 24.0)	16.9	(16.0 - 17.8)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	8.6	(7.0 - 10.3)	11.1	(10.7 - 11.5)		+	
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	11.5	(9.6 - 13.4)	15.4	(15.0 - 15.9)		+	
Sufficient vigorous activity	70.1	(67.6 - 72.6)	66.5	(65.9 - 67.1)	+		
Sufficient moderate activity	27.2	(24.8 - 29.5)	25.3	(24.8 - 25.8)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	41.6	(38.7 - 44.5)	43.5	(42.9 - 44.1)			+
Described themselves as slightly or very overweight	28.7	(26.1 - 31.3)	30.0	(29.5 - 30.6)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	6.1	(4.5 - 7.7)	10.9	(10.4 - 11.3)		+	
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	17.9	(15.8 - 20.0)	20.2	(19.7 - 20.7)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	6.0	(4.5 - 7.5)	6.9	(6.6 - 7.2)			+
Purposely hurt themselves without wanting to die, during the past 12 months	12.6	(10.4 - 14.7)	12.4	(12.0 - 12.8)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	23.2	(20.4 - 25.9)	22.0	(21.5 - 22.5)			+