

Taylor County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Taylor County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	7.3	(4.4 - 10.3)	6.1	(5.8 - 6.4)			+
Smoked cigars on one or more of the past 30 days	7.5	(5.2 - 9.8)	6.9	(6.6 - 7.2)			+
Used smokeless tobacco on one or more of the past 30 days	6.8	(4.3 - 9.3)	3.5	(3.2 - 3.7)	+		
Used any form of tobacco on one or more of the past 30 days	15.2	(11.6 - 18.9)	11.2	(10.8 - 11.7)			+
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	58.2	(52.7 - 63.7)	65.7	(65.1 - 66.3)		+	
Exposed to secondhand smoke during the past 7 days	59.2	(54.5 - 63.9)	44.1	(43.4 - 44.7)	+		
Smoking is allowed in the home	15.6	(12.3 - 19.0)	9.4	(9.0 - 9.8)	+		
Taught about tobacco use in any classes this year	36.2	(30.3 - 42.1)	43.2	(42.3 - 44.1)		+	
Parent or guardian has talked about the dangers of tobacco in the past year	54.0	(48.4 - 59.6)	57.4	(56.8 - 58.0)			+
Served as an active member of SWAT	10.6	(6.9 - 14.2)	3.9	(3.7 - 4.1)	+		
<u>Definitely did not think</u> that smokers have more friends	35.1	(30.5 - 39.6)	32.0	(31.5 - 32.6)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	74.5	(69.4 - 79.5)	72.9	(72.4 - 73.4)			+

Taylor County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Taylor County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	24.4	(19.8 - 29.0)	20.5	(20.1 - 21.0)			+
Current asthma, among all individuals	13.9	(10.2 - 17.7)	11.3	(11.0 - 11.7)			+
Asthma attack in past year, among individuals with lifetime asthma	12.4	(4.9 - 20.0)	16.9	(16.0 - 17.8)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	18.0	(14.2 - 21.8)	11.1	(10.7 - 11.5)	+		
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	17.7	(13.0 - 22.3)	15.4	(15.0 - 15.9)			+
Sufficient vigorous activity	68.0	(62.7 - 73.4)	66.5	(65.9 - 67.1)			+
Sufficient moderate activity	21.0	(16.9 - 25.1)	25.3	(24.8 - 25.8)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	44.6	(40.1 - 49.0)	43.5	(42.9 - 44.1)			+
Described themselves as slightly or very overweight	32.7	(28.0 - 37.5)	30.0	(29.5 - 30.6)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	14.3	(10.8 - 17.8)	10.9	(10.4 - 11.3)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	24.1	(19.8 - 28.4)	20.2	(19.7 - 20.7)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	8.4	(5.7 - 11.1)	6.9	(6.6 - 7.2)			+
Purposely hurt themselves without wanting to die, during the past 12 months	9.2	(6.3 - 12.1)	12.4	(12.0 - 12.8)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	17.2	(13.8 - 20.7)	22.0	(21.5 - 22.5)		+	