

Wakulla County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Wakulla County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	9.5	(6.8 - 12.1)	6.1	(5.8 - 6.4)	+		
Smoked cigars on one or more of the past 30 days	9.7	(7.0 - 12.5)	6.9	(6.6 - 7.2)			+
Used smokeless tobacco on one or more of the past 30 days	7.3	(5.1 - 9.5)	3.5	(3.2 - 3.7)	+		
Used any form of tobacco on one or more of the past 30 days	17.6	(13.8 - 21.3)	11.2	(10.8 - 11.7)	+		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	57.5	(53.0 - 62.1)	65.7	(65.1 - 66.3)		+	
Exposed to secondhand smoke during the past 7 days	53.4	(48.9 - 57.8)	44.1	(43.4 - 44.7)	+		
Smoking is allowed in the home	16.6	(13.6 - 19.5)	9.4	(9.0 - 9.8)	+		
Taught about tobacco use in any classes this year	63.8	(58.7 - 69.0)	43.2	(42.3 - 44.1)	+		
Parent or guardian has talked about the dangers of tobacco in the past year	51.5	(47.4 - 55.5)	57.4	(56.8 - 58.0)		+	
Served as an active member of SWAT	7.0	(5.0 - 9.0)	3.9	(3.7 - 4.1)	+		
<u>Definitely did not think</u> that smokers have more friends	32.2	(28.4 - 36.0)	32.0	(31.5 - 32.6)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	67.9	(63.5 - 72.3)	72.9	(72.4 - 73.4)		+	

Wakulla County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Wakulla County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	26.4	(23.0 - 29.8)	20.5	(20.1 - 21.0)	+		
Current asthma, among all individuals	15.1	(12.1 - 18.0)	11.3	(11.0 - 11.7)	+		
Asthma attack in past year, among individuals with lifetime asthma	14.4	(9.0 - 19.8)	16.9	(16.0 - 17.8)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	13.3	(10.3 - 16.4)	11.1	(10.7 - 11.5)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	14.7	(11.5 - 17.9)	15.4	(15.0 - 15.9)			+
Sufficient vigorous activity	72.5	(68.8 - 76.2)	66.5	(65.9 - 67.1)	+		
Sufficient moderate activity	30.4	(25.9 - 34.9)	25.3	(24.8 - 25.8)	+		
Exercised to lose weight or to keep from gaining weight during the past 30 days	47.7	(43.6 - 51.7)	43.5	(42.9 - 44.1)			+
Described themselves as slightly or very overweight	33.9	(30.0 - 37.8)	30.0	(29.5 - 30.6)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	12.1	(8.7 - 15.6)	10.9	(10.4 - 11.3)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	24.9	(21.5 - 28.4)	20.2	(19.7 - 20.7)	+		
Did not go to school on one or more of the past 30 days because they felt unsafe	7.1	(5.0 - 9.3)	6.9	(6.6 - 7.2)			+
Purposely hurt themselves without wanting to die, during the past 12 months	13.2	(9.8 - 16.6)	12.4	(12.0 - 12.8)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	22.5	(18.7 - 26.4)	22.0	(21.5 - 22.5)			+