

## Baker County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Baker County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	<b>5.5</b>	(2.9 - 8.1)	<b>3.3</b>	(3.0 - 3.5)			<b>+</b>
Smoked cigars on one or more of the past 30 days	<b>5.0</b>	(2.2 - 7.9)	<b>3.9</b>	(3.6 - 4.2)			<b>+</b>
Used smokeless tobacco on one or more of the past 30 days	<b>11.5</b>	(7.8 - 15.1)	<b>2.3</b>	(2.1 - 2.6)	<b>+</b>		
Used any form of tobacco on one or more of the past 30 days	<b>15.7</b>	(10.7 - 20.7)	<b>6.4</b>	(6.0 - 6.8)	<b>+</b>		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	<b>62.9</b>	(56.1 - 69.7)	<b>72.0</b>	(71.2 - 72.8)		<b>+</b>	
Exposed to secondhand smoke during the past 7 days	<b>56.2</b>	(50.4 - 62.1)	<b>41.3</b>	(40.4 - 42.2)	<b>+</b>		
Smoking is allowed in the home	<b>18.2</b>	(14.2 - 22.2)	<b>9.9</b>	(9.5 - 10.4)	<b>+</b>		
Taught about tobacco use in any classes this year	<b>48.6</b>	(39.7 - 57.4)	<b>53.4</b>	(52.0 - 54.7)			<b>+</b>
Parent or guardian has talked about the dangers of tobacco in the past year	<b>61.1</b>	(56.1 - 66.2)	<b>63.9</b>	(63.1 - 64.6)			<b>+</b>
Served as an active member of SWAT	<b>7.5</b>	(4.7 - 10.4)	<b>4.3</b>	(3.9 - 4.7)			<b>+</b>
<u>Definitely did not think</u> that smokers have more friends	<b>33.1</b>	(26.4 - 39.8)	<b>33.7</b>	(32.9 - 34.5)			<b>+</b>
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	<b>72.6</b>	(67.9 - 77.4)	<b>73.7</b>	(73.0 - 74.4)			<b>+</b>

## Baker County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Baker County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	<b>22.8</b>	(17.1 - 28.5)	<b>20.1</b>	(19.5 - 20.7)			<b>+</b>
Current asthma, among all individuals	<b>14.5</b>	(9.8 - 19.2)	<b>12.0</b>	(11.5 - 12.5)			<b>+</b>
Asthma attack in past year, among individuals with lifetime asthma	<b>10.1</b>	(4.2 - 16.1)	<b>18.1</b>	(16.8 - 19.4)		<b>+</b>	
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	<b>15.2</b>	(9.6 - 20.9)	<b>11.6</b>	(11.0 - 12.1)			<b>+</b>
Overweight (i.e., at or above the 85 <sup>th</sup> percentile but below the 95th percentile for body mass index, by age and sex)	<b>15.4</b>	(10.7 - 20.1)	<b>16.5</b>	(15.8 - 17.2)			<b>+</b>
Sufficient vigorous activity	<b>66.0</b>	(60.3 - 71.6)	<b>70.1</b>	(69.2 - 70.9)			<b>+</b>
Sufficient moderate activity	<b>28.4</b>	(22.7 - 34.0)	<b>24.8</b>	(24.1 - 25.4)			<b>+</b>
Exercised to lose weight or to keep from gaining weight during the past 30 days	<b>49.0</b>	(42.2 - 55.9)	<b>44.4</b>	(43.6 - 45.2)			<b>+</b>
Described themselves as slightly or very overweight	<b>37.6</b>	(30.6 - 44.6)	<b>30.2</b>	(29.4 - 30.9)			<b>+</b>
Never or rarely wore a seat belt when riding in a car driven by someone else	<b>16.6</b>	(11.8 - 21.4)	<b>11.4</b>	(10.8 - 12.0)			<b>+</b>
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	<b>23.4</b>	(18.5 - 28.3)	<b>17.8</b>	(17.2 - 18.5)	<b>+</b>		
Did not go to school on one or more of the past 30 days because they felt unsafe	<b>9.1</b>	(6.4 - 11.7)	<b>7.8</b>	(7.3 - 8.3)			<b>+</b>
Purposely hurt themselves without wanting to die, during the past 12 months	<b>12.0</b>	(9.0 - 15.0)	<b>12.6</b>	(12.0 - 13.2)			<b>+</b>
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	<b>18.2</b>	(14.3 - 22.0)	<b>19.2</b>	(18.5 - 19.8)			<b>+</b>

## Baker County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Baker County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	<b>15.4</b>	(11.3 - 19.4)	<b>10.1</b>	(9.6 - 10.6)	<b>+</b>		
Smoked cigars on one or more of the past 30 days	<b>13.7</b>	(9.7 - 17.7)	<b>11.4</b>	(10.9 - 11.9)			<b>+</b>
Used smokeless tobacco on one or more of the past 30 days	<b>16.8</b>	(12.5 - 21.1)	<b>5.6</b>	(5.2 - 6.0)	<b>+</b>		
Used any form of tobacco on one or more of the past 30 days	<b>30.9</b>	(25.4 - 36.5)	<b>17.9</b>	(17.3 - 18.6)	<b>+</b>		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	<b>53.8</b>	(47.4 - 60.3)	<b>58.4</b>	(57.6 - 59.3)			<b>+</b>
Exposed to secondhand smoke during the past 7 days	<b>58.8</b>	(52.8 - 64.9)	<b>47.6</b>	(46.7 - 48.4)	<b>+</b>		
Smoking is allowed in the home	<b>17.9</b>	(13.6 - 22.2)	<b>10.0</b>	(9.5 - 10.5)	<b>+</b>		
Taught about tobacco use in any classes this year	<b>31.9</b>	(23.9 - 39.9)	<b>33.4</b>	(32.4 - 34.4)			<b>+</b>
Parent or guardian has talked about the dangers of tobacco in the past year	<b>40.6</b>	(34.4 - 46.8)	<b>50.4</b>	(49.6 - 51.2)		<b>+</b>	
Served as an active member of SWAT	<b>6.9</b>	(4.2 - 9.5)	<b>4.0</b>	(3.7 - 4.3)			<b>+</b>
<u>Definitely did not think</u> that smokers have more friends	<b>26.9</b>	(22.2 - 31.6)	<b>30.5</b>	(29.8 - 31.2)			<b>+</b>
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	<b>74.0</b>	(69.2 - 78.8)	<b>71.7</b>	(71.0 - 72.4)			<b>+</b>

## Baker County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Baker County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	<b>22.4</b>	(17.7 - 27.1)	<b>20.8</b>	(20.2 - 21.5)			<b>+</b>
Current asthma, among all individuals	<b>12.3</b>	(8.5 - 16.1)	<b>10.6</b>	(10.1 - 11.1)			<b>+</b>
Asthma attack in past year, among individuals with lifetime asthma	<b>10.3</b>	(4.4 - 16.2)	<b>16.3</b>	(15.1 - 17.4)			<b>+</b>
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	<b>16.5</b>	(12.3 - 20.6)	<b>11.1</b>	(10.5 - 11.6)	<b>+</b>		
Overweight (i.e., at or above the 85 <sup>th</sup> percentile but below the 95th percentile for body mass index, by age and sex)	<b>14.1</b>	(10.5 - 17.7)	<b>14.3</b>	(13.7 - 14.9)			<b>+</b>
Sufficient vigorous activity	<b>64.7</b>	(58.4 - 71.0)	<b>62.7</b>	(62.0 - 63.5)			<b>+</b>
Sufficient moderate activity	<b>27.4</b>	(21.6 - 33.2)	<b>26.1</b>	(25.4 - 26.7)			<b>+</b>
Exercised to lose weight or to keep from gaining weight during the past 30 days	<b>48.6</b>	(43.0 - 54.2)	<b>42.3</b>	(41.5 - 43.1)			<b>+</b>
Described themselves as slightly or very overweight	<b>37.8</b>	(32.2 - 43.5)	<b>29.8</b>	(29.0 - 30.5)	<b>+</b>		
Never or rarely wore a seat belt when riding in a car driven by someone else	<b>15.4</b>	(12.1 - 18.7)	<b>11.2</b>	(10.5 - 11.8)	<b>+</b>		
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	<b>21.6</b>	(17.3 - 25.9)	<b>23.1</b>	(22.4 - 23.8)			<b>+</b>
Did not go to school on one or more of the past 30 days because they felt unsafe	<b>7.5</b>	(5.1 - 9.9)	<b>6.9</b>	(6.4 - 7.3)			<b>+</b>
Purposely hurt themselves without wanting to die, during the past 12 months	<b>10.1</b>	(7.1 - 13.2)	<b>12.7</b>	(12.1 - 13.2)			<b>+</b>
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	<b>21.6</b>	(17.1 - 26.1)	<b>24.5</b>	(23.8 - 25.2)			<b>+</b>

## Baker County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Baker County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	<b>9.4</b>	(6.8 - 11.9)	<b>6.1</b>	(5.8 - 6.4)	<b>+</b>		
Smoked cigars on one or more of the past 30 days	<b>8.3</b>	(5.8 - 10.8)	<b>6.9</b>	(6.6 - 7.2)			<b>+</b>
Used smokeless tobacco on one or more of the past 30 days	<b>12.9</b>	(10.0 - 15.9)	<b>3.5</b>	(3.2 - 3.7)	<b>+</b>		
Used any form of tobacco on one or more of the past 30 days	<b>21.5</b>	(17.3 - 25.6)	<b>11.2</b>	(10.8 - 11.7)	<b>+</b>		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	<b>59.6</b>	(54.7 - 64.5)	<b>65.7</b>	(65.1 - 66.3)		<b>+</b>	
Exposed to secondhand smoke during the past 7 days	<b>56.8</b>	(52.3 - 61.4)	<b>44.1</b>	(43.4 - 44.7)	<b>+</b>		
Smoking is allowed in the home	<b>18.1</b>	(14.8 - 21.3)	<b>9.4</b>	(9.0 - 9.8)	<b>+</b>		
Taught about tobacco use in any classes this year	<b>41.8</b>	(35.4 - 48.1)	<b>43.2</b>	(42.3 - 44.1)			<b>+</b>
Parent or guardian has talked about the dangers of tobacco in the past year	<b>50.8</b>	(45.9 - 55.7)	<b>57.4</b>	(56.8 - 58.0)		<b>+</b>	
Served as an active member of SWAT	<b>7.3</b>	(5.3 - 9.3)	<b>3.9</b>	(3.7 - 4.1)	<b>+</b>		
<u>Definitely did not think</u> that smokers have more friends	<b>29.9</b>	(25.6 - 34.1)	<b>32.0</b>	(31.5 - 32.6)			<b>+</b>
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	<b>74.0</b>	(70.2 - 77.8)	<b>72.9</b>	(72.4 - 73.4)			<b>+</b>

## Baker County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Baker County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	<b>22.4</b>	(18.5 - 26.3)	<b>20.5</b>	(20.1 - 21.0)			<b>+</b>
Current asthma, among all individuals	<b>13.6</b>	(10.4 - 16.7)	<b>11.3</b>	(11.0 - 11.7)			<b>+</b>
Asthma attack in past year, among individuals with lifetime asthma	<b>10.7</b>	(6.3 - 15.1)	<b>16.9</b>	(16.0 - 17.8)		<b>+</b>	
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	<b>15.7</b>	(12.1 - 19.3)	<b>11.1</b>	(10.7 - 11.5)	<b>+</b>		
Overweight (i.e., at or above the 85 <sup>th</sup> percentile but below the 95th percentile for body mass index, by age and sex)	<b>15.2</b>	(12.2 - 18.3)	<b>15.4</b>	(15.0 - 15.9)			<b>+</b>
Sufficient vigorous activity	<b>64.8</b>	(60.3 - 69.3)	<b>66.5</b>	(65.9 - 67.1)			<b>+</b>
Sufficient moderate activity	<b>27.9</b>	(23.7 - 32.1)	<b>25.3</b>	(24.8 - 25.8)			<b>+</b>
Exercised to lose weight or to keep from gaining weight during the past 30 days	<b>49.4</b>	(44.7 - 54.0)	<b>43.5</b>	(42.9 - 44.1)	<b>+</b>		
Described themselves as slightly or very overweight	<b>37.6</b>	(33.0 - 42.3)	<b>30.0</b>	(29.5 - 30.6)	<b>+</b>		
Never or rarely wore a seat belt when riding in a car driven by someone else	<b>15.4</b>	(12.4 - 18.4)	<b>10.9</b>	(10.4 - 11.3)	<b>+</b>		
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	<b>21.4</b>	(18.1 - 24.7)	<b>20.2</b>	(19.7 - 20.7)			<b>+</b>
Did not go to school on one or more of the past 30 days because they felt unsafe	<b>7.9</b>	(6.1 - 9.7)	<b>6.9</b>	(6.6 - 7.2)			<b>+</b>
Purposely hurt themselves without wanting to die, during the past 12 months	<b>10.8</b>	(8.5 - 13.0)	<b>12.4</b>	(12.0 - 12.8)			<b>+</b>
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	<b>19.7</b>	(16.7 - 22.7)	<b>22.0</b>	(21.5 - 22.5)			<b>+</b>