

Clay County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Clay County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	3.2	(1.9 - 4.4)	3.3	(3.0 - 3.5)			+
Smoked cigars on one or more of the past 30 days	3.5	(2.2 - 4.9)	3.9	(3.6 - 4.2)			+
Used smokeless tobacco on one or more of the past 30 days	2.0	(1.1 - 2.9)	2.3	(2.1 - 2.6)			+
Used any form of tobacco on one or more of the past 30 days	6.8	(4.9 - 8.8)	6.4	(6.0 - 6.8)			+
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	71.4	(68.3 - 74.5)	72.0	(71.2 - 72.8)			+
Exposed to secondhand smoke during the past 7 days	48.1	(44.5 - 51.7)	41.3	(40.4 - 42.2)	+		
Smoking is allowed in the home	10.9	(8.8 - 13.1)	9.9	(9.5 - 10.4)			+
Taught about tobacco use in any classes this year	66.1	(61.2 - 71.1)	53.4	(52.0 - 54.7)	+		
Parent or guardian has talked about the dangers of tobacco in the past year	63.9	(61.0 - 66.8)	63.9	(63.1 - 64.6)			+
Served as an active member of SWAT	4.0	(2.8 - 5.2)	4.3	(3.9 - 4.7)			+
<u>Definitely did not think</u> that smokers have more friends	33.4	(30.3 - 36.4)	33.7	(32.9 - 34.5)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	75.3	(72.3 - 78.2)	73.7	(73.0 - 74.4)			+

Clay County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Clay County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	19.2	(17.0 - 21.5)	20.1	(19.5 - 20.7)			+
Current asthma, among all individuals	12.8	(10.6 - 15.0)	12.0	(11.5 - 12.5)			+
Asthma attack in past year, among individuals with lifetime asthma	21.7	(16.6 - 26.8)	18.1	(16.8 - 19.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	10.5	(8.7 - 12.3)	11.6	(11.0 - 12.1)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	13.3	(10.8 - 15.7)	16.5	(15.8 - 17.2)		+	
Sufficient vigorous activity	73.7	(70.8 - 76.6)	70.1	(69.2 - 70.9)			+
Sufficient moderate activity	25.5	(23.1 - 27.8)	24.8	(24.1 - 25.4)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	41.8	(38.3 - 45.2)	44.4	(43.6 - 45.2)			+
Described themselves as slightly or very overweight	29.7	(26.5 - 33.0)	30.2	(29.4 - 30.9)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	6.6	(5.1 - 8.1)	11.4	(10.8 - 12.0)		+	
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	17.0	(14.6 - 19.4)	17.8	(17.2 - 18.5)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	5.7	(4.3 - 7.1)	7.8	(7.3 - 8.3)		+	
Purposely hurt themselves without wanting to die, during the past 12 months	12.2	(10.2 - 14.2)	12.6	(12.0 - 13.2)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	18.4	(15.8 - 20.9)	19.2	(18.5 - 19.8)			+

Clay County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Clay County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	12.7	(9.7 - 15.6)	10.1	(9.6 - 10.6)			+
Smoked cigars on one or more of the past 30 days	13.4	(10.5 - 16.3)	11.4	(10.9 - 11.9)			+
Used smokeless tobacco on one or more of the past 30 days	8.3	(5.5 - 11.1)	5.6	(5.2 - 6.0)			+
Used any form of tobacco on one or more of the past 30 days	23.1	(19.0 - 27.2)	17.9	(17.3 - 18.6)	+		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	56.0	(51.4 - 60.7)	58.4	(57.6 - 59.3)			+
Exposed to secondhand smoke during the past 7 days	55.3	(50.6 - 60.0)	47.6	(46.7 - 48.4)	+		
Smoking is allowed in the home	12.0	(9.3 - 14.7)	10.0	(9.5 - 10.5)			+
Taught about tobacco use in any classes this year	24.4	(20.4 - 28.4)	33.4	(32.4 - 34.4)		+	
Parent or guardian has talked about the dangers of tobacco in the past year	46.5	(42.2 - 50.7)	50.4	(49.6 - 51.2)			+
Served as an active member of SWAT	4.5	(2.7 - 6.3)	4.0	(3.7 - 4.3)			+
<u>Definitely did not think</u> that smokers have more friends	27.7	(23.7 - 31.7)	30.5	(29.8 - 31.2)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	75.7	(72.6 - 78.7)	71.7	(71.0 - 72.4)	+		

Clay County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Clay County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	20.5	(17.6 - 23.5)	20.8	(20.2 - 21.5)			+
Current asthma, among all individuals	10.6	(8.3 - 12.9)	10.6	(10.1 - 11.1)			+
Asthma attack in past year, among individuals with lifetime asthma	22.4	(14.8 - 29.9)	16.3	(15.1 - 17.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	9.7	(7.2 - 12.2)	11.1	(10.5 - 11.6)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	13.1	(10.6 - 15.7)	14.3	(13.7 - 14.9)			+
Sufficient vigorous activity	70.3	(66.4 - 74.1)	62.7	(62.0 - 63.5)	+		
Sufficient moderate activity	31.8	(28.1 - 35.6)	26.1	(25.4 - 26.7)	+		
Exercised to lose weight or to keep from gaining weight during the past 30 days	40.3	(36.9 - 43.7)	42.3	(41.5 - 43.1)			+
Described themselves as slightly or very overweight	27.2	(23.8 - 30.5)	29.8	(29.0 - 30.5)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	10.8	(8.0 - 13.7)	11.2	(10.5 - 11.8)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	20.2	(16.8 - 23.6)	23.1	(22.4 - 23.8)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	6.1	(3.4 - 8.9)	6.9	(6.4 - 7.3)			+
Purposely hurt themselves without wanting to die, during the past 12 months	14.6	(11.8 - 17.5)	12.7	(12.1 - 13.2)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	24.8	(21.5 - 28.2)	24.5	(23.8 - 25.2)			+

Clay County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Clay County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	8.0	(6.2 - 9.8)	6.1	(5.8 - 6.4)			+
Smoked cigars on one or more of the past 30 days	8.0	(6.3 - 9.8)	6.9	(6.6 - 7.2)			+
Used smokeless tobacco on one or more of the past 30 days	5.1	(3.4 - 6.7)	3.5	(3.2 - 3.7)			+
Used any form of tobacco on one or more of the past 30 days	14.7	(12.1 - 17.4)	11.2	(10.8 - 11.7)	+		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	64.0	(61.1 - 67.0)	65.7	(65.1 - 66.3)			+
Exposed to secondhand smoke during the past 7 days	51.5	(48.3 - 54.7)	44.1	(43.4 - 44.7)	+		
Smoking is allowed in the home	11.5	(9.7 - 13.4)	9.4	(9.0 - 9.8)			+
Taught about tobacco use in any classes this year	43.9	(39.3 - 48.4)	43.2	(42.3 - 44.1)			+
Parent or guardian has talked about the dangers of tobacco in the past year	55.0	(52.0 - 58.1)	57.4	(56.8 - 58.0)			+
Served as an active member of SWAT	4.3	(3.1 - 5.4)	3.9	(3.7 - 4.1)			+
<u>Definitely did not think</u> that smokers have more friends	30.8	(28.1 - 33.5)	32.0	(31.5 - 32.6)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	75.9	(73.7 - 78.2)	72.9	(72.4 - 73.4)	+		

Clay County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Clay County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	19.8	(17.9 - 21.8)	20.5	(20.1 - 21.0)			+
Current asthma, among all individuals	11.3	(9.6 - 13.0)	11.3	(11.0 - 11.7)			+
Asthma attack in past year, among individuals with lifetime asthma	20.1	(15.0 - 25.2)	16.9	(16.0 - 17.8)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	9.4	(7.8 - 10.9)	11.1	(10.7 - 11.5)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	13.6	(11.8 - 15.5)	15.4	(15.0 - 15.9)			+
Sufficient vigorous activity	72.2	(69.6 - 74.9)	66.5	(65.9 - 67.1)	+		
Sufficient moderate activity	28.7	(26.2 - 31.1)	25.3	(24.8 - 25.8)	+		
Exercised to lose weight or to keep from gaining weight during the past 30 days	41.3	(38.6 - 43.9)	43.5	(42.9 - 44.1)			+
Described themselves as slightly or very overweight	28.9	(26.4 - 31.3)	30.0	(29.5 - 30.6)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	8.9	(7.2 - 10.7)	10.9	(10.4 - 11.3)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	17.9	(15.9 - 20.0)	20.2	(19.7 - 20.7)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	5.2	(3.9 - 6.5)	6.9	(6.6 - 7.2)		+	
Purposely hurt themselves without wanting to die, during the past 12 months	12.9	(11.3 - 14.6)	12.4	(12.0 - 12.8)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	21.3	(19.1 - 23.5)	22.0	(21.5 - 22.5)			+