

## Collier County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Collier County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	<b>1.8</b>	(0.8 - 2.8)	<b>3.3</b>	(3.0 - 3.5)		<b>+</b>	
Smoked cigars on one or more of the past 30 days	<b>3.6</b>	(1.6 - 5.5)	<b>3.9</b>	(3.6 - 4.2)			<b>+</b>
Used smokeless tobacco on one or more of the past 30 days	<b>0.5</b>	(0.0 - 1.3)	<b>2.3</b>	(2.1 - 2.6)		<b>+</b>	
Used any form of tobacco on one or more of the past 30 days	<b>4.7</b>	(2.6 - 6.7)	<b>6.4</b>	(6.0 - 6.8)			<b>+</b>
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	<b>71.0</b>	(66.6 - 75.4)	<b>72.0</b>	(71.2 - 72.8)			<b>+</b>
Exposed to secondhand smoke during the past 7 days	<b>31.9</b>	(27.4 - 36.4)	<b>41.3</b>	(40.4 - 42.2)		<b>+</b>	
Smoking is allowed in the home	<b>5.7</b>	(3.5 - 7.8)	<b>9.9</b>	(9.5 - 10.4)		<b>+</b>	
Taught about tobacco use in any classes this year	<b>43.5</b>	(37.9 - 49.1)	<b>53.4</b>	(52.0 - 54.7)		<b>+</b>	
Parent or guardian has talked about the dangers of tobacco in the past year	<b>61.9</b>	(57.7 - 66.1)	<b>63.9</b>	(63.1 - 64.6)			<b>+</b>
Served as an active member of SWAT	<b>4.2</b>	(2.3 - 6.2)	<b>4.3</b>	(3.9 - 4.7)			<b>+</b>
<u>Definitely did not think</u> that smokers have more friends	<b>34.5</b>	(29.4 - 39.7)	<b>33.7</b>	(32.9 - 34.5)			<b>+</b>
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	<b>69.6</b>	(65.4 - 73.7)	<b>73.7</b>	(73.0 - 74.4)			<b>+</b>

# Collier County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Collier County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	<b>12.0</b>	(9.1 - 14.9)	<b>20.1</b>	(19.5 - 20.7)		<b>+</b>	
Current asthma, among all individuals	<b>6.2</b>	(4.2 - 8.2)	<b>12.0</b>	(11.5 - 12.5)		<b>+</b>	
Asthma attack in past year, among individuals with lifetime asthma	<b>6.8</b>	(0.1 - 13.4)	<b>18.1</b>	(16.8 - 19.4)		<b>+</b>	
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	<b>11.0</b>	(7.2 - 14.8)	<b>11.6</b>	(11.0 - 12.1)			<b>+</b>
Overweight (i.e., at or above the 85 <sup>th</sup> percentile but below the 95th percentile for body mass index, by age and sex)	<b>13.3</b>	(9.2 - 17.5)	<b>16.5</b>	(15.8 - 17.2)			<b>+</b>
Sufficient vigorous activity	<b>76.1</b>	(71.7 - 80.5)	<b>70.1</b>	(69.2 - 70.9)	<b>+</b>		
Sufficient moderate activity	<b>26.1</b>	(22.1 - 30.2)	<b>24.8</b>	(24.1 - 25.4)			<b>+</b>
Exercised to lose weight or to keep from gaining weight during the past 30 days	<b>43.4</b>	(38.5 - 48.3)	<b>44.4</b>	(43.6 - 45.2)			<b>+</b>
Described themselves as slightly or very overweight	<b>25.5</b>	(20.6 - 30.4)	<b>30.2</b>	(29.4 - 30.9)			<b>+</b>
Never or rarely wore a seat belt when riding in a car driven by someone else	<b>12.8</b>	(9.5 - 16.1)	<b>11.4</b>	(10.8 - 12.0)			<b>+</b>
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	<b>18.1</b>	(14.1 - 22.0)	<b>17.8</b>	(17.2 - 18.5)			<b>+</b>
Did not go to school on one or more of the past 30 days because they felt unsafe	<b>8.5</b>	(5.9 - 11.1)	<b>7.8</b>	(7.3 - 8.3)			<b>+</b>
Purposely hurt themselves without wanting to die, during the past 12 months	<b>15.8</b>	(12.4 - 19.3)	<b>12.6</b>	(12.0 - 13.2)			<b>+</b>
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	<b>21.1</b>	(16.7 - 25.4)	<b>19.2</b>	(18.5 - 19.8)			<b>+</b>

## Collier County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Collier County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	<b>12.2</b>	(9.4 - 15.1)	<b>10.1</b>	(9.6 - 10.6)			<b>+</b>
Smoked cigars on one or more of the past 30 days	<b>12.8</b>	(9.7 - 15.9)	<b>11.4</b>	(10.9 - 11.9)			<b>+</b>
Used smokeless tobacco on one or more of the past 30 days	<b>6.1</b>	(4.2 - 8.1)	<b>5.6</b>	(5.2 - 6.0)			<b>+</b>
Used any form of tobacco on one or more of the past 30 days	<b>18.4</b>	(15.0 - 21.8)	<b>17.9</b>	(17.3 - 18.6)			<b>+</b>
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	<b>50.1</b>	(45.5 - 54.6)	<b>58.4</b>	(57.6 - 59.3)		<b>+</b>	
Exposed to secondhand smoke during the past 7 days	<b>45.4</b>	(40.9 - 49.9)	<b>47.6</b>	(46.7 - 48.4)			<b>+</b>
Smoking is allowed in the home	<b>7.7</b>	(5.3 - 10.1)	<b>10.0</b>	(9.5 - 10.5)			<b>+</b>
Taught about tobacco use in any classes this year	<b>29.4</b>	(25.1 - 33.7)	<b>33.4</b>	(32.4 - 34.4)			<b>+</b>
Parent or guardian has talked about the dangers of tobacco in the past year	<b>50.6</b>	(46.7 - 54.6)	<b>50.4</b>	(49.6 - 51.2)			<b>+</b>
Served as an active member of SWAT	<b>5.1</b>	(3.2 - 7.0)	<b>4.0</b>	(3.7 - 4.3)			<b>+</b>
<u>Definitely did not think</u> that smokers have more friends	<b>30.6</b>	(26.7 - 34.4)	<b>30.5</b>	(29.8 - 31.2)			<b>+</b>
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	<b>68.8</b>	(64.7 - 73.0)	<b>71.7</b>	(71.0 - 72.4)			<b>+</b>

## Collier County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Collier County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	<b>13.3</b>	(10.3 - 16.3)	<b>20.8</b>	(20.2 - 21.5)		<b>+</b>	
Current asthma, among all individuals	<b>6.2</b>	(4.2 - 8.1)	<b>10.6</b>	(10.1 - 11.1)		<b>+</b>	
Asthma attack in past year, among individuals with lifetime asthma	<b>18.2</b>	(11.0 - 25.4)	<b>16.3</b>	(15.1 - 17.4)			<b>+</b>
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	<b>8.7</b>	(6.2 - 11.1)	<b>11.1</b>	(10.5 - 11.6)			<b>+</b>
Overweight (i.e., at or above the 85 <sup>th</sup> percentile but below the 95th percentile for body mass index, by age and sex)	<b>14.2</b>	(11.1 - 17.3)	<b>14.3</b>	(13.7 - 14.9)			<b>+</b>
Sufficient vigorous activity	<b>67.9</b>	(63.3 - 72.6)	<b>62.7</b>	(62.0 - 63.5)			<b>+</b>
Sufficient moderate activity	<b>26.8</b>	(23.4 - 30.2)	<b>26.1</b>	(25.4 - 26.7)			<b>+</b>
Exercised to lose weight or to keep from gaining weight during the past 30 days	<b>41.6</b>	(37.7 - 45.5)	<b>42.3</b>	(41.5 - 43.1)			<b>+</b>
Described themselves as slightly or very overweight	<b>27.1</b>	(23.4 - 30.8)	<b>29.8</b>	(29.0 - 30.5)			<b>+</b>
Never or rarely wore a seat belt when riding in a car driven by someone else	<b>11.4</b>	(8.2 - 14.7)	<b>11.2</b>	(10.5 - 11.8)			<b>+</b>
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	<b>25.6</b>	(22.1 - 29.2)	<b>23.1</b>	(22.4 - 23.8)			<b>+</b>
Did not go to school on one or more of the past 30 days because they felt unsafe	<b>6.2</b>	(4.2 - 8.1)	<b>6.9</b>	(6.4 - 7.3)			<b>+</b>
Purposely hurt themselves without wanting to die, during the past 12 months	<b>11.3</b>	(9.1 - 13.5)	<b>12.7</b>	(12.1 - 13.2)			<b>+</b>
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	<b>22.6</b>	(19.3 - 26.0)	<b>24.5</b>	(23.8 - 25.2)			<b>+</b>

## Collier County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Collier County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	<b>6.6</b>	(4.8 - 8.3)	<b>6.1</b>	(5.8 - 6.4)			<b>+</b>
Smoked cigars on one or more of the past 30 days	<b>8.0</b>	(5.9 - 10.2)	<b>6.9</b>	(6.6 - 7.2)			<b>+</b>
Used smokeless tobacco on one or more of the past 30 days	<b>2.6</b>	(1.7 - 3.6)	<b>3.5</b>	(3.2 - 3.7)			<b>+</b>
Used any form of tobacco on one or more of the past 30 days	<b>11.1</b>	(8.7 - 13.4)	<b>11.2</b>	(10.8 - 11.7)			<b>+</b>
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	<b>60.8</b>	(57.1 - 64.5)	<b>65.7</b>	(65.1 - 66.3)		<b>+</b>	
Exposed to secondhand smoke during the past 7 days	<b>38.9</b>	(35.3 - 42.5)	<b>44.1</b>	(43.4 - 44.7)		<b>+</b>	
Smoking is allowed in the home	<b>5.9</b>	(4.5 - 7.4)	<b>9.4</b>	(9.0 - 9.8)		<b>+</b>	
Taught about tobacco use in any classes this year	<b>36.8</b>	(33.0 - 40.5)	<b>43.2</b>	(42.3 - 44.1)		<b>+</b>	
Parent or guardian has talked about the dangers of tobacco in the past year	<b>56.5</b>	(53.5 - 59.6)	<b>57.4</b>	(56.8 - 58.0)			<b>+</b>
Served as an active member of SWAT	<b>4.4</b>	(3.0 - 5.8)	<b>3.9</b>	(3.7 - 4.1)			<b>+</b>
<u>Definitely did not think</u> that smokers have more friends	<b>32.9</b>	(29.7 - 36.2)	<b>32.0</b>	(31.5 - 32.6)			<b>+</b>
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	<b>70.1</b>	(67.1 - 73.0)	<b>72.9</b>	(72.4 - 73.4)			<b>+</b>

## Collier County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Collier County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	<b>13.1</b>	(10.9 - 15.3)	<b>20.5</b>	(20.1 - 21.0)		<b>+</b>	
Current asthma, among all individuals	<b>6.3</b>	(4.9 - 7.8)	<b>11.3</b>	(11.0 - 11.7)		<b>+</b>	
Asthma attack in past year, among individuals with lifetime asthma	<b>13.4</b>	(8.3 - 18.5)	<b>16.9</b>	(16.0 - 17.8)			<b>+</b>
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	<b>9.6</b>	(7.4 - 11.7)	<b>11.1</b>	(10.7 - 11.5)			<b>+</b>
Overweight (i.e., at or above the 85 <sup>th</sup> percentile but below the 95th percentile for body mass index, by age and sex)	<b>13.6</b>	(11.1 - 16.0)	<b>15.4</b>	(15.0 - 15.9)			<b>+</b>
Sufficient vigorous activity	<b>73.0</b>	(69.9 - 76.0)	<b>66.5</b>	(65.9 - 67.1)	<b>+</b>		
Sufficient moderate activity	<b>26.4</b>	(23.8 - 29.0)	<b>25.3</b>	(24.8 - 25.8)			<b>+</b>
Exercised to lose weight or to keep from gaining weight during the past 30 days	<b>42.2</b>	(39.0 - 45.4)	<b>43.5</b>	(42.9 - 44.1)			<b>+</b>
Described themselves as slightly or very overweight	<b>26.2</b>	(23.1 - 29.3)	<b>30.0</b>	(29.5 - 30.6)		<b>+</b>	
Never or rarely wore a seat belt when riding in a car driven by someone else	<b>11.8</b>	(9.5 - 14.1)	<b>10.9</b>	(10.4 - 11.3)			<b>+</b>
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	<b>22.2</b>	(19.4 - 25.0)	<b>20.2</b>	(19.7 - 20.7)			<b>+</b>
Did not go to school on one or more of the past 30 days because they felt unsafe	<b>6.3</b>	(4.7 - 7.9)	<b>6.9</b>	(6.6 - 7.2)			<b>+</b>
Purposely hurt themselves without wanting to die, during the past 12 months	<b>12.7</b>	(10.6 - 14.7)	<b>12.4</b>	(12.0 - 12.8)			<b>+</b>
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	<b>21.4</b>	(18.7 - 24.1)	<b>22.0</b>	(21.5 - 22.5)			<b>+</b>