

Dixie County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Dixie County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	4.2	(1.2 - 7.1)	3.3	(3.0 - 3.5)			+
Smoked cigars on one or more of the past 30 days	1.7	(0.0 - 3.6)	3.9	(3.6 - 4.2)		+	
Used smokeless tobacco on one or more of the past 30 days	7.3	(3.2 - 11.5)	2.3	(2.1 - 2.6)	+		
Used any form of tobacco on one or more of the past 30 days	10.8	(5.2 - 16.5)	6.4	(6.0 - 6.8)			+
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	64.3	(57.6 - 71.1)	72.0	(71.2 - 72.8)		+	
Exposed to secondhand smoke during the past 7 days	56.1	(48.9 - 63.3)	41.3	(40.4 - 42.2)	+		
Smoking is allowed in the home	20.5	(13.8 - 27.2)	9.9	(9.5 - 10.4)	+		
Taught about tobacco use in any classes this year	80.7	(70.7 - 90.7)	53.4	(52.0 - 54.7)	+		
Parent or guardian has talked about the dangers of tobacco in the past year	62.7	(54.8 - 70.7)	63.9	(63.1 - 64.6)			+
Served as an active member of SWAT	15.6	(9.6 - 21.6)	4.3	(3.9 - 4.7)	+		
<u>Definitely did not think</u> that smokers have more friends	30.8	(23.7 - 37.9)	33.7	(32.9 - 34.5)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	78.9	(71.6 - 86.2)	73.7	(73.0 - 74.4)			+

Dixie County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Dixie County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	27.9	(19.4 - 36.3)	20.1	(19.5 - 20.7)			+
Current asthma, among all individuals	12.8	(7.5 - 18.1)	12.0	(11.5 - 12.5)			+
Asthma attack in past year, among individuals with lifetime asthma	14.5	(2.4 - 26.5)	18.1	(16.8 - 19.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	10.7	(7.2 - 14.2)	11.6	(11.0 - 12.1)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	21.8	(14.3 - 29.4)	16.5	(15.8 - 17.2)			+
Sufficient vigorous activity	72.2	(66.6 - 77.9)	70.1	(69.2 - 70.9)			+
Sufficient moderate activity	20.5	(13.5 - 27.4)	24.8	(24.1 - 25.4)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	46.1	(39.1 - 53.1)	44.4	(43.6 - 45.2)			+
Described themselves as slightly or very overweight	36.5	(29.8 - 43.3)	30.2	(29.4 - 30.9)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	20.9	(14.4 - 27.5)	11.4	(10.8 - 12.0)	+		
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	20.7	(14.6 - 26.8)	17.8	(17.2 - 18.5)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	5.0	(1.1 - 8.8)	7.8	(7.3 - 8.3)			+
Purposely hurt themselves without wanting to die, during the past 12 months	10.1	(5.4 - 14.9)	12.6	(12.0 - 13.2)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	15.6	(8.9 - 22.3)	19.2	(18.5 - 19.8)			+

Dixie County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Dixie County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	15.8	(10.9 - 20.7)	10.1	(9.6 - 10.6)	+		
Smoked cigars on one or more of the past 30 days	18.3	(11.7 - 24.9)	11.4	(10.9 - 11.9)			+
Used smokeless tobacco on one or more of the past 30 days	20.7	(11.2 - 30.1)	5.6	(5.2 - 6.0)	+		
Used any form of tobacco on one or more of the past 30 days	35.3	(25.5 - 45.1)	17.9	(17.3 - 18.6)	+		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	44.3	(35.8 - 52.9)	58.4	(57.6 - 59.3)		+	
Exposed to secondhand smoke during the past 7 days	55.4	(48.8 - 62.0)	47.6	(46.7 - 48.4)	+		
Smoking is allowed in the home	20.8	(15.9 - 25.7)	10.0	(9.5 - 10.5)	+		
Taught about tobacco use in any classes this year	65.7	(58.4 - 73.0)	33.4	(32.4 - 34.4)	+		
Parent or guardian has talked about the dangers of tobacco in the past year	46.1	(38.6 - 53.7)	50.4	(49.6 - 51.2)			+
Served as an active member of SWAT	26.2	(18.4 - 34.0)	4.0	(3.7 - 4.3)	+		
<u>Definitely did not think</u> that smokers have more friends	32.9	(25.4 - 40.4)	30.5	(29.8 - 31.2)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	68.6	(58.7 - 78.4)	71.7	(71.0 - 72.4)			+

Dixie County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Dixie County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	27.3	(19.7 - 34.9)	20.8	(20.2 - 21.5)			+
Current asthma, among all individuals	13.7	(6.9 - 20.5)	10.6	(10.1 - 11.1)			+
Asthma attack in past year, among individuals with lifetime asthma	15.5	(4.3 - 26.8)	16.3	(15.1 - 17.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	18.8	(11.3 - 26.2)	11.1	(10.5 - 11.6)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	12.6	(6.3 - 18.9)	14.3	(13.7 - 14.9)			+
Sufficient vigorous activity	65.3	(58.2 - 72.4)	62.7	(62.0 - 63.5)			+
Sufficient moderate activity	28.4	(20.1 - 36.8)	26.1	(25.4 - 26.7)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	44.0	(34.1 - 53.9)	42.3	(41.5 - 43.1)			+
Described themselves as slightly or very overweight	36.0	(28.4 - 43.5)	29.8	(29.0 - 30.5)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	25.4	(16.2 - 34.7)	11.2	(10.5 - 11.8)	+		
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	30.0	(22.4 - 37.5)	23.1	(22.4 - 23.8)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	11.6	(6.6 - 16.7)	6.9	(6.4 - 7.3)			+
Purposely hurt themselves without wanting to die, during the past 12 months	13.3	(7.6 - 19.0)	12.7	(12.1 - 13.2)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	19.6	(13.9 - 25.3)	24.5	(23.8 - 25.2)			+

Dixie County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Dixie County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	9.6	(5.9 - 13.4)	6.1	(5.8 - 6.4)			+
Smoked cigars on one or more of the past 30 days	8.4	(4.7 - 12.1)	6.9	(6.6 - 7.2)			+
Used smokeless tobacco on one or more of the past 30 days	12.3	(7.0 - 17.5)	3.5	(3.2 - 3.7)	+		
Used any form of tobacco on one or more of the past 30 days	20.5	(14.0 - 26.9)	11.2	(10.8 - 11.7)	+		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	55.2	(48.6 - 61.9)	65.7	(65.1 - 66.3)		+	
Exposed to secondhand smoke during the past 7 days	56.1	(51.2 - 61.0)	44.1	(43.4 - 44.7)	+		
Smoking is allowed in the home	21.0	(16.6 - 25.4)	9.4	(9.0 - 9.8)	+		
Taught about tobacco use in any classes this year	72.1	(65.3 - 78.9)	43.2	(42.3 - 44.1)	+		
Parent or guardian has talked about the dangers of tobacco in the past year	54.9	(48.5 - 61.4)	57.4	(56.8 - 58.0)			+
Served as an active member of SWAT	20.3	(15.7 - 24.9)	3.9	(3.7 - 4.1)	+		
<u>Definitely did not think</u> that smokers have more friends	32.3	(27.0 - 37.6)	32.0	(31.5 - 32.6)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	74.7	(68.2 - 81.2)	72.9	(72.4 - 73.4)			+

Dixie County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Dixie County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	28.4	(23.0 - 33.9)	20.5	(20.1 - 21.0)	+		
Current asthma, among all individuals	13.3	(9.4 - 17.2)	11.3	(11.0 - 11.7)			+
Asthma attack in past year, among individuals with lifetime asthma	15.9	(7.9 - 23.8)	16.9	(16.0 - 17.8)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	14.2	(10.1 - 18.3)	11.1	(10.7 - 11.5)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	16.6	(11.6 - 21.5)	15.4	(15.0 - 15.9)			+
Sufficient vigorous activity	68.4	(63.5 - 73.3)	66.5	(65.9 - 67.1)			+
Sufficient moderate activity	24.1	(18.8 - 29.4)	25.3	(24.8 - 25.8)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	44.3	(38.8 - 49.7)	43.5	(42.9 - 44.1)			+
Described themselves as slightly or very overweight	35.3	(30.5 - 40.2)	30.0	(29.5 - 30.6)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	22.5	(16.7 - 28.3)	10.9	(10.4 - 11.3)	+		
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	23.8	(18.8 - 28.7)	20.2	(19.7 - 20.7)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	8.1	(4.7 - 11.5)	6.9	(6.6 - 7.2)			+
Purposely hurt themselves without wanting to die, during the past 12 months	11.4	(7.8 - 14.9)	12.4	(12.0 - 12.8)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	17.3	(12.7 - 21.8)	22.0	(21.5 - 22.5)			+