

Franklin County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Franklin County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	11.8	(3.8 - 19.7)	3.3	(3.0 - 3.5)	+		
Smoked cigars on one or more of the past 30 days	10.7	(2.7 - 18.7)	3.9	(3.6 - 4.2)			+
Used smokeless tobacco on one or more of the past 30 days	8.0	(2.1 - 13.9)	2.3	(2.1 - 2.6)			+
Used any form of tobacco on one or more of the past 30 days	20.3	(7.9 - 32.7)	6.4	(6.0 - 6.8)	+		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	48.4	(36.9 - 60.0)	72.0	(71.2 - 72.8)		+	
Exposed to secondhand smoke during the past 7 days	65.7	(54.0 - 77.4)	41.3	(40.4 - 42.2)	+		
Smoking is allowed in the home	32.9	(22.1 - 43.7)	9.9	(9.5 - 10.4)	+		
Taught about tobacco use in any classes this year	66.2	(48.1 - 84.3)	53.4	(52.0 - 54.7)			+
Parent or guardian has talked about the dangers of tobacco in the past year	62.2	(54.0 - 70.4)	63.9	(63.1 - 64.6)			+
Served as an active member of SWAT	33.7	(17.1 - 50.3)	4.3	(3.9 - 4.7)	+		
<u>Definitely did not think</u> that smokers have more friends	27.1	(19.1 - 35.2)	33.7	(32.9 - 34.5)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	69.2	(59.7 - 78.8)	73.7	(73.0 - 74.4)			+

Franklin County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Franklin County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	19.0	(10.7 - 27.3)	20.1	(19.5 - 20.7)			+
Current asthma, among all individuals	8.4	(4.2 - 12.5)	12.0	(11.5 - 12.5)			+
Asthma attack in past year, among individuals with lifetime asthma	14.3	(0.0 - 29.2)	18.1	(16.8 - 19.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	11.9	(3.8 - 20.1)	11.6	(11.0 - 12.1)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	17.4	(7.9 - 26.8)	16.5	(15.8 - 17.2)			+
Sufficient vigorous activity	58.9	(50.3 - 67.5)	70.1	(69.2 - 70.9)		+	
Sufficient moderate activity	26.2	(17.0 - 35.3)	24.8	(24.1 - 25.4)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	50.8	(42.1 - 59.5)	44.4	(43.6 - 45.2)			+
Described themselves as slightly or very overweight	32.5	(22.1 - 42.9)	30.2	(29.4 - 30.9)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	14.2	(7.8 - 20.6)	11.4	(10.8 - 12.0)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	29.6	(18.4 - 40.8)	17.8	(17.2 - 18.5)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	10.4	(3.5 - 17.3)	7.8	(7.3 - 8.3)			+
Purposely hurt themselves without wanting to die, during the past 12 months	22.4	(12.2 - 32.5)	12.6	(12.0 - 13.2)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	30.5	(21.6 - 39.3)	19.2	(18.5 - 19.8)	+		

Franklin County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Franklin County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	27.1	(18.3 - 35.9)	10.1	(9.6 - 10.6)	+		
Smoked cigars on one or more of the past 30 days	31.7	(21.8 - 41.6)	11.4	(10.9 - 11.9)	+		
Used smokeless tobacco on one or more of the past 30 days	16.8	(7.4 - 26.1)	5.6	(5.2 - 6.0)	+		
Used any form of tobacco on one or more of the past 30 days	48.0	(34.3 - 61.7)	17.9	(17.3 - 18.6)	+		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	34.8	(21.2 - 48.4)	58.4	(57.6 - 59.3)		+	
Exposed to secondhand smoke during the past 7 days	70.1	(58.2 - 81.9)	47.6	(46.7 - 48.4)	+		
Smoking is allowed in the home	20.9	(14.1 - 27.7)	10.0	(9.5 - 10.5)	+		
Taught about tobacco use in any classes this year	31.9	(19.5 - 44.2)	33.4	(32.4 - 34.4)			+
Parent or guardian has talked about the dangers of tobacco in the past year	55.9	(38.7 - 73.2)	50.4	(49.6 - 51.2)			+
Served as an active member of SWAT	12.2	(4.4 - 19.9)	4.0	(3.7 - 4.3)	+		
<u>Definitely did not think</u> that smokers have more friends	35.5	(23.7 - 47.2)	30.5	(29.8 - 31.2)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	62.5	(51.2 - 73.7)	71.7	(71.0 - 72.4)			+

Franklin County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Franklin County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	32.3	(23.9 - 40.7)	20.8	(20.2 - 21.5)	+		
Current asthma, among all individuals	19.4	(12.5 - 26.3)	10.6	(10.1 - 11.1)	+		
Asthma attack in past year, among individuals with lifetime asthma	26.0	(6.6 - 45.5)	16.3	(15.1 - 17.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	18.8	(10.2 - 27.4)	11.1	(10.5 - 11.6)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	16.3	(5.9 - 26.6)	14.3	(13.7 - 14.9)			+
Sufficient vigorous activity	60.1	(48.4 - 71.9)	62.7	(62.0 - 63.5)			+
Sufficient moderate activity	31.8	(20.7 - 42.8)	26.1	(25.4 - 26.7)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	47.5	(34.3 - 60.7)	42.3	(41.5 - 43.1)			+
Described themselves as slightly or very overweight	40.6	(29.3 - 51.9)	29.8	(29.0 - 30.5)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	29.1	(15.8 - 42.4)	11.2	(10.5 - 11.8)	+		
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	37.2	(27.0 - 47.3)	23.1	(22.4 - 23.8)	+		
Did not go to school on one or more of the past 30 days because they felt unsafe	10.8	(4.2 - 17.4)	6.9	(6.4 - 7.3)			+
Purposely hurt themselves without wanting to die, during the past 12 months	18.1	(8.9 - 27.3)	12.7	(12.1 - 13.2)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	35.3	(25.0 - 45.6)	24.5	(23.8 - 25.2)			+

Franklin County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Franklin County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	16.3	(10.4 - 22.3)	6.1	(5.8 - 6.4)	+		
Smoked cigars on one or more of the past 30 days	19.2	(12.0 - 26.5)	6.9	(6.6 - 7.2)	+		
Used smokeless tobacco on one or more of the past 30 days	11.3	(5.6 - 17.1)	3.5	(3.2 - 3.7)	+		
Used any form of tobacco on one or more of the past 30 days	30.5	(20.2 - 40.8)	11.2	(10.8 - 11.7)	+		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	44.9	(35.7 - 54.0)	65.7	(65.1 - 66.3)		+	
Exposed to secondhand smoke during the past 7 days	66.2	(57.5 - 75.0)	44.1	(43.4 - 44.7)	+		
Smoking is allowed in the home	26.2	(18.8 - 33.5)	9.4	(9.0 - 9.8)	+		
Taught about tobacco use in any classes this year	52.9	(39.7 - 66.0)	43.2	(42.3 - 44.1)			+
Parent or guardian has talked about the dangers of tobacco in the past year	60.4	(51.3 - 69.4)	57.4	(56.8 - 58.0)			+
Served as an active member of SWAT	24.1	(13.8 - 34.4)	3.9	(3.7 - 4.1)	+		
<u>Definitely did not think</u> that smokers have more friends	30.9	(23.4 - 38.4)	32.0	(31.5 - 32.6)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	65.8	(58.4 - 73.2)	72.9	(72.4 - 73.4)			+

Franklin County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Franklin County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	27.1	(20.5 - 33.7)	20.5	(20.1 - 21.0)			+
Current asthma, among all individuals	14.4	(9.6 - 19.2)	11.3	(11.0 - 11.7)			+
Asthma attack in past year, among individuals with lifetime asthma	19.8	(6.7 - 32.9)	16.9	(16.0 - 17.8)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	15.5	(9.4 - 21.6)	11.1	(10.7 - 11.5)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	16.5	(9.8 - 23.3)	15.4	(15.0 - 15.9)			+
Sufficient vigorous activity	60.0	(52.7 - 67.3)	66.5	(65.9 - 67.1)			+
Sufficient moderate activity	27.7	(21.1 - 34.4)	25.3	(24.8 - 25.8)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	49.4	(41.9 - 56.9)	43.5	(42.9 - 44.1)			+
Described themselves as slightly or very overweight	37.8	(29.7 - 45.8)	30.0	(29.5 - 30.6)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	19.6	(12.5 - 26.8)	10.9	(10.4 - 11.3)	+		
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	30.6	(23.2 - 37.9)	20.2	(19.7 - 20.7)	+		
Did not go to school on one or more of the past 30 days because they felt unsafe	10.8	(5.7 - 15.8)	6.9	(6.6 - 7.2)			+
Purposely hurt themselves without wanting to die, during the past 12 months	20.8	(14.0 - 27.6)	12.4	(12.0 - 12.8)	+		
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	32.9	(26.0 - 39.7)	22.0	(21.5 - 22.5)	+		