

Hamilton County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Hamilton County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	4.3	(0.2 - 8.5)	3.3	(3.0 - 3.5)			+
Smoked cigars on one or more of the past 30 days	3.4	(0.3 - 6.5)	3.9	(3.6 - 4.2)			+
Used smokeless tobacco on one or more of the past 30 days	9.5	(3.5 - 15.5)	2.3	(2.1 - 2.6)	+		
Used any form of tobacco on one or more of the past 30 days	13.8	(6.7 - 21.0)	6.4	(6.0 - 6.8)			+
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	71.8	(61.3 - 82.2)	72.0	(71.2 - 72.8)			+
Exposed to secondhand smoke during the past 7 days	51.7	(39.9 - 63.4)	41.3	(40.4 - 42.2)			+
Smoking is allowed in the home	13.2	(5.9 - 20.5)	9.9	(9.5 - 10.4)			+
Taught about tobacco use in any classes this year	42.4	(28.2 - 56.5)	53.4	(52.0 - 54.7)			+
Parent or guardian has talked about the dangers of tobacco in the past year	64.7	(55.9 - 73.4)	63.9	(63.1 - 64.6)			+
Served as an active member of SWAT	19.8	(12.8 - 26.8)	4.3	(3.9 - 4.7)	+		
<u>Definitely did not think</u> that smokers have more friends	34.5	(26.5 - 42.6)	33.7	(32.9 - 34.5)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	67.2	(59.9 - 74.6)	73.7	(73.0 - 74.4)			+

Hamilton County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Hamilton County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	19.8	(12.2 - 27.3)	20.1	(19.5 - 20.7)			+
Current asthma, among all individuals	14.0	(6.8 - 21.2)	12.0	(11.5 - 12.5)			+
Asthma attack in past year, among individuals with lifetime asthma	12.6	(0.0 - 31.7)	18.1	(16.8 - 19.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	16.9	(10.2 - 23.7)	11.6	(11.0 - 12.1)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	22.7	(15.2 - 30.2)	16.5	(15.8 - 17.2)			+
Sufficient vigorous activity	60.0	(48.3 - 71.7)	70.1	(69.2 - 70.9)			+
Sufficient moderate activity	18.3	(12.5 - 24.1)	24.8	(24.1 - 25.4)		+	
Exercised to lose weight or to keep from gaining weight during the past 30 days	45.5	(35.8 - 55.2)	44.4	(43.6 - 45.2)			+
Described themselves as slightly or very overweight	28.2	(18.3 - 38.1)	30.2	(29.4 - 30.9)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	21.0	(14.2 - 27.7)	11.4	(10.8 - 12.0)	+		
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	19.9	(12.9 - 27.0)	17.8	(17.2 - 18.5)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	6.8	(2.0 - 11.6)	7.8	(7.3 - 8.3)			+
Purposely hurt themselves without wanting to die, during the past 12 months	14.0	(7.8 - 20.1)	12.6	(12.0 - 13.2)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	22.0	(15.4 - 28.6)	19.2	(18.5 - 19.8)			+

Hamilton County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Hamilton County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	14.5	(8.9 - 20.1)	10.1	(9.6 - 10.6)			+
Smoked cigars on one or more of the past 30 days	13.1	(5.4 - 20.7)	11.4	(10.9 - 11.9)			+
Used smokeless tobacco on one or more of the past 30 days	9.1	(3.8 - 14.4)	5.6	(5.2 - 6.0)			+
Used any form of tobacco on one or more of the past 30 days	21.9	(14.0 - 29.9)	17.9	(17.3 - 18.6)			+
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	51.8	(44.0 - 59.6)	58.4	(57.6 - 59.3)			+
Exposed to secondhand smoke during the past 7 days	60.5	(49.1 - 71.9)	47.6	(46.7 - 48.4)	+		
Smoking is allowed in the home	17.3	(7.3 - 27.3)	10.0	(9.5 - 10.5)			+
Taught about tobacco use in any classes this year	50.8	(36.1 - 65.6)	33.4	(32.4 - 34.4)	+		
Parent or guardian has talked about the dangers of tobacco in the past year	62.4	(52.5 - 72.2)	50.4	(49.6 - 51.2)	+		
Served as an active member of SWAT	16.7	(7.1 - 26.2)	4.0	(3.7 - 4.3)	+		
<u>Definitely did not think</u> that smokers have more friends	23.4	(14.7 - 32.1)	30.5	(29.8 - 31.2)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	71.5	(62.8 - 80.1)	71.7	(71.0 - 72.4)			+

Hamilton County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Hamilton County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	12.6	(6.8 - 18.5)	20.8	(20.2 - 21.5)		+	
Current asthma, among all individuals	5.0	(1.1 - 8.9)	10.6	(10.1 - 11.1)		+	
Asthma attack in past year, among individuals with lifetime asthma	9.1	(0.0 - 30.2)	16.3	(15.1 - 17.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	17.1	(8.7 - 25.5)	11.1	(10.5 - 11.6)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	18.9	(10.8 - 27.1)	14.3	(13.7 - 14.9)			+
Sufficient vigorous activity	55.8	(45.2 - 66.5)	62.7	(62.0 - 63.5)			+
Sufficient moderate activity	24.5	(17.8 - 31.2)	26.1	(25.4 - 26.7)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	44.0	(35.6 - 52.3)	42.3	(41.5 - 43.1)			+
Described themselves as slightly or very overweight	29.7	(23.8 - 35.7)	29.8	(29.0 - 30.5)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	23.1	(15.1 - 31.1)	11.2	(10.5 - 11.8)	+		
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	31.2	(20.9 - 41.4)	23.1	(22.4 - 23.8)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	10.3	(2.4 - 18.3)	6.9	(6.4 - 7.3)			+
Purposely hurt themselves without wanting to die, during the past 12 months	10.6	(4.9 - 16.3)	12.7	(12.1 - 13.2)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	19.7	(14.2 - 25.3)	24.5	(23.8 - 25.2)			+

Hamilton County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Hamilton County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	8.0	(4.3 - 11.8)	6.1	(5.8 - 6.4)			+
Smoked cigars on one or more of the past 30 days	7.4	(3.3 - 11.4)	6.9	(6.6 - 7.2)			+
Used smokeless tobacco on one or more of the past 30 days	9.3	(5.4 - 13.1)	3.5	(3.2 - 3.7)	+		
Used any form of tobacco on one or more of the past 30 days	16.5	(11.5 - 21.4)	11.2	(10.8 - 11.7)			+
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	63.8	(56.4 - 71.3)	65.7	(65.1 - 66.3)			+
Exposed to secondhand smoke during the past 7 days	54.7	(47.0 - 62.4)	44.1	(43.4 - 44.7)	+		
Smoking is allowed in the home	14.9	(8.8 - 21.0)	9.4	(9.0 - 9.8)			+
Taught about tobacco use in any classes this year	49.4	(40.2 - 58.6)	43.2	(42.3 - 44.1)			+
Parent or guardian has talked about the dangers of tobacco in the past year	64.1	(57.8 - 70.5)	57.4	(56.8 - 58.0)			+
Served as an active member of SWAT	16.4	(11.1 - 21.6)	3.9	(3.7 - 4.1)	+		
<u>Definitely did not think</u> that smokers have more friends	30.6	(25.2 - 36.0)	32.0	(31.5 - 32.6)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	69.3	(64.0 - 74.5)	72.9	(72.4 - 73.4)			+

Hamilton County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Hamilton County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	16.7	(11.9 - 21.6)	20.5	(20.1 - 21.0)			+
Current asthma, among all individuals	10.1	(5.8 - 14.5)	11.3	(11.0 - 11.7)			+
Asthma attack in past year, among individuals with lifetime asthma	11.7	(0.0 - 25.3)	16.9	(16.0 - 17.8)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	15.5	(10.3 - 20.7)	11.1	(10.7 - 11.5)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	22.3	(16.7 - 27.9)	15.4	(15.0 - 15.9)	+		
Sufficient vigorous activity	57.9	(50.3 - 65.5)	66.5	(65.9 - 67.1)		+	
Sufficient moderate activity	20.2	(15.7 - 24.7)	25.3	(24.8 - 25.8)		+	
Exercised to lose weight or to keep from gaining weight during the past 30 days	45.1	(38.6 - 51.6)	43.5	(42.9 - 44.1)			+
Described themselves as slightly or very overweight	28.7	(23.2 - 34.3)	30.0	(29.5 - 30.6)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	21.6	(16.8 - 26.4)	10.9	(10.4 - 11.3)	+		
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	24.7	(18.5 - 30.9)	20.2	(19.7 - 20.7)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	7.3	(4.1 - 10.6)	6.9	(6.6 - 7.2)			+
Purposely hurt themselves without wanting to die, during the past 12 months	11.5	(7.4 - 15.6)	12.4	(12.0 - 12.8)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	20.5	(16.5 - 24.5)	22.0	(21.5 - 22.5)			+