

Hillsborough County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Hillsborough County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	2.8	(1.8 - 3.7)	3.3	(3.0 - 3.5)			+
Smoked cigars on one or more of the past 30 days	3.8	(2.4 - 5.3)	3.9	(3.6 - 4.2)			+
Used smokeless tobacco on one or more of the past 30 days	2.2	(1.0 - 3.5)	2.3	(2.1 - 2.6)			+
Used any form of tobacco on one or more of the past 30 days	5.5	(3.9 - 7.2)	6.4	(6.0 - 6.8)			+
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	72.0	(67.8 - 76.1)	72.0	(71.2 - 72.8)			+
Exposed to secondhand smoke during the past 7 days	35.4	(32.0 - 38.8)	41.3	(40.4 - 42.2)		+	
Smoking is allowed in the home	10.5	(8.3 - 12.7)	9.9	(9.5 - 10.4)			+
Taught about tobacco use in any classes this year	55.0	(49.5 - 60.5)	53.4	(52.0 - 54.7)			+
Parent or guardian has talked about the dangers of tobacco in the past year	59.9	(56.6 - 63.2)	63.9	(63.1 - 64.6)			+
Served as an active member of SWAT	5.5	(3.7 - 7.4)	4.3	(3.9 - 4.7)			+
<u>Definitely did not think</u> that smokers have more friends	36.9	(33.3 - 40.4)	33.7	(32.9 - 34.5)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	70.2	(66.8 - 73.6)	73.7	(73.0 - 74.4)			+

Hillsborough County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Hillsborough County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	22.0	(19.0 - 25.0)	20.1	(19.5 - 20.7)			+
Current asthma, among all individuals	12.0	(10.0 - 14.1)	12.0	(11.5 - 12.5)			+
Asthma attack in past year, among individuals with lifetime asthma	19.9	(14.0 - 25.7)	18.1	(16.8 - 19.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	11.9	(9.5 - 14.2)	11.6	(11.0 - 12.1)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	15.0	(12.3 - 17.7)	16.5	(15.8 - 17.2)			+
Sufficient vigorous activity	72.8	(69.4 - 76.2)	70.1	(69.2 - 70.9)			+
Sufficient moderate activity	29.8	(27.2 - 32.5)	24.8	(24.1 - 25.4)	+		
Exercised to lose weight or to keep from gaining weight during the past 30 days	43.1	(39.9 - 46.4)	44.4	(43.6 - 45.2)			+
Described themselves as slightly or very overweight	27.2	(24.3 - 30.1)	30.2	(29.4 - 30.9)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	11.7	(8.5 - 14.9)	11.4	(10.8 - 12.0)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	18.5	(15.4 - 21.6)	17.8	(17.2 - 18.5)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	10.9	(8.6 - 13.2)	7.8	(7.3 - 8.3)	+		
Purposely hurt themselves without wanting to die, during the past 12 months	13.3	(10.7 - 16.0)	12.6	(12.0 - 13.2)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	17.7	(15.1 - 20.3)	19.2	(18.5 - 19.8)			+

Hillsborough County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Hillsborough County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	7.4	(5.3 - 9.5)	10.1	(9.6 - 10.6)		+	
Smoked cigars on one or more of the past 30 days	10.2	(8.1 - 12.3)	11.4	(10.9 - 11.9)			+
Used smokeless tobacco on one or more of the past 30 days	4.6	(3.0 - 6.2)	5.6	(5.2 - 6.0)			+
Used any form of tobacco on one or more of the past 30 days	14.5	(11.7 - 17.3)	17.9	(17.3 - 18.6)		+	
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	64.6	(61.0 - 68.1)	58.4	(57.6 - 59.3)	+		
Exposed to secondhand smoke during the past 7 days	47.3	(43.4 - 51.2)	47.6	(46.7 - 48.4)			+
Smoking is allowed in the home	9.8	(7.2 - 12.4)	10.0	(9.5 - 10.5)			+
Taught about tobacco use in any classes this year	33.9	(28.8 - 39.1)	33.4	(32.4 - 34.4)			+
Parent or guardian has talked about the dangers of tobacco in the past year	46.7	(42.9 - 50.5)	50.4	(49.6 - 51.2)			+
Served as an active member of SWAT	3.5	(2.0 - 5.0)	4.0	(3.7 - 4.3)			+
<u>Definitely did not think</u> that smokers have more friends	30.9	(27.7 - 34.0)	30.5	(29.8 - 31.2)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	74.9	(71.8 - 78.0)	71.7	(71.0 - 72.4)			+

Hillsborough County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Hillsborough County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	24.1	(21.0 - 27.2)	20.8	(20.2 - 21.5)			+
Current asthma, among all individuals	12.0	(9.4 - 14.5)	10.6	(10.1 - 11.1)			+
Asthma attack in past year, among individuals with lifetime asthma	19.4	(14.5 - 24.3)	16.3	(15.1 - 17.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	11.4	(8.9 - 13.9)	11.1	(10.5 - 11.6)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	15.8	(12.8 - 18.8)	14.3	(13.7 - 14.9)			+
Sufficient vigorous activity	59.3	(55.5 - 63.0)	62.7	(62.0 - 63.5)			+
Sufficient moderate activity	25.6	(22.4 - 28.8)	26.1	(25.4 - 26.7)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	41.9	(38.0 - 45.8)	42.3	(41.5 - 43.1)			+
Described themselves as slightly or very overweight	30.5	(26.8 - 34.3)	29.8	(29.0 - 30.5)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	7.5	(5.2 - 9.8)	11.2	(10.5 - 11.8)		+	
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	19.6	(16.4 - 22.9)	23.1	(22.4 - 23.8)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	9.1	(6.9 - 11.3)	6.9	(6.4 - 7.3)			+
Purposely hurt themselves without wanting to die, during the past 12 months	12.3	(10.1 - 14.5)	12.7	(12.1 - 13.2)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	23.3	(20.0 - 26.6)	24.5	(23.8 - 25.2)			+

Hillsborough County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Hillsborough County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	4.5	(3.3 - 5.7)	6.1	(5.8 - 6.4)		+	
Smoked cigars on one or more of the past 30 days	5.8	(4.6 - 7.0)	6.9	(6.6 - 7.2)			+
Used smokeless tobacco on one or more of the past 30 days	2.7	(1.7 - 3.7)	3.5	(3.2 - 3.7)			+
Used any form of tobacco on one or more of the past 30 days	8.9	(7.2 - 10.5)	11.2	(10.8 - 11.7)		+	
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	68.9	(66.0 - 71.7)	65.7	(65.1 - 66.3)			+
Exposed to secondhand smoke during the past 7 days	41.1	(38.2 - 44.1)	44.1	(43.4 - 44.7)			+
Smoking is allowed in the home	9.4	(7.9 - 11.0)	9.4	(9.0 - 9.8)			+
Taught about tobacco use in any classes this year	44.3	(40.2 - 48.5)	43.2	(42.3 - 44.1)			+
Parent or guardian has talked about the dangers of tobacco in the past year	53.8	(51.2 - 56.5)	57.4	(56.8 - 58.0)		+	
Served as an active member of SWAT	4.0	(2.9 - 5.1)	3.9	(3.7 - 4.1)			+
<u>Definitely did not think</u> that smokers have more friends	33.3	(30.8 - 35.9)	32.0	(31.5 - 32.6)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	72.5	(70.0 - 75.0)	72.9	(72.4 - 73.4)			+

Hillsborough County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Hillsborough County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	22.9	(20.7 - 25.1)	20.5	(20.1 - 21.0)			+
Current asthma, among all individuals	12.1	(10.3 - 13.8)	11.3	(11.0 - 11.7)			+
Asthma attack in past year, among individuals with lifetime asthma	19.5	(15.6 - 23.5)	16.9	(16.0 - 17.8)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	12.1	(10.2 - 13.9)	11.1	(10.7 - 11.5)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	15.8	(13.7 - 18.0)	15.4	(15.0 - 15.9)			+
Sufficient vigorous activity	66.2	(63.4 - 69.1)	66.5	(65.9 - 67.1)			+
Sufficient moderate activity	27.0	(24.9 - 29.2)	25.3	(24.8 - 25.8)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	43.7	(41.0 - 46.4)	43.5	(42.9 - 44.1)			+
Described themselves as slightly or very overweight	29.2	(26.8 - 31.6)	30.0	(29.5 - 30.6)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	9.2	(7.2 - 11.2)	10.9	(10.4 - 11.3)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	18.5	(16.2 - 20.7)	20.2	(19.7 - 20.7)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	9.6	(8.0 - 11.2)	6.9	(6.6 - 7.2)	+		
Purposely hurt themselves without wanting to die, during the past 12 months	12.5	(10.7 - 14.3)	12.4	(12.0 - 12.8)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	20.7	(18.5 - 23.0)	22.0	(21.5 - 22.5)			+