

Indian River County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Indian River County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	3.4	(2.2 - 4.6)	3.3	(3.0 - 3.5)			+
Smoked cigars on one or more of the past 30 days	3.9	(2.1 - 5.8)	3.9	(3.6 - 4.2)			+
Used smokeless tobacco on one or more of the past 30 days	2.0	(0.9 - 3.2)	2.3	(2.1 - 2.6)			+
Used any form of tobacco on one or more of the past 30 days	7.4	(5.1 - 9.7)	6.4	(6.0 - 6.8)			+
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	69.5	(65.3 - 73.7)	72.0	(71.2 - 72.8)			+
Exposed to secondhand smoke during the past 7 days	46.1	(42.0 - 50.2)	41.3	(40.4 - 42.2)			+
Smoking is allowed in the home	9.4	(6.9 - 11.8)	9.9	(9.5 - 10.4)			+
Taught about tobacco use in any classes this year	65.6	(60.3 - 70.9)	53.4	(52.0 - 54.7)	+		
Parent or guardian has talked about the dangers of tobacco in the past year	64.4	(60.4 - 68.4)	63.9	(63.1 - 64.6)			+
Served as an active member of SWAT	6.9	(4.3 - 9.6)	4.3	(3.9 - 4.7)			+
<u>Definitely did not think</u> that smokers have more friends	31.8	(28.1 - 35.6)	33.7	(32.9 - 34.5)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	71.9	(68.6 - 75.3)	73.7	(73.0 - 74.4)			+

Indian River County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Indian River County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	17.2	(14.4 - 19.9)	20.1	(19.5 - 20.7)			+
Current asthma, among all individuals	10.0	(8.1 - 12.0)	12.0	(11.5 - 12.5)			+
Asthma attack in past year, among individuals with lifetime asthma	12.4	(6.3 - 18.4)	18.1	(16.8 - 19.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	9.2	(6.8 - 11.7)	11.6	(11.0 - 12.1)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	17.2	(13.9 - 20.4)	16.5	(15.8 - 17.2)			+
Sufficient vigorous activity	73.1	(69.6 - 76.7)	70.1	(69.2 - 70.9)			+
Sufficient moderate activity	28.5	(24.6 - 32.4)	24.8	(24.1 - 25.4)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	45.1	(40.8 - 49.3)	44.4	(43.6 - 45.2)			+
Described themselves as slightly or very overweight	31.7	(28.1 - 35.3)	30.2	(29.4 - 30.9)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	10.4	(7.7 - 13.1)	11.4	(10.8 - 12.0)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	21.0	(17.5 - 24.6)	17.8	(17.2 - 18.5)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	5.9	(3.9 - 7.9)	7.8	(7.3 - 8.3)			+
Purposely hurt themselves without wanting to die, during the past 12 months	10.7	(8.1 - 13.4)	12.6	(12.0 - 13.2)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	17.9	(15.2 - 20.5)	19.2	(18.5 - 19.8)			+

Indian River County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Indian River County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	13.9	(11.2 - 16.6)	10.1	(9.6 - 10.6)	+		
Smoked cigars on one or more of the past 30 days	16.2	(13.1 - 19.2)	11.4	(10.9 - 11.9)	+		
Used smokeless tobacco on one or more of the past 30 days	6.6	(4.7 - 8.5)	5.6	(5.2 - 6.0)			+
Used any form of tobacco on one or more of the past 30 days	24.3	(20.7 - 27.8)	17.9	(17.3 - 18.6)	+		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	53.9	(49.6 - 58.1)	58.4	(57.6 - 59.3)			+
Exposed to secondhand smoke during the past 7 days	51.9	(47.5 - 56.2)	47.6	(46.7 - 48.4)			+
Smoking is allowed in the home	9.9	(7.6 - 12.1)	10.0	(9.5 - 10.5)			+
Taught about tobacco use in any classes this year	37.9	(31.3 - 44.5)	33.4	(32.4 - 34.4)			+
Parent or guardian has talked about the dangers of tobacco in the past year	49.4	(45.2 - 53.5)	50.4	(49.6 - 51.2)			+
Served as an active member of SWAT	3.8	(1.9 - 5.7)	4.0	(3.7 - 4.3)			+
<u>Definitely did not think</u> that smokers have more friends	29.1	(25.7 - 32.5)	30.5	(29.8 - 31.2)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	71.4	(68.1 - 74.7)	71.7	(71.0 - 72.4)			+

Indian River County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Indian River County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	19.3	(16.4 - 22.1)	20.8	(20.2 - 21.5)			+
Current asthma, among all individuals	10.4	(7.8 - 13.0)	10.6	(10.1 - 11.1)			+
Asthma attack in past year, among individuals with lifetime asthma	17.8	(11.1 - 24.5)	16.3	(15.1 - 17.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	11.2	(8.8 - 13.6)	11.1	(10.5 - 11.6)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	14.2	(11.7 - 16.7)	14.3	(13.7 - 14.9)			+
Sufficient vigorous activity	63.6	(59.6 - 67.6)	62.7	(62.0 - 63.5)			+
Sufficient moderate activity	25.6	(22.3 - 28.8)	26.1	(25.4 - 26.7)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	41.2	(37.3 - 45.1)	42.3	(41.5 - 43.1)			+
Described themselves as slightly or very overweight	29.9	(26.4 - 33.4)	29.8	(29.0 - 30.5)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	13.9	(10.7 - 17.0)	11.2	(10.5 - 11.8)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	26.2	(22.0 - 30.3)	23.1	(22.4 - 23.8)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	7.0	(4.6 - 9.4)	6.9	(6.4 - 7.3)			+
Purposely hurt themselves without wanting to die, during the past 12 months	12.7	(9.9 - 15.5)	12.7	(12.1 - 13.2)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	21.7	(18.3 - 25.2)	24.5	(23.8 - 25.2)			+

Indian River County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Indian River County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	8.3	(6.6 - 10.1)	6.1	(5.8 - 6.4)	+		
Smoked cigars on one or more of the past 30 days	10.4	(8.2 - 12.6)	6.9	(6.6 - 7.2)	+		
Used smokeless tobacco on one or more of the past 30 days	4.4	(3.2 - 5.6)	3.5	(3.2 - 3.7)			+
Used any form of tobacco on one or more of the past 30 days	16.3	(13.7 - 18.9)	11.2	(10.8 - 11.7)	+		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	61.2	(57.9 - 64.6)	65.7	(65.1 - 66.3)		+	
Exposed to secondhand smoke during the past 7 days	49.3	(46.2 - 52.3)	44.1	(43.4 - 44.7)	+		
Smoking is allowed in the home	9.4	(7.7 - 11.1)	9.4	(9.0 - 9.8)			+
Taught about tobacco use in any classes this year	51.1	(46.2 - 56.0)	43.2	(42.3 - 44.1)	+		
Parent or guardian has talked about the dangers of tobacco in the past year	57.0	(54.0 - 60.1)	57.4	(56.8 - 58.0)			+
Served as an active member of SWAT	5.1	(3.6 - 6.7)	3.9	(3.7 - 4.1)			+
<u>Definitely did not think</u> that smokers have more friends	30.0	(27.4 - 32.6)	32.0	(31.5 - 32.6)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	71.1	(68.6 - 73.6)	72.9	(72.4 - 73.4)			+

Indian River County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Indian River County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	18.1	(15.9 - 20.2)	20.5	(20.1 - 21.0)			+
Current asthma, among all individuals	10.3	(8.7 - 12.0)	11.3	(11.0 - 11.7)			+
Asthma attack in past year, among individuals with lifetime asthma	15.9	(10.8 - 21.0)	16.9	(16.0 - 17.8)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	9.9	(8.1 - 11.7)	11.1	(10.7 - 11.5)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	15.8	(13.7 - 17.8)	15.4	(15.0 - 15.9)			+
Sufficient vigorous activity	68.3	(65.5 - 71.1)	66.5	(65.9 - 67.1)			+
Sufficient moderate activity	26.9	(24.3 - 29.6)	25.3	(24.8 - 25.8)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	43.2	(40.2 - 46.3)	43.5	(42.9 - 44.1)			+
Described themselves as slightly or very overweight	31.1	(28.5 - 33.6)	30.0	(29.5 - 30.6)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	12.0	(9.9 - 14.2)	10.9	(10.4 - 11.3)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	23.9	(20.9 - 26.9)	20.2	(19.7 - 20.7)	+		
Did not go to school on one or more of the past 30 days because they felt unsafe	5.9	(4.3 - 7.4)	6.9	(6.6 - 7.2)			+
Purposely hurt themselves without wanting to die, during the past 12 months	11.4	(9.6 - 13.3)	12.4	(12.0 - 12.8)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	20.0	(17.7 - 22.4)	22.0	(21.5 - 22.5)			+