

Jefferson County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Jefferson County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	2.7	(0.0 - 6.8)	3.3	(3.0 - 3.5)			+
Smoked cigars on one or more of the past 30 days	7.3	(0.9 - 13.7)	3.9	(3.6 - 4.2)			+
Used smokeless tobacco on one or more of the past 30 days	4.7	(0.0 - 10.9)	2.3	(2.1 - 2.6)			+
Used any form of tobacco on one or more of the past 30 days	10.5	(1.9 - 19.1)	6.4	(6.0 - 6.8)			+
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	61.0	(48.7 - 73.3)	72.0	(71.2 - 72.8)			+
Exposed to secondhand smoke during the past 7 days	36.4	(25.2 - 47.5)	41.3	(40.4 - 42.2)			+
Smoking is allowed in the home	17.6	(4.4 - 30.7)	9.9	(9.5 - 10.4)			+
Taught about tobacco use in any classes this year	53.0	(40.7 - 65.3)	53.4	(52.0 - 54.7)			+
Parent or guardian has talked about the dangers of tobacco in the past year	57.5	(44.7 - 70.3)	63.9	(63.1 - 64.6)			+
Served as an active member of SWAT	17.2	(6.4 - 27.9)	4.3	(3.9 - 4.7)	+		
<u>Definitely did not think</u> that smokers have more friends	35.5	(23.2 - 47.7)	33.7	(32.9 - 34.5)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	70.0	(57.1 - 82.9)	73.7	(73.0 - 74.4)			+

Jefferson County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Jefferson County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	25.5	(14.8 - 36.3)	20.1	(19.5 - 20.7)			+
Current asthma, among all individuals	12.0	(2.2 - 21.8)	12.0	(11.5 - 12.5)			+
Asthma attack in past year, among individuals with lifetime asthma	15.9	(0.0 - 38.6)	18.1	(16.8 - 19.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	24.4	(13.1 - 35.8)	11.6	(11.0 - 12.1)	+		
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	22.7	(6.2 - 39.3)	16.5	(15.8 - 17.2)			+
Sufficient vigorous activity	51.5	(43.1 - 59.9)	70.1	(69.2 - 70.9)		+	
Sufficient moderate activity	20.6	(11.4 - 29.9)	24.8	(24.1 - 25.4)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	45.3	(28.0 - 62.6)	44.4	(43.6 - 45.2)			+
Described themselves as slightly or very overweight	26.3	(12.3 - 40.3)	30.2	(29.4 - 30.9)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	15.9	(2.0 - 29.8)	11.4	(10.8 - 12.0)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	30.2	(18.9 - 41.5)	17.8	(17.2 - 18.5)	+		
Did not go to school on one or more of the past 30 days because they felt unsafe	8.5	(0.0 - 19.0)	7.8	(7.3 - 8.3)			+
Purposely hurt themselves without wanting to die, during the past 12 months	34.1	(18.3 - 49.9)	12.6	(12.0 - 13.2)	+		
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	30.2	(16.7 - 43.7)	19.2	(18.5 - 19.8)			+

Jefferson County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Jefferson County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	6.3	(0.5 - 12.1)	10.1	(9.6 - 10.6)			+
Smoked cigars on one or more of the past 30 days	13.4	(3.5 - 23.3)	11.4	(10.9 - 11.9)			+
Used smokeless tobacco on one or more of the past 30 days	9.0	(0.8 - 17.1)	5.6	(5.2 - 6.0)			+
Used any form of tobacco on one or more of the past 30 days	18.9	(10.0 - 27.8)	17.9	(17.3 - 18.6)			+
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	59.3	(48.2 - 70.5)	58.4	(57.6 - 59.3)			+
Exposed to secondhand smoke during the past 7 days	42.3	(24.2 - 60.4)	47.6	(46.7 - 48.4)			+
Smoking is allowed in the home	9.3	(1.8 - 16.8)	10.0	(9.5 - 10.5)			+
Taught about tobacco use in any classes this year	42.3	(31.1 - 53.5)	33.4	(32.4 - 34.4)			+
Parent or guardian has talked about the dangers of tobacco in the past year	54.4	(38.2 - 70.7)	50.4	(49.6 - 51.2)			+
Served as an active member of SWAT	20.0	(13.0 - 27.0)	4.0	(3.7 - 4.3)	+		
<u>Definitely did not think</u> that smokers have more friends	23.1	(8.8 - 37.4)	30.5	(29.8 - 31.2)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	64.1	(49.2 - 79.0)	71.7	(71.0 - 72.4)			+

Jefferson County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Jefferson County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	20.8	(11.0 - 30.6)	20.8	(20.2 - 21.5)			+
Current asthma, among all individuals	13.9	(5.1 - 22.7)	10.6	(10.1 - 11.1)			+
Asthma attack in past year, among individuals with lifetime asthma	8.6	(0.0 - 29.3)	16.3	(15.1 - 17.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	7.9	(0.0 - 17.1)	11.1	(10.5 - 11.6)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	22.0	(10.1 - 34.0)	14.3	(13.7 - 14.9)			+
Sufficient vigorous activity	71.2	(58.8 - 83.6)	62.7	(62.0 - 63.5)			+
Sufficient moderate activity	29.8	(16.6 - 43.0)	26.1	(25.4 - 26.7)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	36.6	(25.8 - 47.5)	42.3	(41.5 - 43.1)			+
Described themselves as slightly or very overweight	27.8	(15.6 - 40.0)	29.8	(29.0 - 30.5)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	26.3	(11.4 - 41.2)	11.2	(10.5 - 11.8)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	27.5	(10.2 - 44.7)	23.1	(22.4 - 23.8)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	4.1	(0.0 - 10.0)	6.9	(6.4 - 7.3)			+
Purposely hurt themselves without wanting to die, during the past 12 months	10.2	(0.0 - 26.5)	12.7	(12.1 - 13.2)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	22.3	(7.2 - 37.4)	24.5	(23.8 - 25.2)			+

Jefferson County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Jefferson County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	5.1	(1.2 - 9.0)	6.1	(5.8 - 6.4)			+
Smoked cigars on one or more of the past 30 days	10.1	(4.6 - 15.6)	6.9	(6.6 - 7.2)			+
Used smokeless tobacco on one or more of the past 30 days	6.2	(1.4 - 10.9)	3.5	(3.2 - 3.7)			+
Used any form of tobacco on one or more of the past 30 days	15.2	(8.6 - 21.7)	11.2	(10.8 - 11.7)			+
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	61.5	(54.0 - 69.1)	65.7	(65.1 - 66.3)			+
Exposed to secondhand smoke during the past 7 days	37.4	(27.8 - 47.0)	44.1	(43.4 - 44.7)			+
Smoking is allowed in the home	13.7	(5.6 - 21.7)	9.4	(9.0 - 9.8)			+
Taught about tobacco use in any classes this year	49.0	(40.4 - 57.6)	43.2	(42.3 - 44.1)			+
Parent or guardian has talked about the dangers of tobacco in the past year	56.9	(45.7 - 68.0)	57.4	(56.8 - 58.0)			+
Served as an active member of SWAT	20.9	(14.3 - 27.5)	3.9	(3.7 - 4.1)	+		
<u>Definitely did not think</u> that smokers have more friends	29.7	(21.0 - 38.3)	32.0	(31.5 - 32.6)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	68.1	(60.4 - 75.8)	72.9	(72.4 - 73.4)			+

Jefferson County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Jefferson County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	23.4	(15.7 - 31.1)	20.5	(20.1 - 21.0)			+
Current asthma, among all individuals	13.8	(7.2 - 20.5)	11.3	(11.0 - 11.7)			+
Asthma attack in past year, among individuals with lifetime asthma	13.1	(0.0 - 28.5)	16.9	(16.0 - 17.8)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	13.7	(6.5 - 20.9)	11.1	(10.7 - 11.5)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	25.1	(15.4 - 34.8)	15.4	(15.0 - 15.9)			+
Sufficient vigorous activity	62.1	(52.6 - 71.5)	66.5	(65.9 - 67.1)			+
Sufficient moderate activity	25.2	(17.1 - 33.3)	25.3	(24.8 - 25.8)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	41.4	(32.3 - 50.5)	43.5	(42.9 - 44.1)			+
Described themselves as slightly or very overweight	26.8	(18.2 - 35.3)	30.0	(29.5 - 30.6)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	21.8	(10.8 - 32.8)	10.9	(10.4 - 11.3)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	27.4	(18.0 - 36.9)	20.2	(19.7 - 20.7)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	6.6	(0.8 - 12.4)	6.9	(6.6 - 7.2)			+
Purposely hurt themselves without wanting to die, during the past 12 months	17.2	(6.3 - 28.1)	12.4	(12.0 - 12.8)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	25.2	(15.5 - 34.8)	22.0	(21.5 - 22.5)			+