

Leon County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Leon County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	3.0	(1.5 - 4.5)	3.3	(3.0 - 3.5)			+
Smoked cigars on one or more of the past 30 days	4.1	(2.0 - 6.3)	3.9	(3.6 - 4.2)			+
Used smokeless tobacco on one or more of the past 30 days	2.2	(0.8 - 3.6)	2.3	(2.1 - 2.6)			+
Used any form of tobacco on one or more of the past 30 days	6.3	(3.6 - 8.9)	6.4	(6.0 - 6.8)			+
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	74.1	(69.4 - 78.9)	72.0	(71.2 - 72.8)			+
Exposed to secondhand smoke during the past 7 days	30.3	(25.7 - 35.0)	41.3	(40.4 - 42.2)		+	
Smoking is allowed in the home	9.9	(6.8 - 13.0)	9.9	(9.5 - 10.4)			+
Taught about tobacco use in any classes this year	41.6	(35.7 - 47.4)	53.4	(52.0 - 54.7)		+	
Parent or guardian has talked about the dangers of tobacco in the past year	58.0	(54.2 - 61.8)	63.9	(63.1 - 64.6)		+	
Served as an active member of SWAT	6.0	(3.9 - 8.1)	4.3	(3.9 - 4.7)			+
<u>Definitely did not think</u> that smokers have more friends	30.3	(26.6 - 33.9)	33.7	(32.9 - 34.5)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	73.9	(69.1 - 78.6)	73.7	(73.0 - 74.4)			+

Leon County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Leon County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	18.1	(13.9 - 22.2)	20.1	(19.5 - 20.7)			+
Current asthma, among all individuals	11.6	(8.4 - 14.8)	12.0	(11.5 - 12.5)			+
Asthma attack in past year, among individuals with lifetime asthma	23.9	(14.4 - 33.4)	18.1	(16.8 - 19.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	13.8	(10.7 - 17.0)	11.6	(11.0 - 12.1)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	16.6	(12.2 - 21.0)	16.5	(15.8 - 17.2)			+
Sufficient vigorous activity	68.8	(64.4 - 73.3)	70.1	(69.2 - 70.9)			+
Sufficient moderate activity	23.4	(20.3 - 26.6)	24.8	(24.1 - 25.4)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	45.8	(40.8 - 50.7)	44.4	(43.6 - 45.2)			+
Described themselves as slightly or very overweight	29.6	(25.0 - 34.3)	30.2	(29.4 - 30.9)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	8.0	(5.1 - 10.8)	11.4	(10.8 - 12.0)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	19.3	(15.2 - 23.3)	17.8	(17.2 - 18.5)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	8.6	(5.9 - 11.3)	7.8	(7.3 - 8.3)			+
Purposely hurt themselves without wanting to die, during the past 12 months	13.7	(10.5 - 16.9)	12.6	(12.0 - 13.2)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	18.8	(14.8 - 22.7)	19.2	(18.5 - 19.8)			+

Leon County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Leon County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	8.9	(6.2 - 11.6)	10.1	(9.6 - 10.6)			+
Smoked cigars on one or more of the past 30 days	12.3	(9.3 - 15.4)	11.4	(10.9 - 11.9)			+
Used smokeless tobacco on one or more of the past 30 days	9.3	(7.1 - 11.6)	5.6	(5.2 - 6.0)	+		
Used any form of tobacco on one or more of the past 30 days	19.4	(15.7 - 23.2)	17.9	(17.3 - 18.6)			+
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	58.4	(53.9 - 62.8)	58.4	(57.6 - 59.3)			+
Exposed to secondhand smoke during the past 7 days	42.3	(38.2 - 46.4)	47.6	(46.7 - 48.4)		+	
Smoking is allowed in the home	9.4	(7.0 - 11.7)	10.0	(9.5 - 10.5)			+
Taught about tobacco use in any classes this year	40.2	(34.1 - 46.2)	33.4	(32.4 - 34.4)			+
Parent or guardian has talked about the dangers of tobacco in the past year	48.5	(44.4 - 52.6)	50.4	(49.6 - 51.2)			+
Served as an active member of SWAT	7.0	(5.0 - 9.0)	4.0	(3.7 - 4.3)	+		
<u>Definitely did not think</u> that smokers have more friends	30.4	(26.1 - 34.7)	30.5	(29.8 - 31.2)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	70.9	(67.1 - 74.7)	71.7	(71.0 - 72.4)			+

Leon County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Leon County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	20.6	(17.0 - 24.3)	20.8	(20.2 - 21.5)			+
Current asthma, among all individuals	11.1	(8.2 - 14.1)	10.6	(10.1 - 11.1)			+
Asthma attack in past year, among individuals with lifetime asthma	16.7	(9.9 - 23.5)	16.3	(15.1 - 17.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	10.8	(8.0 - 13.6)	11.1	(10.5 - 11.6)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	12.6	(9.6 - 15.6)	14.3	(13.7 - 14.9)			+
Sufficient vigorous activity	67.3	(64.0 - 70.6)	62.7	(62.0 - 63.5)	+		
Sufficient moderate activity	27.0	(23.0 - 31.0)	26.1	(25.4 - 26.7)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	40.5	(36.6 - 44.4)	42.3	(41.5 - 43.1)			+
Described themselves as slightly or very overweight	29.3	(25.7 - 33.0)	29.8	(29.0 - 30.5)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	10.8	(7.6 - 13.9)	11.2	(10.5 - 11.8)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	22.5	(18.9 - 26.1)	23.1	(22.4 - 23.8)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	5.1	(3.2 - 7.0)	6.9	(6.4 - 7.3)			+
Purposely hurt themselves without wanting to die, during the past 12 months	10.1	(7.3 - 12.8)	12.7	(12.1 - 13.2)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	18.7	(15.1 - 22.2)	24.5	(23.8 - 25.2)		+	

Leon County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Leon County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	4.5	(3.1 - 5.9)	6.1	(5.8 - 6.4)			+
Smoked cigars on one or more of the past 30 days	7.1	(5.3 - 8.8)	6.9	(6.6 - 7.2)			+
Used smokeless tobacco on one or more of the past 30 days	4.3	(3.1 - 5.5)	3.5	(3.2 - 3.7)			+
Used any form of tobacco on one or more of the past 30 days	10.9	(8.6 - 13.1)	11.2	(10.8 - 11.7)			+
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	67.7	(64.2 - 71.3)	65.7	(65.1 - 66.3)			+
Exposed to secondhand smoke during the past 7 days	35.6	(32.2 - 38.9)	44.1	(43.4 - 44.7)		+	
Smoking is allowed in the home	8.7	(6.9 - 10.6)	9.4	(9.0 - 9.8)			+
Taught about tobacco use in any classes this year	42.5	(38.0 - 47.0)	43.2	(42.3 - 44.1)			+
Parent or guardian has talked about the dangers of tobacco in the past year	53.8	(50.8 - 56.7)	57.4	(56.8 - 58.0)		+	
Served as an active member of SWAT	6.0	(4.6 - 7.5)	3.9	(3.7 - 4.1)	+		
<u>Definitely did not think</u> that smokers have more friends	31.3	(28.3 - 34.3)	32.0	(31.5 - 32.6)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	73.5	(70.5 - 76.4)	72.9	(72.4 - 73.4)			+

Leon County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Leon County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	18.4	(15.7 - 21.1)	20.5	(20.1 - 21.0)			+
Current asthma, among all individuals	11.1	(8.9 - 13.3)	11.3	(11.0 - 11.7)			+
Asthma attack in past year, among individuals with lifetime asthma	20.2	(14.4 - 26.0)	16.9	(16.0 - 17.8)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	12.2	(10.0 - 14.3)	11.1	(10.7 - 11.5)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	14.7	(12.1 - 17.4)	15.4	(15.0 - 15.9)			+
Sufficient vigorous activity	69.0	(66.2 - 71.8)	66.5	(65.9 - 67.1)			+
Sufficient moderate activity	25.1	(22.6 - 27.7)	25.3	(24.8 - 25.8)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	44.1	(40.8 - 47.3)	43.5	(42.9 - 44.1)			+
Described themselves as slightly or very overweight	30.2	(27.3 - 33.2)	30.0	(29.5 - 30.6)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	8.6	(6.5 - 10.8)	10.9	(10.4 - 11.3)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	19.6	(17.0 - 22.2)	20.2	(19.7 - 20.7)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	6.3	(4.6 - 8.0)	6.9	(6.6 - 7.2)			+
Purposely hurt themselves without wanting to die, during the past 12 months	11.3	(9.2 - 13.4)	12.4	(12.0 - 12.8)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	18.6	(16.0 - 21.3)	22.0	(21.5 - 22.5)		+	