

Madison County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Madison County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	7.4	(3.3 - 11.5)	3.3	(3.0 - 3.5)			+
Smoked cigars on one or more of the past 30 days	6.3	(3.0 - 9.6)	3.9	(3.6 - 4.2)			+
Used smokeless tobacco on one or more of the past 30 days	4.7	(1.3 - 8.1)	2.3	(2.1 - 2.6)			+
Used any form of tobacco on one or more of the past 30 days	11.9	(6.7 - 17.0)	6.4	(6.0 - 6.8)			+
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	63.9	(56.2 - 71.7)	72.0	(71.2 - 72.8)			+
Exposed to secondhand smoke during the past 7 days	43.8	(36.9 - 50.7)	41.3	(40.4 - 42.2)			+
Smoking is allowed in the home	15.5	(10.1 - 20.9)	9.9	(9.5 - 10.4)			+
Taught about tobacco use in any classes this year	42.3	(34.1 - 50.5)	53.4	(52.0 - 54.7)		+	
Parent or guardian has talked about the dangers of tobacco in the past year	55.3	(48.1 - 62.6)	63.9	(63.1 - 64.6)		+	
Served as an active member of SWAT	25.2	(17.5 - 32.9)	4.3	(3.9 - 4.7)	+		
<u>Definitely did not think</u> that smokers have more friends	35.6	(29.5 - 41.7)	33.7	(32.9 - 34.5)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	73.9	(66.6 - 81.1)	73.7	(73.0 - 74.4)			+

Madison County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Madison County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	16.5	(12.2 - 20.8)	20.1	(19.5 - 20.7)			+
Current asthma, among all individuals	11.2	(8.2 - 14.3)	12.0	(11.5 - 12.5)			+
Asthma attack in past year, among individuals with lifetime asthma	35.8	(17.5 - 54.1)	18.1	(16.8 - 19.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	20.5	(14.0 - 27.0)	11.6	(11.0 - 12.1)	+		
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	16.5	(10.5 - 22.5)	16.5	(15.8 - 17.2)			+
Sufficient vigorous activity	67.1	(61.7 - 72.5)	70.1	(69.2 - 70.9)			+
Sufficient moderate activity	22.4	(16.2 - 28.5)	24.8	(24.1 - 25.4)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	49.7	(42.4 - 57.0)	44.4	(43.6 - 45.2)			+
Described themselves as slightly or very overweight	31.2	(24.2 - 38.1)	30.2	(29.4 - 30.9)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	18.8	(14.5 - 23.1)	11.4	(10.8 - 12.0)	+		
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	14.5	(9.2 - 19.7)	17.8	(17.2 - 18.5)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	7.2	(3.8 - 10.6)	7.8	(7.3 - 8.3)			+
Purposely hurt themselves without wanting to die, during the past 12 months	14.0	(8.7 - 19.4)	12.6	(12.0 - 13.2)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	18.8	(14.1 - 23.4)	19.2	(18.5 - 19.8)			+

Madison County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Madison County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	11.7	(7.6 - 15.7)	10.1	(9.6 - 10.6)			+
Smoked cigars on one or more of the past 30 days	11.8	(7.4 - 16.2)	11.4	(10.9 - 11.9)			+
Used smokeless tobacco on one or more of the past 30 days	11.8	(7.1 - 16.5)	5.6	(5.2 - 6.0)	+		
Used any form of tobacco on one or more of the past 30 days	21.4	(16.7 - 26.2)	17.9	(17.3 - 18.6)			+
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	54.2	(47.0 - 61.3)	58.4	(57.6 - 59.3)			+
Exposed to secondhand smoke during the past 7 days	50.1	(42.4 - 57.8)	47.6	(46.7 - 48.4)			+
Smoking is allowed in the home	18.7	(11.5 - 25.9)	10.0	(9.5 - 10.5)	+		
Taught about tobacco use in any classes this year	38.5	(30.4 - 46.7)	33.4	(32.4 - 34.4)			+
Parent or guardian has talked about the dangers of tobacco in the past year	35.0	(25.5 - 44.5)	50.4	(49.6 - 51.2)		+	
Served as an active member of SWAT	15.1	(10.4 - 19.8)	4.0	(3.7 - 4.3)	+		
<u>Definitely did not think</u> that smokers have more friends	34.1	(27.8 - 40.4)	30.5	(29.8 - 31.2)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	62.2	(55.2 - 69.2)	71.7	(71.0 - 72.4)		+	

Madison County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Madison County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	15.8	(10.5 - 21.1)	20.8	(20.2 - 21.5)			+
Current asthma, among all individuals	8.1	(4.3 - 11.8)	10.6	(10.1 - 11.1)			+
Asthma attack in past year, among individuals with lifetime asthma	25.6	(3.7 - 47.5)	16.3	(15.1 - 17.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	15.9	(9.3 - 22.5)	11.1	(10.5 - 11.6)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	13.8	(7.9 - 19.8)	14.3	(13.7 - 14.9)			+
Sufficient vigorous activity	65.8	(58.1 - 73.5)	62.7	(62.0 - 63.5)			+
Sufficient moderate activity	27.4	(19.4 - 35.5)	26.1	(25.4 - 26.7)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	47.5	(38.7 - 56.4)	42.3	(41.5 - 43.1)			+
Described themselves as slightly or very overweight	31.5	(23.3 - 39.6)	29.8	(29.0 - 30.5)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	19.5	(13.8 - 25.3)	11.2	(10.5 - 11.8)	+		
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	22.6	(13.7 - 31.5)	23.1	(22.4 - 23.8)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	8.9	(4.2 - 13.5)	6.9	(6.4 - 7.3)			+
Purposely hurt themselves without wanting to die, during the past 12 months	15.1	(8.3 - 21.9)	12.7	(12.1 - 13.2)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	21.3	(14.6 - 27.9)	24.5	(23.8 - 25.2)			+

Madison County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Madison County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	9.3	(6.5 - 12.1)	6.1	(5.8 - 6.4)	+		
Smoked cigars on one or more of the past 30 days	8.8	(5.9 - 11.6)	6.9	(6.6 - 7.2)			+
Used smokeless tobacco on one or more of the past 30 days	7.9	(4.5 - 11.3)	3.5	(3.2 - 3.7)	+		
Used any form of tobacco on one or more of the past 30 days	15.9	(11.9 - 19.9)	11.2	(10.8 - 11.7)	+		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	59.6	(54.3 - 65.0)	65.7	(65.1 - 66.3)		+	
Exposed to secondhand smoke during the past 7 days	46.7	(41.9 - 51.5)	44.1	(43.4 - 44.7)			+
Smoking is allowed in the home	15.3	(10.9 - 19.7)	9.4	(9.0 - 9.8)	+		
Taught about tobacco use in any classes this year	39.4	(33.3 - 45.4)	43.2	(42.3 - 44.1)			+
Parent or guardian has talked about the dangers of tobacco in the past year	46.2	(39.2 - 53.3)	57.4	(56.8 - 58.0)		+	
Served as an active member of SWAT	19.5	(15.0 - 24.1)	3.9	(3.7 - 4.1)	+		
<u>Definitely did not think</u> that smokers have more friends	35.5	(30.0 - 40.9)	32.0	(31.5 - 32.6)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	66.7	(60.8 - 72.6)	72.9	(72.4 - 73.4)			+

Madison County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Madison County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	15.5	(12.2 - 18.8)	20.5	(20.1 - 21.0)		+	
Current asthma, among all individuals	9.5	(6.9 - 12.1)	11.3	(11.0 - 11.7)			+
Asthma attack in past year, among individuals with lifetime asthma	34.7	(20.0 - 49.5)	16.9	(16.0 - 17.8)	+		
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	17.2	(12.7 - 21.7)	11.1	(10.7 - 11.5)	+		
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	15.9	(11.2 - 20.5)	15.4	(15.0 - 15.9)			+
Sufficient vigorous activity	68.0	(63.2 - 72.8)	66.5	(65.9 - 67.1)			+
Sufficient moderate activity	23.7	(19.2 - 28.1)	25.3	(24.8 - 25.8)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	48.2	(42.4 - 54.1)	43.5	(42.9 - 44.1)			+
Described themselves as slightly or very overweight	31.1	(26.1 - 36.2)	30.0	(29.5 - 30.6)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	20.5	(16.2 - 24.9)	10.9	(10.4 - 11.3)	+		
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	19.2	(13.9 - 24.6)	20.2	(19.7 - 20.7)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	7.3	(4.2 - 10.5)	6.9	(6.6 - 7.2)			+
Purposely hurt themselves without wanting to die, during the past 12 months	14.9	(10.0 - 19.8)	12.4	(12.0 - 12.8)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	20.4	(15.8 - 25.0)	22.0	(21.5 - 22.5)			+