Martin County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Martin County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	**1.7**	(0.6 - 2.8)	**3.3**	(3.0 - 3.5)		+	
Smoked cigars on one or more of the past 30 days	**1.9**	(0.6 - 3.1)	**3.9**	(3.6 - 4.2)		+	
Used smokeless tobacco on one or more of the past 30 days	**2.3**	(1.0 - 3.7)	**2.3**	(2.1 - 2.6)			+
Used any form of tobacco on one or more of the past 30 days	**3.9**	(2.4 - 5.5)	**6.4**	(6.0 - 6.8)		+	
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	**73.0**	(67.8 - 78.2)	**72.0**	(71.2 - 72.8)			+
Exposed to secondhand smoke during the past 7 days	**38.6**	(33.9 - 43.3)	**41.3**	(40.4 - 42.2)			+
Smoking is allowed in the home	**6.5**	(4.1 - 8.9)	**9.9**	(9.5 - 10.4)		+	
Taught about tobacco use in any classes this year	**69.7**	(63.4 - 76.1)	**53.4**	(52.0 - 54.7)	+		
Parent or guardian has talked about the dangers of tobacco in the past year	**66.6**	(62.3 - 70.8)	**63.9**	(63.1 - 64.6)			+
Served as an active member of SWAT	**4.0**	(2.3 - 5.8)	**4.3**	(3.9 - 4.7)			+
Definitely did not think that smokers have more friends	**52.2**	(47.0 - 57.4)	**33.7**	(32.9 - 34.5)	+		
Definitely did not think that smoking helps young people “look cool” or “fit in”	**78.4**	(74.4 - 82.3)	**73.7**	(73.0 - 74.4)	+		

Martin County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Martin County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	**17.9**	(14.6 - 21.3)	**20.1**	(19.5 - 20.7)			+
Current asthma, among all individuals	**11.6**	(9.0 - 14.2)	**12.0**	(11.5 - 12.5)			+
Asthma attack in past year, among individuals with lifetime asthma	**22.4**	(14.1 - 30.6)	**18.1**	(16.8 - 19.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	**11.0**	(7.8 - 14.2)	**11.6**	(11.0 - 12.1)			+
Overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex)	**13.1**	(9.8 - 16.5)	**16.5**	(15.8 - 17.2)			+
Sufficient vigorous activity	**73.9**	(69.6 - 78.1)	**70.1**	(69.2 - 70.9)			+
Sufficient moderate activity	**30.2**	(26.4 - 34.1)	**24.8**	(24.1 - 25.4)	+		
Exercised to lose weight or to keep from gaining weight during the past 30 days	**40.0**	(34.7 - 45.2)	**44.4**	(43.6 - 45.2)			+
Described themselves as slightly or very overweight	**28.1**	(23.8 -32.3)	**30.2**	(29.4 - 30.9)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	**10.7**	(7.5 - 13.8)	**11.4**	(10.8 - 12.0)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	**17.1**	(13.4- 20.9)	**17.8**	(17.2- 18.5)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	**6.3**	(3.7 - 8.8)	**7.8**	(7.3 - 8.3)			+
Purposely hurt themselves without wanting to die, during the past 12 months	**7.9**	(5.5 - 10.4)	**12.6**	(12.0 - 13.2)		+	
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	**14.3**	(11.0 - 17.5)	**19.2**	(18.5 - 19.8)		+	

Martin County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Martin County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	**9.2**	(6.1 - 12.2)	**10.1**	(9.6 - 10.6)			+
Smoked cigars on one or more of the past 30 days	**8.8**	(5.9 - 11.8)	**11.4**	(10.9 - 11.9)			+
Used smokeless tobacco on one or more of the past 30 days	**5.5**	(3.1 - 8.0)	**5.6**	(5.2 - 6.0)			+
Used any form of tobacco on one or more of the past 30 days	**15.7**	(11.9 - 19.4)	**17.9**	(17.3 - 18.6)			+
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	**54.9**	(50.3 - 59.5)	**58.4**	(57.6 - 59.3)			+
Exposed to secondhand smoke during the past 7 days	**52.7**	(47.9 - 57.5)	**47.6**	(46.7 - 48.4)			+
Smoking is allowed in the home	**8.7**	(6.1 - 11.2)	**10.0**	(9.5 - 10.5)			+
Taught about tobacco use in any classes this year	**38.7**	(33.1 - 44.4)	**33.4**	(32.4 - 34.4)			+
Parent or guardian has talked about the dangers of tobacco in the past year	**51.9**	(47.3 - 56.6)	**50.4**	(49.6 - 51.2)			+
Served as an active member of SWAT	**4.8**	(2.8 - 6.7)	**4.0**	(3.7 - 4.3)			+
Definitely did not think that smokers have more friends	**33.7**	(29.1 - 38.2)	**30.5**	(29.8 - 31.2)			+
Definitely did not think that smoking helps young people “look cool” or “fit in”	**71.9**	(67.9 - 75.8)	**71.7**	(71.0 - 72.4)			+

Martin County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Martin County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	**16.0**	(12.7 - 19.2)	**20.8**	(20.2 - 21.5)		+	
Current asthma, among all individuals	**8.8**	(6.3 - 11.3)	**10.6**	(10.1 - 11.1)			+
Asthma attack in past year, among individuals with lifetime asthma	**16.7**	(7.9 - 25.6)	**16.3**	(15.1 - 17.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	**11.5**	(8.3 - 14.7)	**11.1**	(10.5 - 11.6)			+
Overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex)	**12.9**	(9.4 - 16.4)	**14.3**	(13.7 - 14.9)			+
Sufficient vigorous activity	**66.5**	(62.2 - 70.9)	**62.7**	(62.0 - 63.5)			+
Sufficient moderate activity	**26.9**	(22.8 - 31.0)	**26.1**	(25.4 - 26.7)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	**42.1**	(37.9 - 46.3)	**42.3**	(41.5 - 43.1)			+
Described themselves as slightly or very overweight	**29.8**	(25.9 - 33.7)	**29.8**	(29.0 - 30.5)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	**6.7**	(4.1 - 9.3)	**11.2**	(10.5 - 11.8)		+	
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	**20.8**	(16.8 - 24.8)	**23.1**	(22.4 - 23.8)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	**4.3**	(2.3 - 6.2)	**6.9**	(6.4 - 7.3)		+	
Purposely hurt themselves without wanting to die, during the past 12 months	**10.2**	(7.3 - 13.1)	**12.7**	(12.1 - 13.2)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	**20.9**	(17.7 - 24.2)	**24.5**	(23.8 - 25.2)			+

Martin County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Martin County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	**4.6**	(3.1 - 6.0)	**6.1**	(5.8 - 6.4)			+
Smoked cigars on one or more of the past 30 days	**4.3**	(2.9 - 5.6)	**6.9**	(6.6 - 7.2)		+	
Used smokeless tobacco on one or more of the past 30 days	**3.0**	(1.9 - 4.0)	**3.5**	(3.2 - 3.7)			+
Used any form of tobacco on one or more of the past 30 days	**8.3**	(6.3 - 10.3)	**11.2**	(10.8 - 11.7)		+	
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	**64.3**	(60.6 - 67.9)	**65.7**	(65.1 - 66.3)			+
Exposed to secondhand smoke during the past 7 days	**45.3**	(41.6 - 49.0)	**44.1**	(43.4 - 44.7)			+
Smoking is allowed in the home	**6.8**	(5.2 - 8.4)	**9.4**	(9.0 - 9.8)		+	
Taught about tobacco use in any classes this year	**52.9**	(47.8 - 58.0)	**43.2**	(42.3 - 44.1)	+		
Parent or guardian has talked about the dangers of tobacco in the past year	**58.8**	(55.3 - 62.4)	**57.4**	(56.8 - 58.0)			+
Served as an active member of SWAT	**4.5**	(3.1 - 6.0)	**3.9**	(3.7 - 4.1)			+
Definitely did not think that smokers have more friends	**43.3**	(39.5 - 47.2)	**32.0**	(31.5 - 32.6)	+		
Definitely did not think that smoking helps young people “look cool” or “fit in”	**75.5**	(72.7 - 78.4)	**72.9**	(72.4 - 73.4)			+

Martin County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Martin County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	**16.6**	(14.1 - 19.1)	**20.5**	(20.1 - 21.0)		+	
Current asthma, among all individuals	**9.8**	(8.0 - 11.6)	**11.3**	(11.0 - 11.7)			+
Asthma attack in past year, among individuals with lifetime asthma	**17.6**	(11.6 - 23.5)	**16.9**	(16.0 - 17.8)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	**11.2**	(8.8 - 13.6)	**11.1**	(10.7 - 11.5)			+
Overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex)	**12.9**	(10.4 - 15.5)	**15.4**	(15.0 - 15.9)			+
Sufficient vigorous activity	**70.7**	(67.4 - 74.0)	**66.5**	(65.9 - 67.1)	+		
Sufficient moderate activity	**29.5**	(26.7 - 32.4)	**25.3**	(24.8 - 25.8)	+		
Exercised to lose weight or to keep from gaining weight during the past 30 days	**40.5**	(37.2 - 43.9)	**43.5**	(42.9 - 44.1)			+
Described themselves as slightly or very overweight	**28.0**	(25.1 - 31.0)	**30.0**	(29.5 - 30.6)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	**7.3**	(5.4 - 9.1)	**10.9**	(10.4 - 11.3)		+	
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	**18.2**	(15.6 - 20.9)	**20.2**	(19.7 - 20.7)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	**4.7**	(3.3 - 6.2)	**6.9**	(6.6 - 7.2)		+	
Purposely hurt themselves without wanting to die, during the past 12 months	**8.8**	(7.0 - 10.6)	**12.4**	(12.0 - 12.8)		+	
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	**17.6**	(15.1 - 20.1)	**22.0**	(21.5 - 22.5)		+	