

## Okaloosa County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Okaloosa County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	<b>3.3</b>	(1.5 - 5.0)	<b>3.3</b>	(3.0 - 3.5)			<b>+</b>
Smoked cigars on one or more of the past 30 days	<b>2.3</b>	(0.7 - 3.9)	<b>3.9</b>	(3.6 - 4.2)			<b>+</b>
Used smokeless tobacco on one or more of the past 30 days	<b>2.0</b>	(0.9 - 3.1)	<b>2.3</b>	(2.1 - 2.6)			<b>+</b>
Used any form of tobacco on one or more of the past 30 days	<b>5.4</b>	(3.1 - 7.7)	<b>6.4</b>	(6.0 - 6.8)			<b>+</b>
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	<b>76.8</b>	(72.7 - 80.9)	<b>72.0</b>	(71.2 - 72.8)			<b>+</b>
Exposed to secondhand smoke during the past 7 days	<b>44.8</b>	(39.9 - 49.7)	<b>41.3</b>	(40.4 - 42.2)			<b>+</b>
Smoking is allowed in the home	<b>12.5</b>	(8.7 - 16.3)	<b>9.9</b>	(9.5 - 10.4)			<b>+</b>
Taught about tobacco use in any classes this year	<b>55.1</b>	(48.9 - 61.4)	<b>53.4</b>	(52.0 - 54.7)			<b>+</b>
Parent or guardian has talked about the dangers of tobacco in the past year	<b>62.3</b>	(58.6 - 66.0)	<b>63.9</b>	(63.1 - 64.6)			<b>+</b>
Served as an active member of SWAT	<b>2.7</b>	(1.0 - 4.5)	<b>4.3</b>	(3.9 - 4.7)			<b>+</b>
<u>Definitely did not think</u> that smokers have more friends	<b>39.7</b>	(35.4 - 44.0)	<b>33.7</b>	(32.9 - 34.5)	<b>+</b>		
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	<b>80.1</b>	(76.6 - 83.6)	<b>73.7</b>	(73.0 - 74.4)	<b>+</b>		

## Okaloosa County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Okaloosa County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	<b>20.8</b>	(17.6 - 24.0)	<b>20.1</b>	(19.5 - 20.7)			<b>+</b>
Current asthma, among all individuals	<b>14.6</b>	(11.6 - 17.5)	<b>12.0</b>	(11.5 - 12.5)			<b>+</b>
Asthma attack in past year, among individuals with lifetime asthma	<b>18.3</b>	(11.4 - 25.2)	<b>18.1</b>	(16.8 - 19.4)			<b>+</b>
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	<b>11.7</b>	(8.7 - 14.7)	<b>11.6</b>	(11.0 - 12.1)			<b>+</b>
Overweight (i.e., at or above the 85 <sup>th</sup> percentile but below the 95th percentile for body mass index, by age and sex)	<b>11.2</b>	(7.6 - 14.8)	<b>16.5</b>	(15.8 - 17.2)		<b>+</b>	
Sufficient vigorous activity	<b>74.5</b>	(70.4 - 78.7)	<b>70.1</b>	(69.2 - 70.9)			<b>+</b>
Sufficient moderate activity	<b>25.2</b>	(21.4 - 29.0)	<b>24.8</b>	(24.1 - 25.4)			<b>+</b>
Exercised to lose weight or to keep from gaining weight during the past 30 days	<b>38.1</b>	(33.8 - 42.5)	<b>44.4</b>	(43.6 - 45.2)		<b>+</b>	
Described themselves as slightly or very overweight	<b>26.5</b>	(23.2 - 29.8)	<b>30.2</b>	(29.4 - 30.9)			<b>+</b>
Never or rarely wore a seat belt when riding in a car driven by someone else	<b>7.2</b>	(5.0 - 9.5)	<b>11.4</b>	(10.8 - 12.0)		<b>+</b>	
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	<b>14.4</b>	(10.4 - 18.3)	<b>17.8</b>	(17.2 - 18.5)			<b>+</b>
Did not go to school on one or more of the past 30 days because they felt unsafe	<b>5.5</b>	(3.3 - 7.7)	<b>7.8</b>	(7.3 - 8.3)			<b>+</b>
Purposely hurt themselves without wanting to die, during the past 12 months	<b>12.7</b>	(9.9 - 15.4)	<b>12.6</b>	(12.0 - 13.2)			<b>+</b>
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	<b>18.5</b>	(14.8 - 22.2)	<b>19.2</b>	(18.5 - 19.8)			<b>+</b>

## Okaloosa County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Okaloosa County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	<b>10.7</b>	(8.2 - 13.3)	<b>10.1</b>	(9.6 - 10.6)			<b>+</b>
Smoked cigars on one or more of the past 30 days	<b>10.1</b>	(7.7 - 12.5)	<b>11.4</b>	(10.9 - 11.9)			<b>+</b>
Used smokeless tobacco on one or more of the past 30 days	<b>6.9</b>	(4.6 - 9.3)	<b>5.6</b>	(5.2 - 6.0)			<b>+</b>
Used any form of tobacco on one or more of the past 30 days	<b>18.9</b>	(15.5 - 22.4)	<b>17.9</b>	(17.3 - 18.6)			<b>+</b>
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	<b>58.8</b>	(54.7 - 62.9)	<b>58.4</b>	(57.6 - 59.3)			<b>+</b>
Exposed to secondhand smoke during the past 7 days	<b>48.9</b>	(44.8 - 53.1)	<b>47.6</b>	(46.7 - 48.4)			<b>+</b>
Smoking is allowed in the home	<b>7.6</b>	(5.7 - 9.5)	<b>10.0</b>	(9.5 - 10.5)			<b>+</b>
Taught about tobacco use in any classes this year	<b>37.0</b>	(31.7 - 42.3)	<b>33.4</b>	(32.4 - 34.4)			<b>+</b>
Parent or guardian has talked about the dangers of tobacco in the past year	<b>52.1</b>	(47.5 - 56.7)	<b>50.4</b>	(49.6 - 51.2)			<b>+</b>
Served as an active member of SWAT	<b>2.6</b>	(1.3 - 3.9)	<b>4.0</b>	(3.7 - 4.3)			<b>+</b>
<u>Definitely did not think</u> that smokers have more friends	<b>30.9</b>	(27.9 - 33.8)	<b>30.5</b>	(29.8 - 31.2)			<b>+</b>
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	<b>76.8</b>	(73.5 - 80.1)	<b>71.7</b>	(71.0 - 72.4)	<b>+</b>		

## Okaloosa County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Okaloosa County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	<b>21.2</b>	(18.1 - 24.3)	<b>20.8</b>	(20.2 - 21.5)			<b>+</b>
Current asthma, among all individuals	<b>11.7</b>	(9.3 - 14.1)	<b>10.6</b>	(10.1 - 11.1)			<b>+</b>
Asthma attack in past year, among individuals with lifetime asthma	<b>21.1</b>	(14.0 - 28.3)	<b>16.3</b>	(15.1 - 17.4)			<b>+</b>
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	<b>11.3</b>	(8.9 - 13.7)	<b>11.1</b>	(10.5 - 11.6)			<b>+</b>
Overweight (i.e., at or above the 85 <sup>th</sup> percentile but below the 95th percentile for body mass index, by age and sex)	<b>12.5</b>	(9.6 - 15.4)	<b>14.3</b>	(13.7 - 14.9)			<b>+</b>
Sufficient vigorous activity	<b>63.0</b>	(58.9 - 67.1)	<b>62.7</b>	(62.0 - 63.5)			<b>+</b>
Sufficient moderate activity	<b>28.0</b>	(24.5 - 31.5)	<b>26.1</b>	(25.4 - 26.7)			<b>+</b>
Exercised to lose weight or to keep from gaining weight during the past 30 days	<b>44.9</b>	(40.3 - 49.5)	<b>42.3</b>	(41.5 - 43.1)			<b>+</b>
Described themselves as slightly or very overweight	<b>31.7</b>	(27.7 - 35.6)	<b>29.8</b>	(29.0 - 30.5)			<b>+</b>
Never or rarely wore a seat belt when riding in a car driven by someone else	<b>8.7</b>	(5.9 - 11.6)	<b>11.2</b>	(10.5 - 11.8)			<b>+</b>
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	<b>18.8</b>	(15.5 - 22.1)	<b>23.1</b>	(22.4 - 23.8)		<b>+</b>	
Did not go to school on one or more of the past 30 days because they felt unsafe	<b>5.3</b>	(3.4 - 7.1)	<b>6.9</b>	(6.4 - 7.3)			<b>+</b>
Purposely hurt themselves without wanting to die, during the past 12 months	<b>13.5</b>	(10.6 - 16.4)	<b>12.7</b>	(12.1 - 13.2)			<b>+</b>
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	<b>27.0</b>	(22.9 - 31.0)	<b>24.5</b>	(23.8 - 25.2)			<b>+</b>

## Okaloosa County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Okaloosa County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	<b>6.4</b>	(4.8 - 8.0)	<b>6.1</b>	(5.8 - 6.4)			<b>+</b>
Smoked cigars on one or more of the past 30 days	<b>5.8</b>	(4.1 - 7.5)	<b>6.9</b>	(6.6 - 7.2)			<b>+</b>
Used smokeless tobacco on one or more of the past 30 days	<b>4.1</b>	(2.7 - 5.4)	<b>3.5</b>	(3.2 - 3.7)			<b>+</b>
Used any form of tobacco on one or more of the past 30 days	<b>11.3</b>	(8.9 - 13.7)	<b>11.2</b>	(10.8 - 11.7)			<b>+</b>
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	<b>68.1</b>	(64.9 - 71.2)	<b>65.7</b>	(65.1 - 66.3)			<b>+</b>
Exposed to secondhand smoke during the past 7 days	<b>46.4</b>	(43.0 - 49.8)	<b>44.1</b>	(43.4 - 44.7)			<b>+</b>
Smoking is allowed in the home	<b>9.3</b>	(7.3 - 11.3)	<b>9.4</b>	(9.0 - 9.8)			<b>+</b>
Taught about tobacco use in any classes this year	<b>45.1</b>	(40.5 - 49.7)	<b>43.2</b>	(42.3 - 44.1)			<b>+</b>
Parent or guardian has talked about the dangers of tobacco in the past year	<b>58.3</b>	(55.1 - 61.4)	<b>57.4</b>	(56.8 - 58.0)			<b>+</b>
Served as an active member of SWAT	<b>2.5</b>	(1.5 - 3.5)	<b>3.9</b>	(3.7 - 4.1)		<b>+</b>	
<u>Definitely did not think</u> that smokers have more friends	<b>35.4</b>	(32.6 - 38.1)	<b>32.0</b>	(31.5 - 32.6)	<b>+</b>		
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	<b>78.7</b>	(76.2 - 81.2)	<b>72.9</b>	(72.4 - 73.4)	<b>+</b>		

## Okaloosa County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Okaloosa County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	<b>21.2</b>	(18.9 - 23.4)	<b>20.5</b>	(20.1 - 21.0)			<b>+</b>
Current asthma, among all individuals	<b>13.4</b>	(11.4 - 15.4)	<b>11.3</b>	(11.0 - 11.7)			<b>+</b>
Asthma attack in past year, among individuals with lifetime asthma	<b>19.2</b>	(14.2 - 24.3)	<b>16.9</b>	(16.0 - 17.8)			<b>+</b>
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	<b>11.0</b>	(9.0 - 12.9)	<b>11.1</b>	(10.7 - 11.5)			<b>+</b>
Overweight (i.e., at or above the 85 <sup>th</sup> percentile but below the 95th percentile for body mass index, by age and sex)	<b>11.9</b>	(9.7 - 14.1)	<b>15.4</b>	(15.0 - 15.9)		<b>+</b>	
Sufficient vigorous activity	<b>68.9</b>	(65.6 - 72.1)	<b>66.5</b>	(65.9 - 67.1)			<b>+</b>
Sufficient moderate activity	<b>26.5</b>	(23.8 - 29.2)	<b>25.3</b>	(24.8 - 25.8)			<b>+</b>
Exercised to lose weight or to keep from gaining weight during the past 30 days	<b>41.1</b>	(37.7 - 44.6)	<b>43.5</b>	(42.9 - 44.1)			<b>+</b>
Described themselves as slightly or very overweight	<b>29.1</b>	(26.3 - 31.9)	<b>30.0</b>	(29.5 - 30.6)			<b>+</b>
Never or rarely wore a seat belt when riding in a car driven by someone else	<b>7.3</b>	(5.6 - 9.1)	<b>10.9</b>	(10.4 - 11.3)		<b>+</b>	
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	<b>16.9</b>	(14.1 - 19.7)	<b>20.2</b>	(19.7 - 20.7)			<b>+</b>
Did not go to school on one or more of the past 30 days because they felt unsafe	<b>4.9</b>	(3.4 - 6.3)	<b>6.9</b>	(6.6 - 7.2)		<b>+</b>	
Purposely hurt themselves without wanting to die, during the past 12 months	<b>12.7</b>	(10.6 - 14.8)	<b>12.4</b>	(12.0 - 12.8)			<b>+</b>
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	<b>22.6</b>	(19.6 - 25.6)	<b>22.0</b>	(21.5 - 22.5)			<b>+</b>