

Pasco County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

| Indicator | Middle School | | | | Comparison of Two Rates | | |
|--|----------------|---------------|-------------|---------------|--------------------------|--------------------------|---------------|
| | Pasco County % | 95% CI | State % | 95% CI | County Higher Than State | State Higher Than County | No Difference |
| Smoked cigarettes on one or more of the past 30 days | 4.1 | (2.4 - 5.8) | 3.3 | (3.0 - 3.5) | | | + |
| Smoked cigars on one or more of the past 30 days | 2.8 | (1.5 - 4.0) | 3.9 | (3.6 - 4.2) | | | + |
| Used smokeless tobacco on one or more of the past 30 days | 2.7 | (1.3 - 4.2) | 2.3 | (2.1 - 2.6) | | | + |
| Used any form of tobacco on one or more of the past 30 days | 6.7 | (4.5 - 8.9) | 6.4 | (6.0 - 6.8) | | | + |
| Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one | 72.2 | (68.6 - 75.8) | 72.0 | (71.2 - 72.8) | | | + |
| Exposed to secondhand smoke during the past 7 days | 49.9 | (45.8 - 54.0) | 41.3 | (40.4 - 42.2) | + | | |
| Smoking is allowed in the home | 14.0 | (11.3 - 16.8) | 9.9 | (9.5 - 10.4) | + | | |
| Taught about tobacco use in any classes this year | 54.3 | (50.3 - 58.4) | 53.4 | (52.0 - 54.7) | | | + |
| Parent or guardian has talked about the dangers of tobacco in the past year | 63.5 | (60.7 - 66.2) | 63.9 | (63.1 - 64.6) | | | + |
| Served as an active member of SWAT | 5.9 | (4.2 - 7.5) | 4.3 | (3.9 - 4.7) | | | + |
| <u>Definitely did not think</u> that smokers have more friends | 34.0 | (30.7 - 37.4) | 33.7 | (32.9 - 34.5) | | | + |
| <u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in" | 78.0 | (75.0 - 81.1) | 73.7 | (73.0 - 74.4) | + | | |

Pasco County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

| Indicator | Middle School | | | | Comparison of Two Rates | | |
|---|----------------|---------------|-------------|---------------|--------------------------|--------------------------|---------------|
| | Pasco County % | 95% CI | State % | 95% CI | County Higher Than State | State Higher Than County | No Difference |
| Lifetime asthma | 23.0 | (20.7 - 25.3) | 20.1 | (19.5 - 20.7) | + | | |
| Current asthma, among all individuals | 13.5 | (11.3 - 15.6) | 12.0 | (11.5 - 12.5) | | | + |
| Asthma attack in past year, among individuals with lifetime asthma | 19.2 | (13.9 - 24.6) | 18.1 | (16.8 - 19.4) | | | + |
| Obese (i.e., at or above the 95th percentile for body mass index, by age and sex) | 10.7 | (8.4 - 13.0) | 11.6 | (11.0 - 12.1) | | | + |
| Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex) | 15.8 | (13.3 - 18.4) | 16.5 | (15.8 - 17.2) | | | + |
| Sufficient vigorous activity | 72.7 | (69.7 - 75.8) | 70.1 | (69.2 - 70.9) | | | + |
| Sufficient moderate activity | 27.9 | (25.2 - 30.7) | 24.8 | (24.1 - 25.4) | | | + |
| Exercised to lose weight or to keep from gaining weight during the past 30 days | 44.2 | (40.6 - 47.9) | 44.4 | (43.6 - 45.2) | | | + |
| Described themselves as slightly or very overweight | 29.8 | (26.4 - 33.2) | 30.2 | (29.4 - 30.9) | | | + |
| Never or rarely wore a seat belt when riding in a car driven by someone else | 8.0 | (6.1 - 9.9) | 11.4 | (10.8 - 12.0) | | + | |
| Rode in a car driven by someone who had been drinking alcohol during the past 30 days | 15.9 | (13.4 - 18.4) | 17.8 | (17.2 - 18.5) | | | + |
| Did not go to school on one or more of the past 30 days because they felt unsafe | 6.6 | (4.6 - 8.5) | 7.8 | (7.3 - 8.3) | | | + |
| Purposely hurt themselves without wanting to die, during the past 12 months | 11.8 | (9.7 - 14.0) | 12.6 | (12.0 - 13.2) | | | + |
| Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months | 19.8 | (16.9 - 22.8) | 19.2 | (18.5 - 19.8) | | | + |

Pasco County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

| Indicator | High School | | | | Comparison of Two Rates | | |
|--|----------------|---------------|-------------|---------------|--------------------------|--------------------------|---------------|
| | Pasco County % | 95% CI | State % | 95% CI | County Higher Than State | State Higher Than County | No Difference |
| Smoked cigarettes on one or more of the past 30 days | 11.8 | (9.2 - 14.4) | 10.1 | (9.6 - 10.6) | | | + |
| Smoked cigars on one or more of the past 30 days | 13.4 | (10.9 - 15.9) | 11.4 | (10.9 - 11.9) | | | + |
| Used smokeless tobacco on one or more of the past 30 days | 5.9 | (4.2 - 7.6) | 5.6 | (5.2 - 6.0) | | | + |
| Used any form of tobacco on one or more of the past 30 days | 19.8 | (16.8 - 22.8) | 17.9 | (17.3 - 18.6) | | | + |
| Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one | 55.8 | (52.1 - 59.6) | 58.4 | (57.6 - 59.3) | | | + |
| Exposed to secondhand smoke during the past 7 days | 53.3 | (48.8 - 57.8) | 47.6 | (46.7 - 48.4) | + | | |
| Smoking is allowed in the home | 12.3 | (9.3 - 15.3) | 10.0 | (9.5 - 10.5) | | | + |
| Taught about tobacco use in any classes this year | 29.4 | (25.0 - 33.7) | 33.4 | (32.4 - 34.4) | | | + |
| Parent or guardian has talked about the dangers of tobacco in the past year | 50.6 | (46.5 - 54.7) | 50.4 | (49.6 - 51.2) | | | + |
| Served as an active member of SWAT | 2.1 | (1.0 - 3.1) | 4.0 | (3.7 - 4.3) | | + | |
| <u>Definitely did not think</u> that smokers have more friends | 30.7 | (27.3 - 34.0) | 30.5 | (29.8 - 31.2) | | | + |
| <u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in" | 75.3 | (72.4 - 78.2) | 71.7 | (71.0 - 72.4) | + | | |

Pasco County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

| Indicator | High School | | | | Comparison of Two Rates | | |
|---|----------------|---------------|-------------|---------------|--------------------------|--------------------------|---------------|
| | Pasco County % | 95% CI | State % | 95% CI | County Higher Than State | State Higher Than County | No Difference |
| Lifetime asthma | 21.4 | (18.4 - 24.5) | 20.8 | (20.2 - 21.5) | | | + |
| Current asthma, among all individuals | 11.5 | (9.1 - 13.8) | 10.6 | (10.1 - 11.1) | | | + |
| Asthma attack in past year, among individuals with lifetime asthma | 22.5 | (15.2 - 29.8) | 16.3 | (15.1 - 17.4) | | | + |
| Obese (i.e., at or above the 95th percentile for body mass index, by age and sex) | 10.5 | (7.9 - 13.2) | 11.1 | (10.5 - 11.6) | | | + |
| Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex) | 14.0 | (11.4 - 16.5) | 14.3 | (13.7 - 14.9) | | | + |
| Sufficient vigorous activity | 62.8 | (59.3 - 66.3) | 62.7 | (62.0 - 63.5) | | | + |
| Sufficient moderate activity | 27.5 | (24.3 - 30.6) | 26.1 | (25.4 - 26.7) | | | + |
| Exercised to lose weight or to keep from gaining weight during the past 30 days | 41.7 | (38.2 - 45.2) | 42.3 | (41.5 - 43.1) | | | + |
| Described themselves as slightly or very overweight | 28.6 | (24.9 - 32.2) | 29.8 | (29.0 - 30.5) | | | + |
| Never or rarely wore a seat belt when riding in a car driven by someone else | 7.8 | (5.5 - 10.2) | 11.2 | (10.5 - 11.8) | | + | |
| Rode in a car driven by someone who had been drinking alcohol during the past 30 days | 21.8 | (18.6 - 25.0) | 23.1 | (22.4 - 23.8) | | | + |
| Did not go to school on one or more of the past 30 days because they felt unsafe | 5.9 | (4.1 - 7.7) | 6.9 | (6.4 - 7.3) | | | + |
| Purposely hurt themselves without wanting to die, during the past 12 months | 16.4 | (13.6 - 19.2) | 12.7 | (12.1 - 13.2) | + | | |
| Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months | 25.4 | (22.4 - 28.5) | 24.5 | (23.8 - 25.2) | | | + |

Pasco County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

| Indicator | Youth (Ages 11-17) | | | | Comparison of Two Rates | | |
|--|--------------------|---------------|-------------|---------------|--------------------------|--------------------------|---------------|
| | Pasco County % | 95% CI | State % | 95% CI | County Higher Than State | State Higher Than County | No Difference |
| Smoked cigarettes on one or more of the past 30 days | 7.1 | (5.5 - 8.8) | 6.1 | (5.8 - 6.4) | | | + |
| Smoked cigars on one or more of the past 30 days | 7.5 | (5.9 - 9.1) | 6.9 | (6.6 - 7.2) | | | + |
| Used smokeless tobacco on one or more of the past 30 days | 3.8 | (2.7 - 4.9) | 3.5 | (3.2 - 3.7) | | | + |
| Used any form of tobacco on one or more of the past 30 days | 12.1 | (10.1 - 14.1) | 11.2 | (10.8 - 11.7) | | | + |
| Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one | 64.5 | (61.7 - 67.3) | 65.7 | (65.1 - 66.3) | | | + |
| Exposed to secondhand smoke during the past 7 days | 50.8 | (47.7 - 53.9) | 44.1 | (43.4 - 44.7) | + | | |
| Smoking is allowed in the home | 12.2 | (10.2 - 14.2) | 9.4 | (9.0 - 9.8) | + | | |
| Taught about tobacco use in any classes this year | 41.0 | (37.4 - 44.7) | 43.2 | (42.3 - 44.1) | | | + |
| Parent or guardian has talked about the dangers of tobacco in the past year | 57.2 | (54.4 - 59.9) | 57.4 | (56.8 - 58.0) | | | + |
| Served as an active member of SWAT | 3.9 | (2.9 - 4.9) | 3.9 | (3.7 - 4.1) | | | + |
| <u>Definitely did not think</u> that smokers have more friends | 32.1 | (29.6 - 34.6) | 32.0 | (31.5 - 32.6) | | | + |
| <u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in" | 77.0 | (74.8 - 79.2) | 72.9 | (72.4 - 73.4) | + | | |

Pasco County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

| Indicator | Youth (Ages 11-17) | | | | Comparison of Two Rates | | |
|---|--------------------|---------------|-------------|---------------|--------------------------|--------------------------|---------------|
| | Pasco County % | 95% CI | State % | 95% CI | County Higher Than State | State Higher Than County | No Difference |
| Lifetime asthma | 22.4 | (20.4 - 24.5) | 20.5 | (20.1 - 21.0) | | | + |
| Current asthma, among all individuals | 12.6 | (10.9 - 14.4) | 11.3 | (11.0 - 11.7) | | | + |
| Asthma attack in past year, among individuals with lifetime asthma | 21.7 | (16.9 - 26.5) | 16.9 | (16.0 - 17.8) | | | + |
| Obese (i.e., at or above the 95th percentile for body mass index, by age and sex) | 10.5 | (8.6 - 12.4) | 11.1 | (10.7 - 11.5) | | | + |
| Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex) | 15.1 | (13.2 - 17.1) | 15.4 | (15.0 - 15.9) | | | + |
| Sufficient vigorous activity | 67.6 | (65.0 - 70.2) | 66.5 | (65.9 - 67.1) | | | + |
| Sufficient moderate activity | 27.1 | (24.9 - 29.2) | 25.3 | (24.8 - 25.8) | | | + |
| Exercised to lose weight or to keep from gaining weight during the past 30 days | 42.4 | (39.7 - 45.0) | 43.5 | (42.9 - 44.1) | | | + |
| Described themselves as slightly or very overweight | 28.6 | (25.9 - 31.3) | 30.0 | (29.5 - 30.6) | | | + |
| Never or rarely wore a seat belt when riding in a car driven by someone else | 7.8 | (6.2 - 9.4) | 10.9 | (10.4 - 11.3) | | + | |
| Rode in a car driven by someone who had been drinking alcohol during the past 30 days | 18.7 | (16.6 - 20.8) | 20.2 | (19.7 - 20.7) | | | + |
| Did not go to school on one or more of the past 30 days because they felt unsafe | 6.0 | (4.6 - 7.3) | 6.9 | (6.6 - 7.2) | | | + |
| Purposely hurt themselves without wanting to die, during the past 12 months | 14.3 | (12.5 - 16.2) | 12.4 | (12.0 - 12.8) | | | + |
| Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months | 23.2 | (20.9 - 25.5) | 22.0 | (21.5 - 22.5) | | | + |