

Putnam County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Putnam County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	5.9	(3.8 - 8.1)	3.3	(3.0 - 3.5)	+		
Smoked cigars on one or more of the past 30 days	4.5	(2.6 - 6.5)	3.9	(3.6 - 4.2)			+
Used smokeless tobacco on one or more of the past 30 days	6.2	(3.7 - 8.7)	2.3	(2.1 - 2.6)	+		
Used any form of tobacco on one or more of the past 30 days	11.7	(8.6 - 14.9)	6.4	(6.0 - 6.8)	+		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	63.4	(57.7 - 69.1)	72.0	(71.2 - 72.8)		+	
Exposed to secondhand smoke during the past 7 days	53.3	(48.3 - 58.3)	41.3	(40.4 - 42.2)	+		
Smoking is allowed in the home	22.5	(18.7 - 26.2)	9.9	(9.5 - 10.4)	+		
Taught about tobacco use in any classes this year	62.3	(56.4 - 68.2)	53.4	(52.0 - 54.7)	+		
Parent or guardian has talked about the dangers of tobacco in the past year	59.7	(54.9 - 64.6)	63.9	(63.1 - 64.6)			+
Served as an active member of SWAT	11.2	(8.5 - 13.9)	4.3	(3.9 - 4.7)	+		
<u>Definitely did not think</u> that smokers have more friends	40.8	(35.8 - 45.8)	33.7	(32.9 - 34.5)	+		
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	72.6	(68.8 - 76.5)	73.7	(73.0 - 74.4)			+

Putnam County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Putnam County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	21.3	(17.4 - 25.2)	20.1	(19.5 - 20.7)			+
Current asthma, among all individuals	12.6	(9.6 - 15.6)	12.0	(11.5 - 12.5)			+
Asthma attack in past year, among individuals with lifetime asthma	14.2	(7.5 - 20.8)	18.1	(16.8 - 19.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	16.3	(12.6 - 19.9)	11.6	(11.0 - 12.1)	+		
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	21.4	(17.9 - 24.8)	16.5	(15.8 - 17.2)	+		
Sufficient vigorous activity	64.0	(59.7 - 68.3)	70.1	(69.2 - 70.9)		+	
Sufficient moderate activity	22.7	(18.7 - 26.8)	24.8	(24.1 - 25.4)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	46.3	(42.8 - 49.8)	44.4	(43.6 - 45.2)			+
Described themselves as slightly or very overweight	37.3	(32.7 - 41.8)	30.2	(29.4 - 30.9)	+		
Never or rarely wore a seat belt when riding in a car driven by someone else	17.8	(14.0 - 21.7)	11.4	(10.8 - 12.0)	+		
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	24.5	(20.5 - 28.6)	17.8	(17.2 - 18.5)	+		
Did not go to school on one or more of the past 30 days because they felt unsafe	11.8	(8.8 - 14.8)	7.8	(7.3 - 8.3)	+		
Purposely hurt themselves without wanting to die, during the past 12 months	14.9	(11.6 - 18.1)	12.6	(12.0 - 13.2)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	23.3	(19.2 - 27.4)	19.2	(18.5 - 19.8)			+

Putnam County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Putnam County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	15.1	(4.8 - 25.5)	10.1	(9.6 - 10.6)			+
Smoked cigars on one or more of the past 30 days	25.8	(16.1 - 35.6)	11.4	(10.9 - 11.9)	+		
Used smokeless tobacco on one or more of the past 30 days	13.0	(2.3 - 23.8)	5.6	(5.2 - 6.0)			+
Used any form of tobacco on one or more of the past 30 days	33.7	(23.8 - 43.6)	17.9	(17.3 - 18.6)	+		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	46.3	(33.5 - 59.0)	58.4	(57.6 - 59.3)			+
Exposed to secondhand smoke during the past 7 days	49.4	(35.4 - 63.4)	47.6	(46.7 - 48.4)			+
Smoking is allowed in the home	18.0	(11.4 - 24.5)	10.0	(9.5 - 10.5)	+		
Taught about tobacco use in any classes this year	38.4	(25.6 - 51.1)	33.4	(32.4 - 34.4)			+
Parent or guardian has talked about the dangers of tobacco in the past year	48.9	(34.5 - 63.3)	50.4	(49.6 - 51.2)			+
Served as an active member of SWAT	10.6	(4.4 - 16.9)	4.0	(3.7 - 4.3)	+		
<u>Definitely did not think</u> that smokers have more friends	34.4	(22.0 - 46.9)	30.5	(29.8 - 31.2)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	65.6	(54.3 - 76.8)	71.7	(71.0 - 72.4)			+

Putnam County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Putnam County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	25.8	(11.5 - 40.2)	20.8	(20.2 - 21.5)			+
Current asthma, among all individuals	10.3	(3.4 - 17.3)	10.6	(10.1 - 11.1)			+
Asthma attack in past year, among individuals with lifetime asthma	17.4	(1.8 - 33.0)	16.3	(15.1 - 17.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	18.1	(10.4 - 25.7)	11.1	(10.5 - 11.6)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	12.5	(1.6 - 23.4)	14.3	(13.7 - 14.9)			+
Sufficient vigorous activity	54.3	(42.1 - 66.5)	62.7	(62.0 - 63.5)			+
Sufficient moderate activity	23.7	(15.0 - 32.4)	26.1	(25.4 - 26.7)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	25.8	(15.5 - 36.2)	42.3	(41.5 - 43.1)		+	
Described themselves as slightly or very overweight	25.8	(18.2 - 33.5)	29.8	(29.0 - 30.5)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	32.6	(23.1 - 42.1)	11.2	(10.5 - 11.8)	+		
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	33.7	(23.0 - 44.4)	23.1	(22.4 - 23.8)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	20.0	(8.7 - 31.3)	6.9	(6.4 - 7.3)	+		
Purposely hurt themselves without wanting to die, during the past 12 months	18.9	(9.0 - 28.8)	12.7	(12.1 - 13.2)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	28.1	(18.7 - 37.5)	24.5	(23.8 - 25.2)			+

Putnam County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Putnam County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	6.1	(4.0 - 8.2)	6.1	(5.8 - 6.4)			+
Smoked cigars on one or more of the past 30 days	6.3	(3.9 - 8.6)	6.9	(6.6 - 7.2)			+
Used smokeless tobacco on one or more of the past 30 days	6.5	(4.2 - 8.9)	3.5	(3.2 - 3.7)	+		
Used any form of tobacco on one or more of the past 30 days	13.6	(10.3 - 16.8)	11.2	(10.8 - 11.7)			+
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	61.8	(56.4 - 67.2)	65.7	(65.1 - 66.3)			+
Exposed to secondhand smoke during the past 7 days	52.9	(48.1 - 57.7)	44.1	(43.4 - 44.7)	+		
Smoking is allowed in the home	21.2	(17.7 - 24.7)	9.4	(9.0 - 9.8)	+		
Taught about tobacco use in any classes this year	59.0	(53.1 - 64.9)	43.2	(42.3 - 44.1)	+		
Parent or guardian has talked about the dangers of tobacco in the past year	58.1	(53.4 - 62.8)	57.4	(56.8 - 58.0)			+
Served as an active member of SWAT	10.9	(8.4 - 13.4)	3.9	(3.7 - 4.1)	+		
<u>Definitely did not think</u> that smokers have more friends	40.6	(35.9 - 45.3)	32.0	(31.5 - 32.6)	+		
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	72.7	(69.1 - 76.3)	72.9	(72.4 - 73.4)			+

Putnam County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Putnam County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	21.9	(18.1 - 25.6)	20.5	(20.1 - 21.0)			+
Current asthma, among all individuals	12.5	(9.7 - 15.2)	11.3	(11.0 - 11.7)			+
Asthma attack in past year, among individuals with lifetime asthma	13.8	(8.1 - 19.5)	16.9	(16.0 - 17.8)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	16.2	(12.8 - 19.6)	11.1	(10.7 - 11.5)	+		
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	20.5	(17.1 - 23.8)	15.4	(15.0 - 15.9)	+		
Sufficient vigorous activity	62.3	(58.2 - 66.4)	66.5	(65.9 - 67.1)			+
Sufficient moderate activity	22.7	(19.0 - 26.4)	25.3	(24.8 - 25.8)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	44.3	(40.6 - 47.9)	43.5	(42.9 - 44.1)			+
Described themselves as slightly or very overweight	35.9	(31.7 - 40.1)	30.0	(29.5 - 30.6)	+		
Never or rarely wore a seat belt when riding in a car driven by someone else	18.9	(15.2 - 22.6)	10.9	(10.4 - 11.3)	+		
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	24.8	(21.0 - 28.7)	20.2	(19.7 - 20.7)	+		
Did not go to school on one or more of the past 30 days because they felt unsafe	12.4	(9.6 - 15.3)	6.9	(6.6 - 7.2)	+		
Purposely hurt themselves without wanting to die, during the past 12 months	15.0	(12.0 - 17.9)	12.4	(12.0 - 12.8)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	23.9	(20.0 - 27.8)	22.0	(21.5 - 22.5)			+