

Suwannee County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Suwannee County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	5.4	(3.0 - 7.9)	3.3	(3.0 - 3.5)			+
Smoked cigars on one or more of the past 30 days	7.0	(3.3 - 10.8)	3.9	(3.6 - 4.2)			+
Used smokeless tobacco on one or more of the past 30 days	9.6	(5.1 - 14.2)	2.3	(2.1 - 2.6)	+		
Used any form of tobacco on one or more of the past 30 days	13.7	(8.8 - 18.5)	6.4	(6.0 - 6.8)	+		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	66.1	(60.0 - 72.1)	72.0	(71.2 - 72.8)			+
Exposed to secondhand smoke during the past 7 days	49.8	(44.9 - 54.8)	41.3	(40.4 - 42.2)	+		
Smoking is allowed in the home	12.7	(9.1 - 16.3)	9.9	(9.5 - 10.4)			+
Taught about tobacco use in any classes this year	29.8	(23.0 - 36.6)	53.4	(52.0 - 54.7)		+	
Parent or guardian has talked about the dangers of tobacco in the past year	57.7	(51.3 - 64.1)	63.9	(63.1 - 64.6)			+
Served as an active member of SWAT	8.8	(6.3 - 11.4)	4.3	(3.9 - 4.7)	+		
<u>Definitely did not think</u> that smokers have more friends	34.4	(28.4 - 40.3)	33.7	(32.9 - 34.5)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	73.0	(67.2 - 78.9)	73.7	(73.0 - 74.4)			+

Suwannee County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Middle School				Comparison of Two Rates		
	Suwannee County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	19.4	(16.0 - 22.7)	20.1	(19.5 - 20.7)			+
Current asthma, among all individuals	12.4	(9.7 - 15.1)	12.0	(11.5 - 12.5)			+
Asthma attack in past year, among individuals with lifetime asthma	22.8	(13.0 - 32.6)	18.1	(16.8 - 19.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	14.9	(10.1 - 19.6)	11.6	(11.0 - 12.1)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	18.9	(14.8 - 23.0)	16.5	(15.8 - 17.2)			+
Sufficient vigorous activity	68.6	(63.4 - 73.9)	70.1	(69.2 - 70.9)			+
Sufficient moderate activity	26.7	(21.0 - 32.5)	24.8	(24.1 - 25.4)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	44.1	(38.2 - 49.9)	44.4	(43.6 - 45.2)			+
Described themselves as slightly or very overweight	29.3	(24.3 - 34.3)	30.2	(29.4 - 30.9)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	14.2	(10.2 - 18.2)	11.4	(10.8 - 12.0)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	24.1	(18.6 - 29.7)	17.8	(17.2 - 18.5)	+		
Did not go to school on one or more of the past 30 days because they felt unsafe	7.5	(5.1 - 10.0)	7.8	(7.3 - 8.3)			+
Purposely hurt themselves without wanting to die, during the past 12 months	10.5	(6.7 - 14.4)	12.6	(12.0 - 13.2)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	16.2	(12.8 - 19.6)	19.2	(18.5 - 19.8)			+

Suwannee County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Suwannee County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	15.5	(10.8 - 20.3)	10.1	(9.6 - 10.6)	+		
Smoked cigars on one or more of the past 30 days	15.5	(11.2 - 19.7)	11.4	(10.9 - 11.9)			+
Used smokeless tobacco on one or more of the past 30 days	11.7	(7.3 - 16.1)	5.6	(5.2 - 6.0)	+		
Used any form of tobacco on one or more of the past 30 days	26.3	(19.9 - 32.7)	17.9	(17.3 - 18.6)	+		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	49.8	(43.8 - 55.8)	58.4	(57.6 - 59.3)		+	
Exposed to secondhand smoke during the past 7 days	58.2	(53.3 - 63.0)	47.6	(46.7 - 48.4)	+		
Smoking is allowed in the home	17.0	(12.0 - 21.9)	10.0	(9.5 - 10.5)	+		
Taught about tobacco use in any classes this year	31.7	(25.4 - 37.9)	33.4	(32.4 - 34.4)			+
Parent or guardian has talked about the dangers of tobacco in the past year	45.8	(38.7 - 52.8)	50.4	(49.6 - 51.2)			+
Served as an active member of SWAT	8.0	(4.8 - 11.2)	4.0	(3.7 - 4.3)	+		
<u>Definitely did not think</u> that smokers have more friends	32.7	(25.6 - 39.9)	30.5	(29.8 - 31.2)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	70.4	(65.0 - 75.7)	71.7	(71.0 - 72.4)			+

Suwannee County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	High School				Comparison of Two Rates		
	Suwannee County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	21.1	(16.4 - 25.8)	20.8	(20.2 - 21.5)			+
Current asthma, among all individuals	10.9	(7.2 - 14.5)	10.6	(10.1 - 11.1)			+
Asthma attack in past year, among individuals with lifetime asthma	10.2	(1.6 - 18.8)	16.3	(15.1 - 17.4)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	13.4	(9.6 - 17.2)	11.1	(10.5 - 11.6)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	16.1	(10.6 - 21.6)	14.3	(13.7 - 14.9)			+
Sufficient vigorous activity	59.5	(54.1 - 64.8)	62.7	(62.0 - 63.5)			+
Sufficient moderate activity	23.1	(18.2 - 28.1)	26.1	(25.4 - 26.7)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	46.1	(40.7 - 51.5)	42.3	(41.5 - 43.1)			+
Described themselves as slightly or very overweight	35.2	(30.2 - 40.2)	29.8	(29.0 - 30.5)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	14.9	(10.7 - 19.2)	11.2	(10.5 - 11.8)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	25.1	(19.7 - 30.5)	23.1	(22.4 - 23.8)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	5.4	(3.1 - 7.7)	6.9	(6.4 - 7.3)			+
Purposely hurt themselves without wanting to die, during the past 12 months	13.4	(9.1 - 17.6)	12.7	(12.1 - 13.2)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	24.5	(19.8 - 29.2)	24.5	(23.8 - 25.2)			+

Suwannee County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Suwannee County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Smoked cigarettes on one or more of the past 30 days	8.3	(5.9 - 10.8)	6.1	(5.8 - 6.4)			+
Smoked cigars on one or more of the past 30 days	8.7	(6.3 - 11.0)	6.9	(6.6 - 7.2)			+
Used smokeless tobacco on one or more of the past 30 days	9.2	(6.3 - 12.1)	3.5	(3.2 - 3.7)	+		
Used any form of tobacco on one or more of the past 30 days	16.7	(13.0 - 20.5)	11.2	(10.8 - 11.7)	+		
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	60.1	(55.5 - 64.6)	65.7	(65.1 - 66.3)		+	
Exposed to secondhand smoke during the past 7 days	53.1	(49.3 - 56.8)	44.1	(43.4 - 44.7)	+		
Smoking is allowed in the home	14.7	(11.7 - 17.6)	9.4	(9.0 - 9.8)	+		
Taught about tobacco use in any classes this year	32.1	(27.1 - 37.1)	43.2	(42.3 - 44.1)		+	
Parent or guardian has talked about the dangers of tobacco in the past year	53.3	(48.0 - 58.6)	57.4	(56.8 - 58.0)			+
Served as an active member of SWAT	8.0	(5.8 - 10.1)	3.9	(3.7 - 4.1)	+		
<u>Definitely did not think</u> that smokers have more friends	34.0	(29.4 - 38.6)	32.0	(31.5 - 32.6)			+
<u>Definitely did not think</u> that smoking helps young people "look cool" or "fit in"	73.1	(69.2 - 76.9)	72.9	(72.4 - 73.4)			+

Suwannee County: 2012 At a Glance

Data from the 2012 Florida Youth Tobacco Survey (FYTS)

Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Suwannee County %	95% CI	State %	95% CI	County Higher Than State	State Higher Than County	No Difference
Lifetime asthma	19.5	(16.6 - 22.4)	20.5	(20.1 - 21.0)			+
Current asthma, among all individuals	11.5	(9.2 - 13.8)	11.3	(11.0 - 11.7)			+
Asthma attack in past year, among individuals with lifetime asthma	16.6	(9.6 - 23.6)	16.9	(16.0 - 17.8)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	13.3	(10.3 - 16.2)	11.1	(10.7 - 11.5)			+
Overweight (i.e., at or above the 85 th percentile but below the 95th percentile for body mass index, by age and sex)	18.1	(14.5 - 21.8)	15.4	(15.0 - 15.9)			+
Sufficient vigorous activity	65.9	(61.9 - 69.9)	66.5	(65.9 - 67.1)			+
Sufficient moderate activity	25.6	(21.6 - 29.7)	25.3	(24.8 - 25.8)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	46.1	(41.9 - 50.2)	43.5	(42.9 - 44.1)			+
Described themselves as slightly or very overweight	32.0	(28.3 - 35.7)	30.0	(29.5 - 30.6)			+
Never or rarely wore a seat belt when riding in a car driven by someone else	13.3	(10.4 - 16.1)	10.9	(10.4 - 11.3)			+
Rode in a car driven by someone who had been drinking alcohol during the past 30 days	24.3	(20.3 - 28.4)	20.2	(19.7 - 20.7)			+
Did not go to school on one or more of the past 30 days because they felt unsafe	5.8	(4.1 - 7.5)	6.9	(6.6 - 7.2)			+
Purposely hurt themselves without wanting to die, during the past 12 months	11.3	(8.3 - 14.4)	12.4	(12.0 - 12.8)			+
Felt so sad or hopeless almost every day for two weeks or more in a row that they stopped doing some usual activities during the past 12 months	18.9	(15.9 - 21.9)	22.0	(21.5 - 22.5)			+