



# Brevard County: 2014 at a Glance

Data from the 2014 Florida Youth Tobacco Survey (FYTS)



Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Brevard County %	95% CI	State %	95% CI	County Significantly Higher Than State	State Significantly Higher Than County	No Significant Difference
Smoked cigarettes on one or more of the past 30 days	<b>4.9</b>	(3.5-6.3)	<b>4.3</b>	(4.1-4.6)			+
Smoked cigars on one or more of the past 30 days	<b>5.5</b>	(3.9-7.1)	<b>5.4</b>	(5.2-5.7)			+
Used smokeless tobacco on one or more of the past 30 days	<b>3.3</b>	(2.2-4.4)	<b>3.3</b>	(3.1-3.5)			+
Used electronic cigarettes on one or more of the past 30 days	<b>9.1</b>	(7.3-11.0)	<b>7.2</b>	(6.8-7.5)			+
Used hookah on one or more of the past 30 days	<b>6.5</b>	(5.0-8.0)	<b>7.1</b>	(6.7-7.5)			+
Used any form of tobacco on one or more of the past 30 days*	<b>9.2</b>	(7.1-11.3)	<b>9.2</b>	(8.8-9.6)			+
Used any form of tobacco (excluding cigarettes) on one or more of the past 30 days	<b>15.1</b>	(12.6-17.6)	<b>14.8</b>	(14.2-15.3)			+
Thought using electronic cigarettes is less harmful than smoking cigarettes	<b>75.4</b>	(71.7-79.1)	<b>69.4</b>	(68.4-70.3)	+		
Thought using hookah is less harmful than smoking cigarettes	<b>53.3</b>	(47.4-59.2)	<b>52.8</b>	(51.7-54.0)			+
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	<b>67.7</b>	(64.6-70.8)	<b>67.1</b>	(66.4-67.7)			+
Exposed to secondhand smoke during the past seven days (in a room or car)	<b>43.0</b>	(39.6-46.4)	<b>37.5</b>	(36.9-38.2)	+		
Smoking is allowed in the home	<b>10.5</b>	(8.6-12.4)	<b>8.4</b>	(8.0-8.7)			+

\*This excludes specialty tobacco products, which include bidis, kreteks, pipe tobacco, snus, e-cigarettes, flavored cigarettes, flavored cigars, and flavored smokeless tobacco.



# Brevard County: 2014 at a Glance

Data from the 2014 Florida Youth Tobacco Survey (FYTS)



Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Brevard County %	95% CI	State %	95% CI	County Significantly Higher Than State	State Significantly Higher Than County	No Significant Difference
Students who were taught about tobacco use in any classes this year	<b>37.7</b>	(34.0-41.5)	<b>38.0</b>	(37.1-39.0)			+
Parent or guardian has talked about the dangers of tobacco in the past year	<b>67.0</b>	(64.2-69.7)	<b>62.4</b>	(61.8-63.0)	+		
Served as an active member of Students Working Against Tobacco (SWAT)	<b>2.0</b>	(1.2-2.8)	<b>2.8</b>	(2.7-3.0)			+
Definitely did not think that smokers have more friends	<b>25.1</b>	(22.7-27.5)	<b>28.2</b>	(27.6-28.8)		+	
Definitely did not think that smoking helps young people "look cool" or "fit in"	<b>69.7</b>	(66.9-72.6)	<b>68.8</b>	(68.2-69.3)			+
Lifetime asthma	<b>19.6</b>	(17.5-21.7)	<b>20.8</b>	(20.3-21.3)			+
Current asthma	<b>11.2</b>	(9.5-12.9)	<b>11.5</b>	(11.1-11.9)			+
Asthma attack in past year	<b>12.6</b>	(8.7-16.6)	<b>17.7</b>	(16.7-18.7)		+	
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	<b>10.9</b>	(9.2-12.6)	<b>12.2</b>	(11.8-12.7)			+
Overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex)	<b>16.5</b>	(14.2-18.7)	<b>15.9</b>	(15.4-16.4)			+
Physically active for at least 60 minutes per day during the past seven days	<b>24.1</b>	(21.8-26.4)	<b>22.7</b>	(22.2-23.3)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	<b>43.7</b>	(41.1-46.3)	<b>43.7</b>	(43.1-44.4)			+
Described themselves as slightly or very overweight	<b>31.4</b>	(28.8-34.1)	<b>30.7</b>	(30.2-31.3)			+