



# Dixie County: 2014 at a Glance

Data from the 2014 Florida Youth Tobacco Survey (FYTS)



Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Dixie County %	95% CI	State %	95% CI	County Significantly Higher Than State	State Significantly Higher Than County	No Significant Difference
Smoked cigarettes on one or more of the past 30 days	<b>10.3</b>	(6.5-14.1)	<b>4.3</b>	(4.1-4.6)	+		
Smoked cigars on one or more of the past 30 days	<b>8.1</b>	(5.1-11.2)	<b>5.4</b>	(5.2-5.7)			+
Used smokeless tobacco on one or more of the past 30 days	<b>16.0</b>	(10.7-21.3)	<b>3.3</b>	(3.1-3.5)	+		
Used electronic cigarettes on one or more of the past 30 days	<b>5.2</b>	(2.6-7.8)	<b>7.2</b>	(6.8-7.5)			+
Used hookah on one or more of the past 30 days	<b>3.8</b>	(1.8-5.8)	<b>7.1</b>	(6.7-7.5)		+	
Used any form of tobacco on one or more of the past 30 days*	<b>24.5</b>	(18.8-30.3)	<b>9.2</b>	(8.8-9.6)	+		
Used any form of tobacco (excluding cigarettes) on one or more of the past 30 days	<b>25.1</b>	(19.2-31.1)	<b>14.8</b>	(14.2-15.3)	+		
Thought using electronic cigarettes is less harmful than smoking cigarettes	<b>62.7</b>	(53.6-71.8)	<b>69.4</b>	(68.4-70.3)			+
Thought using hookah is less harmful than smoking cigarettes	<b>14.3</b>	(6.1-22.4)	<b>52.8</b>	(51.7-54.0)		+	
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	<b>52.4</b>	(46.1-58.6)	<b>67.1</b>	(66.4-67.7)		+	
Exposed to secondhand smoke during the past seven days (in a room or car)	<b>52.4</b>	(47.7-57.1)	<b>37.5</b>	(36.9-38.2)	+		
Smoking is allowed in the home	<b>19.0</b>	(14.7-23.3)	<b>8.4</b>	(8.0-8.7)	+		

\*This excludes specialty tobacco products, which include bidis, kreteks, pipe tobacco, snus, e-cigarettes, flavored cigarettes, flavored cigars, and flavored smokeless tobacco.



# Dixie County: 2014 at a Glance

Data from the 2014 Florida Youth Tobacco Survey (FYTS)



Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Dixie County %	95% CI	State %	95% CI	County Significantly Higher Than State	State Significantly Higher Than County	No Significant Difference
Students who were taught about tobacco use in any classes this year	<b>53.7</b>	(44.0-63.5)	<b>38.0</b>	(37.1-39.0)	+		
Parent or guardian has talked about the dangers of tobacco in the past year	<b>60.9</b>	(55.8-65.9)	<b>62.4</b>	(61.8-63.0)			+
Served as an active member of Students Working Against Tobacco (SWAT)	<b>22.1</b>	(17.8-26.5)	<b>2.8</b>	(2.7- 3.0)	+		
Definitely did not think that smokers have more friends	<b>25.2</b>	(20.7-29.7)	<b>28.2</b>	(27.6-28.8)			+
Definitely did not think that smoking helps young people "look cool" or "fit in"	<b>68.6</b>	(63.8-73.4)	<b>68.8</b>	(68.2-69.3)			+
Lifetime asthma	<b>26.7</b>	(22.8-30.6)	<b>20.8</b>	(20.3-21.3)	+		
Current asthma	<b>15.4</b>	(11.7-19.1)	<b>11.5</b>	(11.1-11.9)			+
Asthma attack in past year	<b>22.3</b>	(13.4-31.3)	<b>17.7</b>	(16.7-18.7)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	<b>17.8</b>	(12.3-23.3)	<b>12.2</b>	(11.8-12.7)			+
Overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex)	<b>17.5</b>	(13.5-21.5)	<b>15.9</b>	(15.4-16.4)			+
Physically active for at least 60 minutes per day during the past seven days	<b>32.8</b>	(28.2-37.5)	<b>22.7</b>	(22.2-23.3)	+		
Exercised to lose weight or to keep from gaining weight during the past 30 days	<b>51.2</b>	(44.9-57.5)	<b>43.7</b>	(43.1-44.4)	+		
Described themselves as slightly or very overweight	<b>38.5</b>	(33.3-43.7)	<b>30.7</b>	(30.2-31.3)	+		