



# Lake County: 2014 at a Glance

Data from the 2014 Florida Youth Tobacco Survey (FYTS)



Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Lake County %	95% CI	State %	95% CI	County Significantly Higher Than State	State Significantly Higher Than County	No Significant Difference
Smoked cigarettes on one or more of the past 30 days	5.6	(4.1-7.1)	4.3	(4.1-4.6)			+
Smoked cigars on one or more of the past 30 days	6.1	(4.6-7.6)	5.4	(5.2-5.7)			+
Used smokeless tobacco on one or more of the past 30 days	4.2	(2.9-5.4)	3.3	(3.1-3.5)			+
Used electronic cigarettes on one or more of the past 30 days	7.2	(5.4-8.9)	7.2	(6.8-7.5)			+
Used hookah on one or more of the past 30 days	5.8	(4.5-7.2)	7.1	(6.7-7.5)			+
Used any form of tobacco on one or more of the past 30 days*	11.3	(9.1-13.6)	9.2	(8.8-9.6)			+
Used any form of tobacco (excluding cigarettes) on one or more of the past 30 days	15.3	(12.8-17.9)	14.8	(14.2-15.3)			+
Thought using electronic cigarettes is less harmful than smoking cigarettes	70.0	(65.7-74.4)	69.4	(68.4-70.3)			+
Thought using hookah is less harmful than smoking cigarettes	46.4	(39.9-52.9)	52.8	(51.7-54.0)			+
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	66.6	(63.3-69.9)	67.1	(66.4-67.7)			+
Exposed to secondhand smoke during the past seven days (in a room or car)	42.2	(38.5-45.9)	37.5	(36.9-38.2)	+		
Smoking is allowed in the home	9.9	(7.8-11.9)	8.4	(8.0-8.7)			+

\*This excludes specialty tobacco products, which include bidis, kreteks, pipe tobacco, snus, e-cigarettes, flavored cigarettes, flavored cigars, and flavored smokeless tobacco.



# Lake County: 2014 at a Glance

Data from the 2014 Florida Youth Tobacco Survey (FYTS)



Indicator	Youth (Ages 11-17)				Comparison of Two Rates		
	Lake County %	95% CI	State %	95% CI	County Significantly Higher Than State	State Significantly Higher Than County	No Significant Difference
Students who were taught about tobacco use in any classes this year	<b>39.6</b>	(34.2-44.9)	<b>38.0</b>	(37.1-39.0)			+
Parent or guardian has talked about the dangers of tobacco in the past year	<b>58.6</b>	(55.1-62.0)	<b>62.4</b>	(61.8-63.0)			+
Served as an active member of Students Working Against Tobacco (SWAT)	<b>1.8</b>	(0.9-2.8)	<b>2.8</b>	(2.7- 3.0)			+
Definitely did not think that smokers have more friends	<b>28.3</b>	(25.6-31.0)	<b>28.2</b>	(27.6-28.8)			+
Definitely did not think that smoking helps young people "look cool" or "fit in"	<b>72.2</b>	(69.1-75.3)	<b>68.8</b>	(68.2-69.3)			+
Lifetime asthma	<b>19.8</b>	(17.6-22.1)	<b>20.8</b>	(20.3-21.3)			+
Current asthma	<b>11.6</b>	(9.4-13.8)	<b>11.5</b>	(11.1-11.9)			+
Asthma attack in past year	<b>20.9</b>	(15.1-26.6)	<b>17.7</b>	(16.7-18.7)			+
Obese (i.e., at or above the 95th percentile for body mass index, by age and sex)	<b>14.0</b>	(11.6-16.3)	<b>12.2</b>	(11.8-12.7)			+
Overweight (i.e., at or above the 85th percentile but below the 95th percentile for body mass index, by age and sex)	<b>14.0</b>	(11.9-16.1)	<b>15.9</b>	(15.4-16.4)			+
Physically active for at least 60 minutes per day during the past seven days	<b>23.7</b>	(21.1-26.3)	<b>22.7</b>	(22.2-23.3)			+
Exercised to lose weight or to keep from gaining weight during the past 30 days	<b>44.9</b>	(42.0-47.8)	<b>43.7</b>	(43.1-44.4)			+
Described themselves as slightly or very overweight	<b>32.1</b>	(28.9-35.4)	<b>30.7</b>	(30.2-31.3)			+