



Weeks 21-22: May 22-June 4, 2011



Summary

The Florida Department of Health (FDOH) monitors multiple surveillance systems such as the Electronic Surveillance System for the Early Notification of Community-based Epidemics (ESSENCE), the Florida Pneumonia and Influenza Mortality Surveillance System (FPIMSS), notifiable disease reports (Merlin), National Respiratory and Enteric Virus Surveillance System (NREVSS), EpiCom, and Florida ILINet in order to track influenza activity in the state. **This is the first abbreviated summer version of the Florida Flu Review for summer 2011. The full report will return in week 40.**

- State:**
- ILI activity is low in many of our monitoring systems. Mandatory submissions of county influenza activity levels has ended for the summer months, so no county level influenza activity data is presented here.
 - ESSENCE and ILINet ILI activity levels are near previous non-pandemic seasons at this time in most regions of Florida. Due to decreased reporting in the summer months, regional data from ILINet sentinels will not be reported until the beginning of the 2011-12 season.
 - In week 22 there were no positive flu results from the state laboratory, and in week 21 there was one positive lab result for 2009 H1N1 influenza A. Other viruses known to be currently circulating, potentially causing ILI, include adenovirus, rhinovirus, parainfluenza, and respiratory syncytial virus (RSV).
 - No outbreaks of ILI or influenza were reported in weeks 21-22.

June 8, 2011

Posted on the Bureau of Epidemiology website: http://www.doh.state.fl.us/Disease_ctrl/epi/htopics/flu/reports.htm

Produced by: Bureau of Epidemiology, Florida Department of Health (FDOH)

Contributors: Colin Malone, MPH; Brian Lang; Aaron Kite-Powell, MS; Brian Fox, MA; Kate Goodin, MPH; Leah Eisenstein, MPH; Lillian Stark, PhD, MPH, MS; Valerie Mock; Julian Everett; Janet Hamilton, MPH; Richard Hopkins, MD, MSPH

TABLE 1: Summary of Florida Influenza-Like Illness (ILI) Activity for Week 22

Measure	Difference from previous week	Current week 22	Previous week 21
Percent of visits to ILINet providers for ILI	▼ 0.1	0.5%	0.6%
Percent of emergency department visits (from ESSENCE) due to ILI	▼ 0.1	1.5%	1.6%
Percent of hospital admissions (from ESSENCE) due to ILI	No Change	0.3%	0.3%
Number of laboratory specimens that were positive for influenza (out of total number tested)	▼ 1	0 (of 9)	1 (of 11)

In this Issue:

Summary	1
NREVSS Respiratory Virus Surveillance	1
Outpatient Influenza-like Illness Surveillance Network (ILINET)-Statewide	2
ESSENCE Syndromic Surveillance Summary-Statewide	2
Florida Pneumonia and Influenza Mortality Surveillance (FPIMSS)	2

NREVSS Respiratory Virus Surveillance


The National Respiratory and Enteric Virus Surveillance System (NREVSS) collects data from laboratory facilities around the country on a weekly basis. NREVSS monitors temporal and geographic patterns of respiratory syncytial virus (RSV), human parainfluenza viruses (HPIV), respiratory and enteric adenoviruses and rotavirus. Florida has over 30 participating laboratory facilities.

FIGURE 1 shows the percentage of positive tests for multiple respiratory viruses reported by NREVSS-participating laboratories in Florida

The six respiratory viruses summarized in Figure 1 are:

- RSV
- Parainfluenza 1-3 (HPIV)
- Adenovirus
- Human Metapneumo Virus
- Rhinovirus
- Influenza

Figure 1: Percentage of Positive Respiratory Virus Tests as Collected by NREVSS, Florida, 2009-2011, as of June 8, 2011




— %RSV — %Parainfluenza — %Adenovirus — % Human Metapneumo Virus — % Rhinovirus — % Influenza

FIGURE 2 shows the percentage of visits for influenza-like illness* reported by ILINet Sentinel Providers statewide.

The percentage of visits for ILINet sentinel providers continues to be low.

Note: Small numbers of sentinels reporting year-round may make summer ILI estimates less reliable.




*ILI = Influenza-like illness, fever >100°F AND sore throat and/or cough *in the absence* of another known cause.
 **The 2010—2011 threshold for moderate activity is calculated from ILINet data. The threshold for moderate activity is the mean percentage of patient visits for ILI during influenza weeks for the previous three seasons plus two standard deviations. Only weeks with 10% or greater of laboratory specimens testing positive are included in the calculation. Due to wide variability in regional level data, it is not appropriate to apply the state baseline to regional data.
 ^There is no week 53 during the 2007-2008, 2009-2010, and 2010-2011 seasons; the week 53 data point for those seasons is an average of weeks 52 and 1.

ESSENCE Syndromic Surveillance Summary-Statewide

The Electronic Surveillance System for the Early Notification of Community-based Epidemics (ESSENCE) currently collects data daily from 163 hospital emergency departments (ED). Data are processed into 11 different syndrome categories based on the patient's chief complaint. One category is influenza-like illness (ILI), which comprises chief complaints that include "influenza" or "flu," or either fever and cough or sore throat.

FIGURE 3 shows ESSENCE data on ILI visits to Emergency Departments as a percentage of all ED Visits.

Overall activity for influenza-like illness reported in ESSENCE is near expected levels for this time of year.



Florida Pneumonia and Influenza Mortality Surveillance

The Florida Department of Health started the Florida Pneumonia and Influenza Mortality Surveillance System (FPIMSS) in 2006 in order to more timely assess the number of pneumonia and influenza deaths occurring in the state. This system was modeled on the CDC's 122 cities surveillance system. Each week, the vital statistics office in the 24 most populous counties in Florida manually reviews the death certificates received for the previous week. Any mention of pneumonia or influenza on the death certificate, with certain prescribed exceptions, is counted as a pneumonia or influenza death. These counts, by age group, are then reported to the state via the EpiGateway web-interface. We are now using a Serfling model to more accurately calculate our predicted values for weekly pneumonia and influenza mortality.

FIGURE 4 shows Pneumonia and Influenza Deaths for 24 Florida Counties, 2007-2008, 2008-2009, 2009-2010, and 2010-11

For week 22 (ending June 4, 2011) there were:

- 109 deaths reported
- Upper bound of 95% confidence interval for prediction: 177 deaths
- NO excess deaths

23 out of 24 participating counties reported their data for week 22.

