



## Week 2: January 11 - 17, 2015



### Summary

#### National influenza activity:

**Influenza activity is elevated nationally.**

- The Centers for Disease Control and Prevention (CDC) has identified an antigenically drifted influenza A (H3N2) strain circulating nationally and in Florida that is different from the strain of influenza A (H3N2) contained in the current 2014-15 influenza vaccine formulations.
- The CDC indicates this season's vaccine is offering reduced protection, as such, use of neuraminidase inhibitor antiviral medications, for treatment and prevention of influenza, is more important than ever. High risk individuals with suspected flu should be treated with antivirals as early as possible (even prior to laboratory confirmation). More information can be found here: <http://www.floridahealth.gov/diseases-and-conditions/influenza/documents/Other/influenza-letter-for-health-care-providers.pdf>.
  - The CDC indicates that antiviral medications are underutilized; one study estimates antivirals were only used one out of five times where antivirals use would be recommended.

#### State Influenza and influenza-like illness (ILI) activity:

- **Flu activity remains high in Florida and is widespread.**
  - The 2014-15 flu season began early and is in full swing in Florida.
- Although influenza activity has decreased in recent weeks in some surveillance systems, overall activity levels remain high and it is too early to tell if the season has peaked.
- **Seasons like this one, where influenza A (H3) is the predominantly circulating strain, are typically associated with higher morbidity and mortality, particularly in the 65+ age group.**
  - More hospitalizations and deaths are typical of H3N2-seasons, which hit young children and older people harder.
- **While biggest increases in ED visits for ILI have most recently been identified in the 65+ age group, activity is still greatest in children.**
  - Forty-seven (66%) of reported outbreaks of ILI have been in facilities that primarily serve the 65+ years old age group.
- **In the past week, the number of pneumonia and influenza associated deaths, particularly in those over the age of 65 have increased to above levels seen during previous years at this time.** Increases in hospitalizations and deaths at this point in the season are expected during severe flu years, like this one.
  - During flu season, increases in ED visits typically come before increases hospitalizations and deaths.
- It is likely that flu deaths will reach higher levels later this season since mortality tends to lag behind other indicators.
- **In Florida, the most common influenza subtype detected at the Bureau of Public Health Laboratories (BPHL) in recent weeks has been influenza A (H3).**
- In the past week, 24 of 37 (64.9%) specimens submitted for influenza testing at BPHL were PCR positive for seasonal strains of influenza: 13 were positive for influenza A (H3), nine were influenza A not yet subtyped, and two influenza B not yet subtyped.
- **Eight outbreaks of influenza (two or more cases of influenza or ILI in a specific setting) were reported to EpiCom in week 2.**
- **No pediatric influenza-associated deaths were reported in week 2.**


January 21, 2015

Posted on the Bureau of Epidemiology website:


<http://www.floridahealth.gov/floridaflu>

Produced by: Bureau of Epidemiology, Florida Department of Health

Contributors: Heather Rubino, PhD; Ellen Dugan, MPH, Leah Eisenstein, MPH; Lea Heberlein-Larson, MPH; Valerie Mock; Janet Hamilton, MPH




**Map 1: County Influenza Activity Week 2, 2015**



Fifteen counties reported moderate influenza activity. For more information, see page 6.

**Map 2: Influenza and ILI Outbreaks Week 2, 2015**



**Seventy-one** outbreaks of ILI or influenza have been reported since Week 40, 2014. For more information, see page 10.

#### In this Issue:

Summary	1
ILINet ILI-Statewide	2
ESSENCE-FL Syndromic Surveillance Summary-Statewide	3
ESSENCE-FL Syndromic Surveillance Summary-Regional	3
Bureau of Public Health Laboratories Viral Surveillance	5
County Influenza and ILI Activity	6
Pediatric Influenza-Associated Mortality	8
ESSENCE-FL Pneumonia and Influenza Mortality	9
NREVSS Respiratory Virus Surveillance	10
Influenza and ILI Outbreaks Reported in EpiCom	10
Florida ILI Surveillance System Summary	11

Descriptions of Florida influenza and ILI surveillance systems can be found on page 11.

TABLE 1: Summary of Florida Influenza-Like Illness (ILI) Activity for Week 2

Measure	Difference from Previous Week	Current Week 2	Previous Week 1	Page of Report
Overall statewide activity code reported to CDC	No Change	Widespread	Widespread	1
Percent of visits to ILINet providers for ILI	▲ 0.2	2.5%	2.3%	2
Percent of ED and UCC visits (from ESSENCE-FL) due to ILI	▼ 0.7	3.3%	4.0%	3
Percent of laboratory specimens that were positive for influenza	▲ 14.9	64.9%	50.0%	5
Number of counties reporting moderate influenza activity	No Change	15	15	6
Number of counties reporting widespread influenza activity	▲ 1	1	0	6
Number of counties reporting increasing influenza activity	No Change	13	13	6
Number of counties reporting decreasing influenza activity	▲ 2	21	19	6
Number of ILI outbreaks reported in EpiCom	▼ 8	9	17	10


## ILINet Influenza-Like Illness-Statewide

ILINet is a nationwide surveillance system composed of sentinel providers. Florida has 107 providers enrolled in ILINet who submit weekly ILI and total visit counts, as well as submit ILI specimens to the BPHL for confirmatory testing.

**FIGURE 1** shows the percentage of visits for ILI\* reported by ILINet sentinel providers statewide.

The percent of visits to ILINet sentinel outpatient physicians for ILI is at or near levels seen in previous years at this time.


FIGURE 1: Percentage of Visits for Influenza-Like Illness (ILI)\* Reported by ILINet Sentinel Providers Statewide, Week 40, 2011 through Week 2, 2015 as Reported by 5:00 p.m. January 20, 2015



**FIGURE 2** shows ILI visit counts reported by ILINet sentinel providers statewide by age group.

In week 2, the number of ILI visits to ILINet sentinels decreased in the all age groups in concordance with the decline seen in previous weeks.

FIGURE 2: Influenza-Like Illness (ILI)\* Visit Counts Reported by ILINet Sentinel Providers Statewide by Age Group Week 40, 2011 through Week 2, 2015 as Reported by 5:00 p.m. January 20, 2015



[Return to Top](#)

†Data presented here are counts, not proportions as included in Figure 1. This is because age group denominator data is not available through ILINet.

\*ILI = Influenza-like illness, fever >100°F AND sore throat and/or cough *in the absence* of another known cause.


ESSENCE-FL collects data daily from 219 EDs and UCCs. These data are processed into 11 different syndrome categories based on the patient's chief complaint. One of the categories is ILI, which is composed of chief complaints that include the words "influenza" or "flu," or complaints that contain fever plus cough or sore throat.

**FIGURE 3** shows ESSENCE-FL data on ILI visits to EDs and UCCs as a percentage of all visits.

The percent of visits to EDs and UCCs for ILI has been above levels seen in previous years at this time, but has declined in recent weeks.

There continues to be increased levels of influenza in pregnant women presenting to EDs for care that is above expected levels for this time of year. Pregnant women are among those at high risk for severe complications due to influenza infection. More information can be found here: [http://www.floridahealth.gov/diseases-and-conditions/influenza/\\_documents/Other/influenza-guidance-for-health-care-providers.pdf](http://www.floridahealth.gov/diseases-and-conditions/influenza/_documents/Other/influenza-guidance-for-health-care-providers.pdf)


FIGURE 3: Percentage of Influenza Like-Illness visits from Emergency Department (ED) and Urgent Care Center (UCC) Chief Complaints, ESSENCE-FL Participating Facilities (N=219), Week 40, 2011 through Week 2, 2015 accessed January 21, 2015



**FIGURE 4** shows percentage of ILI among all ED and UCC visits by age.


The percent of ED and UCC visits for ILI is similar to levels seen in previous years in all age groups at this time. While the proportion of ED and UCC visits for ILI has decreased in almost all age groups in the past few weeks, activity still remains highest in children.

FIGURE 4: Percentage of Influenza Like-Illness visits from Emergency Department (ED) and Urgent Care Center (UCC) Chief Complaints by Age, ESSENCE-FL Participating Facilities (N=219), Week 40, 2011 through Week 2, 2015 accessed January 21, 2015




## ESSENCE-FL Syndromic Surveillance-Regional

Map 3: Emergency Departments and Urgent Care Centers Reporting Data to ESSENCE-FL by Regional Domestic Security Task Force (RDSTF), January 21, 2015 (N=219)



**FIGURE 5 - FIGURE 11** describe Emergency Department (ED) and Urgent Care Center (UCC) chief complaint data from ESSENCE-FL by Regional Domestic Security Task Force (RDSTF) regions.

FIGURE 5: Percentage of of Influenza Like-Illness Visits from ED Chief Complaints, RDSTF Region 1 ESSENCE-FL Participating Hospitals (N=14), Week 40, 2011 through Week 2, 2015 accessed January 21, 2015



After having been elevated, ED and UCC visits for ILI in RDSTF Regions 1-7 are at or near levels seen during previous years at this time.

FIGURE 6: Percentage of of Influenza Like-Illness Visits from ED and UCC Chief Complaints, RDSTF Region 2 ESSENCE-FL Participating Facilities (N=7), Week 40, 2011 through Week 2, 2015 accessed January 21, 2015




FIGURE 7: Percentage of of Influenza Like-Illness Visits from ED Chief Complaints, RDSTF Region 3 ESSENCE-FL Participating Facilities (N=20), Week 40, 2011 through Week 2, 2015 accessed January 21, 2015




FIGURE 8: Percentage of of Influenza Like-Illness Visits from ED and UCC Chief Complaints, RDSTF Region 4 ESSENCE-FL Participating Facilities (N=42), Week 40, 2011 through Week 2, 2015 accessed January 21, 2015




FIGURE 9: Percentage of ILI visits from ED and UCC Chief Complaints, ESSENCE-FL Participating Facilities RDSTF Region 5 ESSENCE-FL Participating Facilities (N=61), Week 40, 2011 through Week 2, 2015 accessed January 21, 2015




FIGURE 10: Percentage of of Influenza Like-Illness Visits from ED and UCC Chief Complaints, RDSTF Region 6 ESSENCE-FL Participating Facilities (N=23), Week 40, 2011 through Week 2, 2015 accessed January 21, 2015





FIGURE 11: Percentage of Influenza Like-Illness Visits from ED and UCC Chief Complaints, ESSENCE-FL Participating Facilities, RDSTF Region 7 ESSENCE-FL Participating Facilities (N=52), Week 40, 2011 through Week 2, 2014 accessed January 21, 2015



**TABLE 2** shows the number of specimens tested by BPHL, how many are influenza positive and their subtypes.


**Table 2: Bureau of Public Health Laboratories (BPHL) Viral Surveillance for Week 2 by Lab Event Date\* as reported by 10:00 a.m. January 21, 2015**

	Current Week 2	Previous Week 1
<b>Total Specimens Tested</b>	<b>37</b>	<b>90</b>
Influenza positive specimens (% of total)	24 (64.9%)	45 (50.0%)
Influenza A (2009 H1N1) (% of influenza positives)	-	-
Influenza A (H3) (% of influenza positives)	13 (54.2%)	40 (88.9%)
Influenza A not yet subtyped (% of influenza positives)	9 (37.5%)	1 (2.2%)
Influenza B Yamagata (% of influenza positives)	-	2 (4.4%)
Influenza B Victoria (% of influenza positives)	-	1 (2.2%)
Influenza B not yet subtyped (% of influenza positives)	2 (8.3%)	1 (2.2%)


**FIGURE 12 - FIGURE 13** use BPHL viral surveillance data to track the progress of influenza infection over time. They include weekly information on how many specimens are tested by the BPHL, what proportion of those test positive for influenza and what subtypes are identified.

- **Influenza A (H3) and influenza B have been identified by BPHL this season.**
- **In recent weeks, influenza specimens submitted to BPHL tested positive for influenza A (H3), influenza B Yamagata lineage and influenza B Victoria lineage.**
- Influenza A (H3) has been the most common strain of influenza detected by BPHL so far in the 2014-2015 influenza season.
- The drifted Influenza A (H3) strain has been detected in Florida.

**FIGURE 12: Number of Influenza-Positive Specimens Tested by the Florida Bureau of Public Health Laboratories (BPHL) by Subtype by Lab Event Date\*, Week 1, 2012 to Week 2, 2015 as Accessed in Merlin by 10:00 a.m. January 21, 2015**



**FIGURE 13: Number of Specimens Tested by Florida Bureau of Public Health Laboratories (BPHL) and Percent Positive for Influenza by Lab Event Date\* Week 1, 2012 to Week 2, 2015 as Accessed in Merlin by 10:00 a.m. January 21, 2015**



\*Please note that lab event date is defined as the earliest of the following dates associated with the lab: date specimen collected, date received by the laboratory, date reported or date inserted.

For county-specific laboratory data, please refer to the Flu Lab Report in Merlin.

For instructions on how to use the Flu Lab Report, please see the Guide to Flu Lab Report on the Bureau of Epidemiology website:


<http://www.floridahealth.gov/diseases-and-conditions/influenza/documents/flulabreportguide.pdf>

As of 11:30 a.m. January 21, 2015, a total of 67 (100%) counties reported their weekly level of influenza activity. *Please note that data reported by counties after the deadline Tuesday at 5 p.m. are recorded but may not be included in the activity map for previous weeks.*


**TABLE 3: Weekly County Influenza Activity for Week 2 (ending January 17, 2015) as Reported by 11:30 a.m. January 21, 2015**

Activity Level	Week 2 Number of Counties	Week 1 Number of Counties	Week 2 Counties
No Report	0	0	-
No Activity	6	5	Bay, Desoto, Dixie, Hamilton, Leon, Taylor
Mild	45	47	Alachua, Bradford, Broward, Calhoun, Citrus, Clay, Collier, Columbia, Dade, Duval, Escambia, Flagler, Franklin, Gadsden, Gilchrist, Gulf, Hendry, Hernando, Highlands, Indian River, Jackson, Jefferson, Lafayette, Lake, Levy, Madison, Manatee, Martin, Monroe, Nassau, Okaloosa, Osceola, Palm Beach, Pasco, Pinellas, Santa Rosa, Sarasota, Seminole, St. Johns, Sumter, Suwannee, Union, Wakulla, Walton, Washington
Moderate	15	15	Baker, Brevard, Charlotte, Glades, Hardee, Hillsborough, Holmes, Lee, Marion, Okeechobee, Orange, Polk, Putnam, St. Lucie, Volusia
Widespread	1	0	Liberty

**Map 4: Weekly County Influenza Activity Level for Week 2 Reported by 11:30 a.m. January 21, 2015**




**Map 5: Weekly County Influenza Activity Trend for Week 2 Reported by 11:30 a.m. January 21, 2015**



County influenza activity data are reported to BOE through EpiGateway on a weekly basis by each county influenza coordinator. Specific information is requested about laboratory results, outbreak reports and surveillance system activity. Figures 14-23, displayed below, reflect a county's assessment of influenza activity within their county as a whole as well as influenza activity within specific settings. For week 2, 21 counties indicated that activity was decreasing, 33 indicated activity was about the same as previous weeks and 13 indicated that activity was increasing.

**FIGURE 14: Assessment of Overall Influenza Activity Trend**



**FIGURE 14** shows the assessment of the overall influenza activity trend in each county as reported by CHD influenza coordinators for week 2 as of 11:30 a.m. January 21, 2015.



Counties are asked to evaluate influenza activity in certain settings within their county. Each setting has a scale for activity that ranges from none or minimal activity to very high activity. What defines each of the values varies by facility type, but the example of the assessment in elementary, middle and high schools is included below. More detailed information on the meanings of the levels for each setting can be found on the webpage also included below.

No or very minimal activity -- Scattered cases of ILI with no increase in absenteeism or disruption of school activities.


Moderate activity -- Absenteeism elevated above baseline (in range of 10 to 25%) in some but fewer than half of schools where it is known; occasional children sent home because of ILI.

High activity -- Absenteeism elevated above baseline (in range of 10 to 25%) in more than half of schools; most schools sending several or many children home each day because of ILI.


Very high activity -- Absenteeism high enough to force curtailment of some or all school activities.

**FIGURE 15 - FIGURE 18** show the activity levels in various facilities by county as reported by CHD influenza coordinators for week 2 as of 11:30 a.m., January 21, 2015.


**FIGURE 15: Assessment of Influenza Activity in Elementary, Middle, and High Schools**




**FIGURE 16: Assessment of Influenza Activity in Colleges and Universities**



**FIGURE 17: Assessment of Influenza Activity in Jails/Prisons**




**FIGURE 18: Assessment of Influenza Activity in Retirement Facilities**




**FIGURE 19 - FIGURE 23** show the activity levels in various facilities by county as reported by CHD influenza coordinators for week 2 as of 11:30 a.m., January 21, 2015.


**FIGURE 19: Assessment of Influenza Activity in Nursing Homes**




**FIGURE 20: Assessment of Influenza Activity in Healthcare Facilities**




**FIGURE 21: Assessment of Influenza Activity in Day Care Centers**



**FIGURE 22: Assessment of Influenza Activity in Businesses**



**FIGURE 23: Assessment of Influenza Activity in State or Local Government Offices**



## Pediatric Influenza-Associated Mortality

No influenza-associated pediatric deaths were reported in week 2.

Three influenza-associated pediatric deaths have been reported so far in the 2014-15 influenza season.




FDOH Bureau of Vital Statistics and county health departments (CHDs) collect death record data electronically in all 67 Florida counties, which can be accessed using ESSENCE-FL. For pneumonia and influenza (P&I) surveillance, death record literals are queried in ESSENCE-FL using a free-text query that searches for references to P&I on death certificates. Any mention of P&I in the death certificate literals, with certain exceptions, is counted as a P&I death. Current season P&I death numbers are preliminary estimates, and may change as more data are received. *The most recent data available are displayed here. ESSENCE-FL vital statistics death records data are currently considered to be complete through week 1, 2015.*

**FIGURE 24** shows the count of preliminary estimated P&I deaths for all Florida counties, the number of deaths predicted using a multi-year regression model and the upper bound of the 95% confidence interval for this prediction.

- For week 1 (ending Jan 10, 2015):
- 259 preliminary estimated P&I deaths were reported.
  - Upper bound of 95% confidence interval for prediction: 262 deaths.
  - No excess deaths.


Figure 24: Vital Statistics Statewide Pneumonia and Influenza Deaths, Multi-Year Regression Model  
Week 17, 2000 - Week 1, 2015, Reported into ESSENCE-FL



**FIGURE 25** shows P&I deaths for all Florida counties, week 40, 2010 - week 2 2015, as reported into ESSENCE-FL. Deaths due to P&I are above levels seen during previous years at this time.

- As of week 2 (ending Jan 17, 2015):
- 3,373 P&I deaths have been reported so far in the 2014-15 influenza season.

FIGURE 25: Vital Statistics Statewide Pneumonia and Influenza Deaths, Reported into ESSENCE-FL,  
Week 40, 2011-Week 2, 2015




**FIGURE 26** shows P&I deaths for all Florida counties by age group, week 40, 2012 - week 2, 2015, as reported into ESSENCE-FL.

Increases in the preliminary number of P&I deaths are expected for this time of year.

**Seasons where influenza A (H3N2) is the predominantly circulating strain are associated with higher mortality and morbidity, particularly in the over 65 age group.**

It is likely that flu deaths will reach higher levels later this season since mortality tends to lag behind other indicators.


FIGURE 26: Vital Statistics Florida Pneumonia and Influenza Deaths by Age Group, Reported into ESSENCE-FL,  
Week 40, 2012 through Week 2, 2015



\* Death records data reported into ESSENCE-FL are currently considered to be complete through week 1, 2015.

The National Respiratory and Enteric Virus Surveillance System (NREVSS) collects data from laboratories around the country on a weekly basis. NREVSS monitors temporal and geographic patterns of six common respiratory viruses. Eight Florida facilities reported in week 2.

**FIGURE 27** shows the percentage of positive tests for multiple respiratory viruses reported by NREVSS-participating laboratories in Florida.



The 6 respiratory viruses summarized in Figure 27 are:

- Respiratory Syncytial Virus (RSV)
- Parainfluenza 1-3
- Adenovirus
- Human Metapneumovirus (HMPV)
- Rhinovirus
- Influenza

## Influenza and ILI Outbreaks Reported in EpiCom

In week 2, 2014, nine outbreaks of influenza or ILI were reported in to EpiCom.

### Hillsborough County

- A school reported 14 children with ILI. Four specimens tested positive for influenza A by rapid antigen test at local healthcare providers. Infection control measures were reviewed with facility leadership. Investigation is ongoing.
- An assisted living facility reported 15 residents with ILI. Two residents were hospitalized. Two specimens tested positive for influenza A by rapid antigen test at local healthcare providers. Fourteen of the fifteen ill residents (93%) had received the 2014-15 vaccine. Chemoprophylaxis was recommended for all residents. Investigation is ongoing.
- A school reported 15 students with ILI. Eight specimens tested positive for influenza A by rapid antigen test at local healthcare providers. Infection control measures were reviewed with facility leadership. Investigation is ongoing.
- An assisted living facility reported 29 residents and two staff with ILI. Two residents were hospitalized. Two specimens tested positive for influenza A by rapid antigen test at local healthcare providers. The 2014-15 vaccination rate for residents and staff was 90% and 75%, respectively. Chemoprophylaxis was recommended for all residents. Investigation is ongoing.

### Liberty County

- A correctional facility reported 68 inmates with ILI. At a local healthcare provider, three specimens tested positive for influenza A, two specimens tested positive for influenza B, and two specimens were negative for influenza. Chemoprophylaxis was recommended. Investigation is ongoing.

### Pinellas County

- A long term care facility reported eight residents and one staff with ILI. Eight additional residents reported minor respiratory symptoms. Four residents tested positive for influenza A by rapid antigen test at local providers. Chemoprophylaxis was recommended at the facility. All four residents that tested positive for influenza A had received the 2014-15 vaccine. Infection control measures were reviewed with facility leadership. Investigation is ongoing.

### Polk County

- A nursing facility reported 18 residents with ILI. One resident was hospitalized. Two specimens tested positive for influenza A by rapid antigen test at local providers. Chemoprophylaxis was recommended for all residents. The 2014-15 vaccination rate for residents was about 100%. Infection control measures were reviewed with facility leadership. Investigation is ongoing.

### Sarasota County

- A nursing facility reported eight residents and three staff with ILI. Two specimens tested positive for influenza A by rapid antigen test at local healthcare providers; one being collected from a hospitalized resident. Chemoprophylaxis was recommended for all residents and staff. Eight of nine ill residents (88%) and none of the ill staff had received the 2014-15 vaccine. Infection control measures were reviewed with facility leadership. Investigation is ongoing.
- A nursing facility reported six residents and four staff with ILI. One specimen tested positive for influenza A at local healthcare providers and five samples tested negative for influenza. Five of six (75%) ill residents and three out of four (75%) ill staff had received the 2014-15 vaccine. Chemoprophylaxis was recommended for all residents and staff. Investigation is ongoing.

**Seventy-one** outbreaks of influenza or ILI have been reported into EpiCom so far in the 2014-2015 season.

Map 6: Influenza and ILI outbreaks by county, Week 2, 2015 as Reported by 11:30 a.m., January 21, 2015




TABLE 4 : Summary of Florida Influenza and ILI Outbreaks by Facility Status, Week 40-2, 2015

Setting	Number of outbreaks	Implicated Viruses
Schools	13	<ul style="list-style-type: none"> <li>• One outbreak due to influenza (H3)</li> <li>• Seven outbreaks due to influenza A unspecified</li> <li>• Two outbreaks due to influenza A unspecified and influenza B unspecified</li> <li>• Three outbreaks, virus information not yet available</li> </ul>
Daycares	9	<ul style="list-style-type: none"> <li>• Three outbreaks due to influenza (H3)</li> <li>• Three outbreaks due to influenza A unspecified</li> <li>• One outbreak due to influenza A unspecified and influenza B unspecified</li> <li>• One outbreak due to RSV</li> <li>• One outbreak due to parainfluenza III</li> </ul>
Jails and prisons	1	<ul style="list-style-type: none"> <li>• One outbreak due to influenza A unspecified and influenza B unspecified</li> </ul>
Nursing homes and long term care facilities	47	<ul style="list-style-type: none"> <li>• Seven outbreaks due to influenza A (H3)</li> <li>• Twenty-nine outbreaks due to influenza A unspecified</li> <li>• Two outbreaks due to influenza A unspecified and influenza B unspecified</li> <li>• One outbreak due to influenza A (H3) and rhinovirus</li> <li>• One outbreak due to influenza A unspecified and RSV</li> <li>• One outbreak due to influenza A (H1N1)</li> <li>• One outbreak due to influenza B Yamagata</li> <li>• One outbreak due to influenza B unspecified</li> <li>• One outbreak due to RSV</li> <li>• Four outbreaks, virus information not yet available</li> </ul>
Colleges and universities, private businesses, local and state government offices, retirement homes, healthcare facilities, other	0	<ul style="list-style-type: none"> <li>• No outbreaks</li> </ul>
<b>Total</b>	<b>71</b>	<ul style="list-style-type: none"> <li>• Eleven outbreaks due to influenza A (H3)</li> <li>• Thirty-eight outbreaks due to influenza A unspecified</li> <li>• Five outbreaks due to influenza A unspecified and influenza B unspecified</li> <li>• One outbreak due to influenza A unspecified and RSV</li> <li>• One outbreak due to influenza B Yamagata</li> <li>• One outbreak due to influenza B unspecified</li> <li>• One outbreak due to H1N1</li> <li>• Two outbreaks due to RSV</li> <li>• One outbreak due to parainfluenza III</li> <li>• Seven outbreaks, virus information not yet available</li> </ul>

## Florida ILI Surveillance System Summary

### Florida ILINet

#### *Measures trends in ILI visits to outpatient doctor's offices*

Network of volunteer healthcare providers who:  
 Report ILI and total visit counts every week  
 Submit specimens for confirmatory testing

### ESSENCE-FL Syndromic Surveillance

#### *Measures trends in ILI visits and hospital admissions from emergency departments and urgent care clinics*

EDs and UCCs electronically transmit visit data into ESSENCE-FL daily  
 Visit data summarized in the Florida Flu Review include:  
 Percent of ED/urgent care visits due to ILI  
 Percent of ED/urgent care visitors with ILI who are admitted to the hospital

### ESSENCE-FL Vital Statistics Portal

#### *Measures influenza mortality by using death certificates with pneumonia or influenza listed as a cause of death.*

Death certificate data from the Bureau of Vital Statistics can be accessed through ESSENCE-FL and are used for pneumonia and influenza mortality surveillance

### County Influenza Activity in EpiGateway

#### *Uses data provided by CHDs to create a county-by-county breakdown of influenza and ILI activity around the state*

CHD epidemiologists report their county's influenza and ILI surveillance data weekly into the EpiGateway website  
 Influenza activity is classified as: No Activity, Mild, Moderate or Widespread  
 Setting-specific influenza activity and influenza trend is also reported

### Outbreak Reporting in EpiCom

#### *Tracks influenza and ILI outbreak investigations by CHDs and shows what types of influenza are responsible for outbreaks and where outbreaks are occurring*

CHD epidemiologists report outbreaks of influenza or ILI into EpiCom, Florida's online disease communication system  
 Outbreaks are defined as two or more cases of influenza or ILI in a specific setting

### BPHL

BPHL performs confirmatory testing and subtyping on surveillance specimens from ILINet sentinel providers, outbreak investigations, patients with severe or unusual influenza presentations and medical examiners

### Case-Based Influenza Surveillance

#### **Pediatric Influenza-Associated Mortality**

Deaths in children with laboratory-confirmed influenza infection are reportable in Florida

#### **Influenza due to Novel or Pandemic Strains**

Patients with influenza infection due to novel or pandemic strains are reportable in Florida

### National Respiratory and Enteric Virus Surveillance System (NREVSS)

#### *Measures trends in different viruses that cause respiratory disease*

Network of laboratories who report counts of test results for common respiratory viruses, including influenza, RSV, rhinovirus and others